


***Efficacy and safety of antidotes
for acute poisoning by cyanides***

Volume II

Technical Report No. 121



***Efficacy and safety of antidotes
for acute poisoning by cyanides***

Volume II

Technical Report No. 121

Brussels, November 2013

ISSN-0773-8072-121 (print)
ISSN-2079-1526-121 (online)

ECETOC Technical Report No. 121

© Copyright – ECETOC AISBL

European Centre for Ecotoxicology and Toxicology of Chemicals
2 Avenue E. Van Nieuwenhuyse (Bte 8), B-1160 Brussels, Belgium.

All rights reserved. No part of this publication may be reproduced, copied, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise without the prior written permission of the copyright holder. Applications to reproduce, store, copy or translate should be made to the Secretary General. ECETOC welcomes such applications. Reference to the document, its title and summary may be copied or abstracted in data retrieval systems without subsequent reference.

The content of this document has been prepared and reviewed by experts on behalf of ECETOC with all possible care and from the available scientific information. It is provided for information only. ECETOC cannot accept any responsibility or liability and does not provide a warranty for any use or interpretation of the material contained in the publication.

Efficacy and Safety of Antidotes for Acute Poisoning by Cyanides

CONTENTS

1. SUMMARY AND CONCLUSIONS	1
2. INTRODUCTION	VOLUME I
2.1 Background	
2.2 Terms of reference	
2.3 Pathophysiology, pharmacokinetics and antidotal mode of action for cyanide	
3. METHODOLOGY	VOLUME I
3.1 Selection of experts	
3.2 Procedure	
3.3 Materials	
3.3.1 Criteria for exclusion	
3.4 Methods	
3.4.1 By antidote	
3.4.2 By poisoning circumstance	
3.4.3 Practicality	
3.4.4 Assessment of efficacy and effectiveness	
3.4.5 Additional cases for dicobalt edetate	
4. ANTIDOTES	VOLUME I
4.1 Oxygen	
4.1.1 Introduction	
4.1.2 Identity	
4.1.3 Practicality	
4.1.4 Pharmacokinetics and metabolism	
4.1.6 Safety	
4.1.7 Case series and casuistics of cyanide / cyanogenic poisonings treated with or without oxygen	
4.1.8 Assessment of efficacy and effectiveness in clinical use	
Comparing the successes of antidote therapy with and without oxygen application, no appreciable differences were seen:	
4.1.9 Gaps in knowledge	
4.1.10 Conclusion	
4.2 Sodium thiosulphate	
4.2.1 Introduction	
4.2.2 Identity	
4.2.3 Practicality	
4.2.4 Pharmacokinetics and metabolism	
4.2.5 Mode of action, dosage and time of onset	
4.2.6 Safety	
4.2.7 Case series and casuistics of cyanide / cyanogenic poisonings treated with STS	
4.2.8 Assessment of efficacy and effectiveness in clinical use	

4.2.9 Conclusion

4.3 Nitrites

4.3.1 Introduction

4.3.2 Identity

4.3.3 Physico-chemical and pharmaceutical properties

4.3.4 Practicality

4.3.5 Pharmacokinetics and metabolism

4.3.6 Mode of action, dosage and time of onset

4.3.7 Safety

4.3.8 Case series and casuistics of cyanide / cyanogenic poisonings treated with nitrites

4.3.9 Occurrence of adverse effects in clinical use

4.3.10 Assessment of efficacy and effectiveness in clinical use

4.3.11 Gaps in knowledge

4.3.12 Conclusion

4.4 4-Dimethylaminophenol

4.4.1 Introduction

4.4.2 Identity

4.4.3 Practicality

4.4.4 Mode of action, dosage and time of onset

4.4.5 Safety

4.4.6 Case series and casuistics of cyanide / cyanogenic poisonings treated with DMAP

4.4.7 Occurrence of adverse effects in clinical use

4.4.8 Assessment of efficacy and effectiveness clinical use

4.4.9 Gaps in knowledge

4.4.10 Conclusion

4.5 Dicobalt edetate

4.5.1 Introduction

4.5.2 Identity

4.5.3 Practicality

4.5.4 Pharmacokinetics and metabolism

4.5.5 Mode of action, dosage and time of onset

4.5.6 Safety

4.5.7 Case series and casuistics of cyanide / cyanogenic poisonings treated with Dicobalt-EDTA

4.5.8 Assessment of efficacy and effectiveness in clinical use

4.5.9 Conclusion

4.6 Hydroxocobalamin

4.6.1 Introduction

4.6.2 Identity

4.6.3 Physico-chemical and pharmaceutical properties

4.6.4 Practicality

4.6.5 Pharmacokinetics and metabolism

4.6.6 Mode of action, dosage and time of onset

4.6.7 Safety

4.6.8 Case series and casuistics of cyanide / cyanogenic poisonings treated with hydroxocobalamin

4.6.9 Occurrence of adverse effects in clinical use

4.6.10	Assessment of efficacy and effectiveness in clinical use	
4.6.11	Gaps in knowledge	
4.6.12	Conclusion	
4.7	Pre-clinical drugs	
4.7.1	Introduction	
4.7.2	3-Mercaptopyruvate and prodrugs	
4.7.3	α -Ketoglutarate	
4.7.4	Cysteine and N-Acetyl-cysteine	
4.7.5	Cobinamide	
4.7.6	Dihydroxyacetone	
4.7.7	Hemoglobin-glutamer	
4.7.8	Conclusion	
5.	POISONING CIRCUMSTANCES	VOLUME I
5.1	Direct poisoning (HCN and Cyanide salts)	
5.1.1	Introduction	
5.1.2	Identity	
5.1.3	Pharmacokinetics and toxicity	
5.1.4	Casuistics	
5.1.5	Treatment and/or prophylaxis	
5.1.6	Occurrence of adverse effects in clinical use	
5.1.7	Assessment of efficacy and effectiveness of antidotes in clinical use	
5.1.8	Discussion and conclusion	
5.2	Cyanogenic substances	
5.2.1	Cassava	
5.2.2	Amygdalin (laetrile)	
5.2.3	Nitriles	
5.2.4	Sodium nitroprusside	
5.3	Fire smoke (mixed intoxications)	
5.3.1	Toxicity - Cyanide concentrations	
5.3.2	Case series and casuistics	
5.3.3	Gaps in knowledge	
5.3.4	Conclusion	
5.4	Poisoning with an unknown substance	
5.4.1	Introduction	
5.4.2	Approach to treating a patient suspected of having had an overdose or having been poisoned by an unknown substance	
5.4.3	Symptomatology of Cyanide Poisoning	
5.4.4	Review of Case Reports	
5.4.5	Conclusion	
6.	CONCLUSIONS AND RECOMMENDATIONS	VOLUME I
6.1	Classification of antidotes to cyanide and mode of action	
6.2	Practicality	
6.2.2	STS Error! Bookmark not defined.	
6.2.3	Methaemoglobin-forming agents	

6.2.4	Co-EDTA	
6.2.5	HOCO	
6.3	Efficacy and effectiveness	
6.3.1	STS Error! Bookmark not defined.	
6.3.2	Amyl nitrite	
6.3.3	Sodium nitrite	
6.3.4	AN/SN/STS	
6.3.5	DMAP	
6.3.6	Co-EDTA	
6.3.7	HOCO	
6.4	Poisoning circumstances and antidotes	
6.4.1	Direct poisonings – HCN and salts	
6.4.2	Cyanogens	
6.4.3	Smoke	
6.4.4	Unknown poisonings	
6.5	Summary	
LIST OF ABBREVIATIONS		4
BIBLIOGRAPHY		6
APPENDIX A: QUESTIONNAIRE		141
A.1	Personal Cases	141
A.2	The questionnaire	143
B.	ECETOC questionnaire on poisoning cases with cyanides	144
20	Responses received (10-Aug-10)	145
APPENDIX B: DATABASE OF > 400 CASES		146
APPENDIX C: CASUISTICS		147
APPENDIX D: EFFECTS OF HOCO ON LABORATORY MEASUREMENTS AND PARAMETERS		246
MEMBERS OF THE TASK FORCE		248
MEMBERS OF THE SCIENTIFIC COMMITTEE		249

1. SUMMARY AND CONCLUSIONS

Cyanide poisoning is a very serious, albeit rare, event with possible neurological sequelae that may result in severe disability, and death. It can occur after exposure by ingestion, inhalation and dermal absorption of hydrocyanic acid, cyanide salts, or cyanogenic compounds such as acetone cyanohydrin, nitriles, biological substances like cassava or amygdalin (Vitamin B17), or sodium nitroprusside (SNP) (used as a medication for hypertension). Fire smoke inhalation may also cause cyanide poisoning under certain conditions of combustion.

Despite the 1993 overview on “Antidotes of Poisoning by Cyanide” (IPCS/CEC Evaluation of Antidotes Series) and the review of commonly used cyanide antidotes in the 2007 ECETOC Report in the JACC (Joint Assessment of Commodity Chemicals) series “Cyanides of Hydrogen, Sodium and Potassium, and Acetone Cyanohydrin” there has been no recognised consensus on the relative efficacy, efficiency, safety, and practicality of the various antidote treatments being used across the world. This situation although undesirable was sustained by the absence of any comprehensive review which was in turn due to the complexity of the clinical picture, since cyanide antidotes are employed in different circumstances of poisoning. In order to address this situation ECETOC established a new Task Force to review the available evidence on the efficacy, efficiency, safety, and practicality of antidote regimes under the different poisoning circumstances in which they were used. The four circumstances considered were direct poisonings with hydrocyanic acid or its salts, poisoning with cyanogenic compounds (above), fire smoke inhalation, and poisoning with initially unknown substances.

The Task Force reviewed the available literature until mid-2010 and collected further cases with a questionnaire from industry and Poison Control Centres. The Poisoning Severity Score (PSS), a well-established score system addressing all organ systems for grading of poisoning severity, was adjusted slightly to meet the specifics of cyanide poisoning (PSSa – ‘a’ denotes adapted).

The reported cases were then evaluated for antidote use. In the majority of cases single antidotes were given although in some combinations of antidotes were used. Cases were split into treatment groups based on the sequence in which different antidotes were administered, if more than one antidote was given. For each such sequence the efficacy based on the PSSa and the safety of the respective antidote or antidote combinations were described. This sequential view allowed in many cases for a statistical analysis of efficacy which was included in the single antidote assessments together with pharmaceutical data, pharmacokinetic data, safety/side effects, practicality, an overview of reported cases and case series. Single casuistic case reports shortly described in tables were collected in an appendix as summaries to make the antidote chapters more readable. An overall assessment was made for each antidote.

Pre-clinical antidotes were addressed based on data from animal experiments. In particular alpha-ketoglutarate and cobinamide, and to a lesser extent, dihydroxyacetone and pyruvate (pro-drugs) seem to have promising potential as cyanide antidotes.

It was apparent that although all direct acting antidotes (excluding sodium thiosulphate) may reverse cardiac arrest or at least facilitate resuscitation, they cannot prevent neurological sequelae due to hypoxia in all cases.

Sodium thiosulphate appears to act more rapidly than previously thought, but is effective, when administered alone, only in moderate poisoning (PSSa 2). Amyl nitrite, when administered alone, was found to be effective in moderate to severe poisonings (PSSa 2.5). Hydroxocobalamin, when administered alone, was found to be effective in severe poisonings (PSSa 3). The combinations of sodium nitrite and sodium thiosulphate, with or without amyl nitrite (PSSa 3), and of dimethylaminophenol and sodium thiosulphate (PSSa 2) were also found to be at least partially effective. These methaemoglobin forming agents all require combination with sodium thiosulphate to capture cyanide released during physiological methaemoglobin reduction. Dicobalt edetate has rarely been given, so that no comprehensive evaluation was possible. In any case, the intrinsic toxicity of dicobalt edetate significantly reduces its applicability. In regard of its effectiveness and low toxicity sodium thiosulphate can be given after all direct acting antidotes in situations when a delayed formation of cyanides cannot be excluded.

Oxygen, in contrast to the other antidotes discussed, has no effect on cyanide toxicokinetics, and is therefore not regarded as an antidote in its own right. Oxygen was only partially effective in severe poisonings (PSSa 3), and other antidotes were fully effective without oxygen. The administration of normobaric oxygen in combination with other antidotes to cyanide is very effective and possibly at least additive. It is however recommended that in conditions where oxygen is not available, treatment with antidotes to cyanide should not be delayed.

For the different circumstances of cyanide poisoning mechanism of cyanide formation, course, case series, case tables and an overview of antidote uses in the respective conditions were addressed, before recommending (a) specific antidote(s) for the condition. These recommendations could not only deal with efficacy and safety, but had also to account for practicility, e.g. cold storage requirement in a tropical country, or high price in developing countries. These combined aspects prompted recommendations for different situations / circumstances and severities shown in the following table.

Table 1: Antidotes^a recommended for acute poisoning by cyanides

Circumstance ^b	Severity of poisoning	
	Mild or moderate	Severe
Direct		
HCN or its salts	None or STS (HOCO)	AN/SN or DMAP, followed by STS
Cyanogenic compound		
Cassava	None or STS	STS, in very rare most severe cases SN followed by STS
Laetrile/Amygdalin	None or STS	AN/SN/STS or HOCO(/STS) (or DMAP/STS)
Nitriles	None or STS	STS, in very severe cases SN/STS or HOCO(/STS) or DMAP/STS
SNP	STS	STS
Smoke		
CO and cyanide	None or STS	HOCO (followed by STS). Neither AN/SN nor DMAP
Unknown	None or STS	HOCO or AN, followed by STS. If then required SN/STS (or DMAP/STS)
Child	None or STS	HOCO and/or STS
Mass poisoning	None	AN (or DMAP i.m.)
First aider	None	AN

^a STS, sodium thiosulphate; HOCO, hydroxocobalamin; AN, amyl nitrite; SN, sodium nitrite; DMAP, dimethylaminophenol; i.m., intramuscular (injection).

^b HCN, hydrogen cyanide; SNP, sodium nitroprusside; CO, carbon monoxide.

The basis of cyanide poisoning treatment, independent of the source, are the usual measures of life support i.e. oxygen, and if required mechanical ventilation and safeguarding of adequate circulation by e.g. catecholamines.

ABBREVIATIONS

-	Not applied, to (in range)
ACH	Acetone cyanohydrin
AN	Amyl nitrite
ATC	Anatomical Therapeutic Chemical Classification System (World Health Organisation)
bw	Bodyweight
CAS	Chemical Abstracts Service (American Chemical Society)
CO	Carbon monoxide
COHb	Carboxyhaemoglobin
Co-EDTA	Dicobalt edetate
d	Day
DMAP	Dimethylaminophenol
FiO ₂	Fraction of inspired oxygen
g	Gram
h	Hour(s)
h-	Hecto-
Hb	Haemoglobin
HCN	Hydrogen cyanide
HOCO	Hydroxocobalamin
i.m.	Intramuscular
i.o.	Intraosseous
i.p.	Intraperitoneal
i.v.	Intravenous
ICSC	International Chemical Safety Card (International Labour Organisation)
IPCS	International Programme on Chemical Safety
k-	Kilo-
KCN	Potassium cyanide
l	Litre
µ-	Micro-
m-	Milli-
MetHb	Methaemoglobin
NaCN	Sodium cyanide
NS	Not stated
Ph Eur	European Pharmacopoeia (Council of Europe)
p.o.	Per os (oral)
PSS	Poisoning Severity Score
PSSa	PSSAadapted to cyanide
s	Second
s.c.	Subcutaneous
SN	Sodium nitrite
SNP	Sodium nitroprusside

STS	Sodium thiosulphate
USP	United States Pharmacopeia
WHO	World Health Organisation

BIBLIOGRAPHY

~References quoted

The following references were consulted by the Task Force and quoted in the report.

Aaron CK. 1998. Cyanide antidotes. In Goldfrank LR, Flomenbaum N, Lewin NA, Weisman RS, Howland MA, Hoffman RS, eds, Goldfrank's Toxicological Emergencies, 6th ed. Appleton and Lange, Stamford, CT, USA, pp 1583-1585.

Ageeva, TS. 1970. The condition of the metabolism of mediating substances in workers producing acrylonitrile. Tr Sarat Med Inst 78:10-13 [Russian].

Aguilar R, Soy D, Nogué S. 2006. Availability of essential antidotes in Catalonia hospitals (Spain). Clin Toxicol 44:554-555.

Ahmed, AE; Loh, JP; Ghanayem, B; et al. (1992) Studies on the mechanism of acetonitrile toxicity: I. Whole body autoradiographic distribution and macromolecular interaction of ^{14}C -acetonitrile in mice. Pharmacol Toxicol 70:322-330.

Ahmed AE, Patel K. 1981. Acrylonitrile. *In vivo* metabolism in rats and mice. Drug Metabol Dispos 9:219-222.

Ahmed AE, Farooqui MYH. 1982. Comparative toxicities of aliphatic nitriles. Toxicol Lett 12:157-163.

Ahmed AE, Trieff NM. 1983. Aliphatic nitriles: Metabolism and toxicity. Progress in Drug Metabolism 7:229-294.

Ahmed AE, Soliman SA, Loh JP, Hussein GI. 1989. Studies on the mechanism of haloacetonitriles toxicity: Inhibition of rat hepatic glutathione S-transferases *in vitro*. Toxicol Appl Pharmacol. 100(2):271-279.

Aitken D, West D, Smith F, Poznanski W, Cowan J, Hurtig J, Peterson E, Benoit B. 1977. Cyanide toxicity following nitroprusside induced hypotension. Can Anaesth Soc J 24(6):651-660.

Akanji AO, Famuyiwa OO. 1993. The effects of chronic cassava consumption, cyanide intoxication and protein malnutrition on glucose tolerance in growing rats. Br J Nutr 69:269-276.

Akimenko VK. 1976. Detection of cyanide resistant respiration among Candida lipolytic yeasts. Mikrobiologiya 45:146-150 [Russian].

Akintonwa A, Tunwashe OL. 1992. Fatal cyanide poisoning from cassava-based meal. *Hum Exp Toxicol* 11:47-49.

Alarie Y. 2002. Toxicity of fire smoke. *Crit Rev Toxicol* 32:259-289.

Albaum, HG; Tepperman, J; Bodansky, O. (1946) The in vivo inactivation by cyanide of cytochrome oxidase and its effects on high energy phosphorous compounds in the brain. *J Biol Chem* 64:45-51.

Aldous C, Norris J, Baltar A, Wilson R, Ho I, Hume A. 1984. Formation of cyanohydrin and antagonism of cyanide by carbonyl compounds. *Pharmacologist* 26:230.

Alston RE, Turner BL. 1963. Cyanogenic substances. In Alston RE, Turner BL, eds, *Biochemical systematics*. Prentice-Hall, Englewood Cliffs, New Jersey, USA, pp 181-190.

Aly FW. 1957. Behandlung einer schweren Cyannatriumvergiftung mit Methylenblau und Katalysin. *Arztl Wochensh* 12:1014-1018.

Amdur ML. 1959. Accidental group exposure to acetonitrile: A clinical study. *J Occup Med* 1:627-633.

Anderson RA, Thomson I, Harland WA. 1979. The importance of cyanide and organic nitriles in fire fatalities. *Fire Mater* 3:91-99.

Anonymous. 1972. Food and cyanide. *Br Med J* 2:481-482.

Anonymous. 1987. Intoxication par les cyanures. *Contact et Urgences* 29:12-15.

Anonymous. 2001. Acetonitril. In: H.Greim, Ed, *Gesundheitsschädliche Arbeitsstoffe, Toxikologisch-arbeitsmedizinische Begründungen von MAK-Werten*. Wiley-VCH, München, Germany.

Anonymous. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 0 cyanide/nitrile cases]. Personal communication. Laboratory for Clinical and Forensic Toxicology and Drug Analyses, University Hospital, Groningen, The Netherlands.

Ansell M, Lewis FAS. 1970. A review of cyanide concentrations found in human organs. A survey of literature concerning cyanide metabolism, 'normal', non-fatal, and fatal body cyanide levels. *J Forensic Med* 17:148-155.

Appel KE, Bolt M. 1981. Effect of antidotes in acute acrylonitrile poisoning. *Naunyn-Schmiedeberg's Arch Pharmacol* 316:Suppl, R 15.

Appel KE, Peter H, Bolt HM. 1981. Effect of potential antidotes on the acute toxicity of acrylonitrile. *Int Arch Occup Environ Health* 49:157-163.

Ardelt BK, Borowitz JL, Isom GE. 1989. Brain lipid peroxidation and antioxidant protectant mechanisms following acute cyanide intoxication. *Toxicology* 56:147-154.

Ardelt BK, Borowitz JL, Maduh EU, Swain SL, Isom GE. 1994. Cyanide-induced lipid peroxidation in different organs: subcellular distribution and hydroperoxide generation in neuronal cells. *Toxicology* 89:127-137.

Aregheore EM, Agunbiade OO. 1991. The toxic effects of cassava (*Manihot esculenta* Grantz) diets on humans: a review. *Vet Hum Toxicol* 33:274-275.

Arieff AI. 1992. Is measurement of venous oxygen saturation useful in the diagnosis of cyanide poisoning. *Am J Med* 93:582-583.

Arnone G. 1952. Contributo alla sindrome di avvelenamento da acido cianidrico [Syndrome of hydrocyanic acid poisoning]. *Sicil Sanit* 5:711-714.

Arron GP, Spalding MH, Edwards GE. 1979. Isolation and oxidative properties of intact mitochondria from the leaves of *Sedum praealtum*: A crassulacean acid metabolism plant. *Plant Physiol* 64:182-186.

Astier A, Baud FJ. 1995. Simultaneous determination of hydroxocobalamin and its cyanide complex cyanocobalamin in human plasma by high-performance liquid chromatography: Application to pharmacokinetic studies after high-dose hydroxocobalamin as an antidote for severe cyanide poisoning. *J Chromatogr B* 667:129-135.

Astier A, Baud FJ. 1996. Complexation of intracellular cyanide by hydroxocobalamin using a human cellular model. *Hum Exp Toxicol* 15:19-25.

Atkinson A, Rutter DA, Sargeant K. 1974. Enzyme antidote for experimental cyanide poisoning. *Lancet* 2:1446.

ATSDR. 1990. Acrylonitrile. Agency for Toxic Substances and Disease Registry, US Department of Health and Human Services, Public Health Service, Atlanta, Georgia, USA. www.atsdr.cdc.gov/toxprofiles/tp125.pdf

ATSDR. 1993. Cyanide toxicity. *American Family Physician* 48:107-114. Agency for Toxic Substances and Disease Registry, US Department of Health and Human Services, Public Health Service, Atlanta, Georgia, USA.

ATSDR. 1997. Medical management guidelines for acrylonitrile. Agency for Toxic Substances and Disease Registry, US Department of Health and Human Services, Public Health Service, Atlanta, Georgia, USA. www.atsdr.cdc.gov/mhmi/mmg125.html

Attucci S, Carde J P, Raymond P, Saint-Ges V, Spiteri A, Pradet A. 1991. Oxidative phosphorylation by mitochondria extracted from dry sunflower seeds. *Plant Physiol* 95:390-398.

Auzepy P, Veissieres JF, Deparis M. 1974. Letter: Anaphylactic shock due to hydroxocobalamine. *Nouv Presse Med* 19:3-152 [French].

Aw TC, Bishop CM. 1981. Cyanide poisoning. *J Soc Occup Med* 31:173-174.

Ayanru JO. 1976. The tropical amblyopia syndrome (or tropical nutritional amblyopia) in the Mid-Western State of Nigeria. *Afr J Med Med Sci* 5:41-48.

Bader M, Wrbitzky R. 2006. Follow-up biomonitoring after accidental exposure to acrylonitrile: Implications for protein adducts as a dose monitor for short-term exposures. *Toxicol Lett* 162(2-3):125-131.

Bain JTB, Knowles EL. 1967. Successful treatment of cyanide poisoning. *Br Med J* 2:763.

Bakker JG, Jongens SMJ, van Neer FCJ, Neis JM. 1991. Occupational contact dermatitis due to acrylonitrile. *Contact Dermatitis* 24:50-53.

Balahura RJ, Wilkins RG. 1983. Ligational effects on reduction of myoglobin and horseradish peroxidase by inorganic reagents. *Biochim Biophys Acta* 724:465-72.

Ballantyne B. 1975. Blood, brain and cerebrospinal fluid cyanide concentrations in experimental acute cyanide poisoning. *J Forensic Sci Soc* 15:51-56.

Ballantyne B. 1976. Changes in blood cyanide as a function of storage time and temperature. *J Forensic Sci Soc* 16:305-310.

Ballantyne B. 1977a. Factors in the analysis of whole blood thiocyanate. *Clin Toxicol* 11:195-210.

Ballantyne B. 1977b. *In vitro* production of cyanide in normal human blood and the influence of thiocyanate and storage temperature. *Clin Toxicol* 11:173-193.

Ballantyne B. 1983. Artifacts in the definition of toxicity by cyanides and cyanogens. *Fundam Appl Toxicol* 3:400-408.

Ballantyne B. 1987. Toxicology of cyanides. In Ballantyne B, Marrs T, eds, Clinical and experimental toxicology of cyanides. John Wright, Bristol, UK, pp 41-126.

Ballantyne B, Marrs TC. 1987. Postmortem features and criteria for the diagnosis of acute lethal cyanide poisoning. In Ballantyne B, Marrs T, eds, Clinical and experimental toxicology of cyanides. John Wright, Bristol, UK, pp 217-247.

Ballantyne B, Bright J, Williams P. 1973. An experimental assessment of decreases in measurable cyanide levels in biological fluids. J Forens Sci Soc 13:111-117.

Ballantyne B, Bright JE, Williams P. 1974. The post-mortem rate of transformation of cyanide. Forensic Sci 3:71-76.

Ballesteros García MM, Fernández Carrión F, Casanueva Mateos L, Sánchez Díaz JJ, Gómez de Quero P, Ramos Casado V, Mar Molinero F. 2003. Intoxicación por nitroprusiato. An Pediatr 58:81-82.

Baranowski Z. 1985. Alternative pathway of respiration in *Physarum polycephalum* plasmodia. Cell Biol Int Rep 9:85-90.

Baranowski Z, Shraideh Z, Korohoda W, Wohlfarth-Bottermann KE. 1983. Consequences of impeding in mitochondrial function in *Physarum polycephalum*. II: Influence on 'de novo' generation of contractile activities and responses to glucose and blue light. Eur J Cell Biol 31:187-196.

Barcroft J. 1931. The toxicity of atmospheres containing hydrocyanic acid gas. J Hyg (Lond) 31:1-34.

Barillo DJ, Goode R, Esch V. 1994. Cyanide poisoning in victims of fire: Analysis of 364 cases and review of the literature. J Burn Care Rehabil 15:46-57.

Barillo DJ. 2009. Diagnosis and treatment of cyanide toxicity. J Burn Care Res 30:148-152.

Barker MH, Lindberg HA, Wald MH. 1941. Further experiences with thiocyanates. J Am Med Assoc 117:1591-1594.

Barker MH. 1936. The blood cyanates in the treatment of hypertension. J Am Med Assoc 106:762-767.

Bartelheimer E, Friedberg K, Lendle L. 1962. Chemische Reaktionsmöglichkeiten von Kobaltverbindungen als Voraussetzung ihrer Toxizität und Antidotwirksamkeit gegen Blausäure. Arch Int Pharmacodyn 139:99-108.

BASF, Hampshire Chemicals, ICI. 1993. Code of practice for chemicals with major hazards. Hydrogen cyanide; properties, uses, storage and handling. The safe design, construction and use of plants producing or consuming hydrogen cyanide. Unpublished report. BASF, Hampshire Chemicals, ICI, UK.

BASF. 1999. Safety data sheet according to 91/155/EEC, Acrylnitril ST 2. BASF, Ludwigshafen, Germany.

BASF. 2000. Internal safety data sheet, Cyanwasserstoff roh ber.HCN 100%. BASF, Ludwigshafen, Germany.

BASF. 2000. Safety data sheet according to 91/155/EEC, Cyannatrium solution ca. 30%. BASF, Ludwigshafen, Germany.

BASF. 2001. Thiocyanate in urine, metabolite of hydrocyanic acid (HCN). Personal communication by Will W, Occupational Medical and Health Protection Department. BASF, Ludwigshafen, Germany.

Baskin SI, Brewer TG. 1997. Cyanide poisoning. In Sidell FR, Takafuji ET, Franz DR, eds, Medical aspects of chemical and biological warfare. Part I: Textbook of military medicine. US Department of the Army, Office of the Surgeon General, Borden Institute, Washington, DC, USA, pp 271-286.

Baskin SI, Froehlich HL, Groff WA. 1986. The dissociation of reversal of cyanide (CN) toxicity and methemoglobin formation by nitrite (N) in the isolated heart. Fed Proc 45:196.

Baskin SI, Horowitz AM, Neally EW. 1992. The antidotal action of sodium nitrite and sodium thiosulfate against cyanide poisoning. J Clin Pharmacol 32:368-375.

Bassani RA, Fagian MM, Bassani JW, Vercesi AE. 1998. Changes in calcium uptake rate by rat cardiac mitochondria during postnatal development. J Mol Cell Cardiol 30:2013-2023.

Bastigkeit M. 2000. Blausäureintoxikation: Neues Antidot macht die Therapie verträglich – aber teuer. Rettungsdienst, Sonderdruck 1-4, Orphan Europe.

Bastigkeit M. 2002. Weg mit alten pharmakologischen Zöpfen - Ein (sehr) kritischer Überblick. Rettungsdienst 25:38-43.

Bastigkeit M. 2006. Antidote: Welche sind sinnvoll für den Rettungsdienst ? Rettungsdienst 26:1092-1095

Baud FJ. 1999. Personal communication, Cyanide Congress (Orphan Europe), Fulda.

Baud FJ. 2004/5. Dicobalt edetate. [French/English translation].

Baud FJ. 2007. Clarifications regarding interference of hydroxocobalamin with carboxyhemoglobin measurements in victims of smoke inhalation. *Ann Emerg Med* 50:625-626 [Comment on Lee et al, 2007; author's reply 626.]

Baud FJ. 2007. Cyanide: Critical issues in diagnosis and treatment. *Hum Exp Toxicol* 26:191-201.

Baud FJ. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 14 cyanide cases, June 2009]. Personal communication. Lariboisière Hospital, Paris, France.

Baud FJ, Borron SW. 2008. In reply: Hydroxocobalamin for acute cyanide poisoning in smoke inhalation. *Ann Emerg Med* 51:108-109.

Baud FJ, Lapostolle F. 2009. Matched comparison of plasma lactate concentrations in cyanide poisoned and drug poisoned patients (18). Presented at EAPCCT congress, May 2009, Stockholm, Sweden. [EAPCCT.abstracts.]

Baud FJ, Mégarbane B. 2009. Does the toxidrome of acute cyanide poisoning exist? (199). Presented at EAPCCT congress, May 2009, Stockholm, Sweden. [EAPCCT.abstracts]

Baud FJ, Bismuth C, Astier A, Djeghout H, Aubriot D. 1986. Toxicocinétique des cyanures et des thiocyanates lors d'un accident professionnel au propionitrile. *Arch Mal Prof* 47:85-86.

Baud FJ, Astier A, Toffis V, Barriot P. 1987. Plasma kinetics of hydroxocobalamin and cyanocobalamin in a treated cyanide poisoning. Unpublished data. Hôpital Fernand Widal, Paris, France.

Baud FJ, Toffis V, Certain B, Muzynski J, Astier A, Bismuth C. 1989. Hydroxocobalamine antidote des cyanures. Vérification cinétique de sa transformation en cyanocobalamine dans l'organisme intoxiqué. *Thérapie* 44:469.

Baud FJ, Barriot P, Toffis V, Astier A, Ledantec P, Briou HJ, Bismuth C. 1990. Démonstration de l'effet complexant de l'hydroxocobalamine chez l'intoxiqué par cyanure. *J Toxicol Clin Exp* 10:193-194.

Baud FJ, Barriot P, Toffis V, Riou B, Vicaut E, Lecarpentier Y, Bourdon R, Astier A, Bismuth C. 1991. Elevated blood cyanide concentrations in victims of smoke inhalation. *N Engl J Med* 325:1761-1766.

Baud FJ, Borron SW, Bavoux E, Astier A, Hoffman JR. 1996. Relation between plasma lactate and blood cyanide concentrations in acute cyanide poisoning. *BMJ* 312: 26-27 [with Cook TM. 1996. Plasma lactate and blood cyanid in acute cyanide poisoning. Letter to the editor and authors' reply. *BMJ* 312:1039].

Baud FJ, Borron SW, Mégarbane B, Trout H, Lapostolle F, Vicaut E, Debray M, Bismuth C. 2002. The value of lactic acidosis in the assessment of the severity of acute cyanide poisoning. *Crit Care Med* 30:2044-2050.

Beamer WC, Shealy RM, Prough DS. 1983. Acute cyanide poisoning from laetrile ingestion. *Ann Emerg Med* 12:449-451.

Beasley D.M.G, Glass WI. 1998. Cyanide poisoning pathophysiology and treatment recommendations. *Occup Med* 48:427-431.

Beasley RWR, Blown RJ, Lunau FW, Taylor BM. 1978. Amyl nitrite and all that. *J Soc Occup Med* 28:142-143.

Beavis AD, Vercesi A E. 1992. Anion uniport in plant mitochondria is mediated by a Mg(2+)-insensitive inner membrane anion channel. *J Biol Chem* 267:3079-3087.

Bebarta VS, Little C, Fragoso M, Heard K. 2007. A swine model for poison-induced cardiac arrest using intravenous potassium cyanide. *Clin Toxicol* 45:606.

Becker CE. 1985. The role of cyanide in fires. *Vet Hum Toxicol* 27:487-490.

Bell J. 1972. The acute toxicity of four common poisons to the opossum, *Trichosurus vulpecula*. *N Z Vet J* 20:212-214.

Benabid ML, Decorps M, Merry C. 1987. ³¹P nuclear magnetic resonance *in vitro* spectroscopy of metabolic changes induced in the awake rat brain during KCN intoxication and its reversal by hydroxocobalamin. *J Neurochem* 48:804-808.

Benaissa ML, Hantson P, Bismuth C, Baud FJ. 1995. Mercury oxycyanide and mercuric cyanide poisoning: Two cases. *Intensive Care Med* 21:1051-1053.

Benaissa L, Hantson P, Laforge M, Borron S, Baud F. 1999. Cyanides and toxic cyanogens. *Encyclopédie médico-chirurgicale. Toxicologie-Pathologie professionnelle*, 3rd quarter, no.124, pp 7 [in French].

Benz FW and Nerland DE. 2005. Effect of cytochrome P450 inhibitors and anticonvulsants on the acute toxicity of acrylonitrile. *Arch Toxicol* 79 : 610-614.

Benz FW, Nerland DE, Corbett D, Li J. 1997a. Biological markers of acute acrylonitrile intoxication in rats as a function of dose and time. *Fundam Appl Toxicol* 36:141-148.

Benz FW, Nerland DE, Li J, Corbett D. 1997b. Dose dependence of covalent binding of acrylonitrile to tissue protein and globin in rats. *Fundam Appl Toxicol* 36:149-156.

Benz FW, Nerland DE, Pierce WM, Babiuk C. 1990. Acute acrylonitrile toxicity: Studies on the mechanism of antidotal effect of D- and L-cysteine and their N-acetyl derivatives in the rat. *Toxicol Appl Pharmacol* 102:142-150.

Benz FW, Pierce WM, Nerland DE, Cai J. 2001. Covalent binding of acrylonitrile to specific liver glutathione S transferases *in vivo*. *Chem Res Toxicol* 14:799-806.

Benz FW. 1998. Acute acrylonitrile intoxication – Antidotal assessment. Crisp Data Base, National Institutes of Health, USA.

Berlin CM. 1970. The treatment of cyanide poisoning in children. *Pediatrics* 46:793-796.

Berninger TA, von Meyer L, Siess E, Schon O, Goebel FD. 1989. Leber's hereditary optic atrophy: Further evidence for a defect of cyanide metabolism. *Br J Ophthalmol* 73:314-316.

Bernt AS, Kerde C, Prokop O. 1962. Zur Frage der Verwertbarkeit von Cyanidbefunden im Leichenmaterial. *Dtsch Z Ges Gericht Med* 51:522.

Berumen U. 1983. Dog poisons man. *JAMA* 249:353.

Beutler B, Baluda M. 1963. Methemoglobin reduction: Studies on the interaction between cell populations and of the role of methylene blue. *Blood* 22:323 [Mechanism of action against methemoglobonaemia and side effects].

Bhattacharya R, Tulsawani R. 2009. Protective role of alpha-ketoglutarate against massive doses of cyanide in rats. *J Environ Biol* 30:515-520.

Bhattacharya R, Jeevaratnam K, Raza SK, Das Gupta S. 1991. Cyanide antagonism in a rodent model. *Arch Toxicol* 14:231-235.

Bhattacharya R, Jeevaratnam K, Raza SK, Das Gupta S. 1993. Protection against cyanide poisoning by the co-administration of sodium nitrite and hydroxylamine in rats. *Hum Exp Toxicol* 12:33-36.

Bhattacharya R, Kumar D, Sugendran K, Pant SC, Tulsawani RK, Vijayaraghavan R. 2001. Acute toxicity studies for α ketoglutarate: A promising antidote for cyanide poisoning. *J Appl Toxicol* 2:495-499.

Bhattacharya R, Lakshamana Rao PV, Parida MM, Jana AM. 1999. Antidotal efficacy of antioxidants against cyanide poisoning *in vitro*. Def Sci J 49:55-63.

Bhattacharya R, Lakshamana Rao PV. 2001. Pharmacological interventions of cyanide-induced cytotoxicity and DNA damage in isolated rat thymocytes and their protective efficacy *in vivo*. Toxicol Lett 199:59-70.

Bhattacharya R, Lakshmana Rao PV, Vijayaraghavan R. 2002. *In vitro* and *in vivo* attenuation of experimental cyanide poisoning by α -ketoglutarate. Toxicol Lett 128:185-195.

Bhattacharya R, Pant SC, Kumar D, Dube SN. 1995. Toxicity evaluation of two treatment regimens for cyanide poisoning. J Appl Toxicol 15:439-441.

Bhattacharya R, Rao PVL. 1997. Cyanide induced DNA fragmentation in mammalian cell cultures. Toxicology 123:207-215.

Bhattacharya R, Tulsawani R, Kumar D, Vijayaraghavan R, Sekhar K. 2004. Effect of α ketoglutarate on cyanide induced biochemical changes in the brain and various cardiorespiratory parameters in rats. Proc Chem Biol Med Treat Sym (CBMTS V), Spiez Laboratory, Switzerland.

Bhattacharya R, Vijayaraghavan R. 1991. Cyanide intoxication in mice through different routes and its prophylaxis by α ketoglutarate. Biomed Environ Sci, 4:452-460.

Bhattacharya R, Vijayaraghavan R. 2002. Promising role of alpha-ketoglutarate in protecting against the lethal effects of cyanide. Hum Exp Toxicol 21:297-303.

Bhattacharya R. 2004. α -Ketoglutarate: A promising antidote to cyanide poisoning. In Flora SJS, J.A.Romano, S.I.Baskin, K.Sekhar, eds, Pharmacological perspectives of toxic chemicals and their antidotes. Narosa Publishing House, New Delhi, India pp 411-430.

Bhattacharya R. 2000. Antidotes to cyanide poisoning: present status. Indian J Pharmacol 32:94 101.

Bhattacharya R and Flora HJS. 2009. Detection and estimation of cyanide. In Handbook of Toxicology of Chemical Warfare Agents. Edited by Ramesh C. Gupta, First Edition. Elsevier Press. 258-259.

Bhattacharya R., Gujar N, Singh R, Rao P, Araghavan V. 2011. Toxicity of alpha-ketoglutarate following 14-days repeated oral administration in Wistar rats. Cell Mol Biol 57:1543-1549.

Bickler PE, Buck LT. 1996. Effects of fructose-1,6-bisphosphate on glutamate release and ATP loss from rat brain slices during hypoxia. J Neurochem 67:1463-1468.

Binder L, Fredrickson L. 1991. Poisonings in laboratory personnel and health care professionals. *Am J Emerg Med* 9:11-15.

Binet L, Wellers G, Dubrisay J. 1951. La valeur et le mécanisme de l'action antidote du tétrathionate de sodium dans l'intoxication cyanhydrique. *Presse Med* 59:641-643.

Birnbaum G, Allegra J, PlosayJJ. 1978. Cyanide poisoning. *Lancet* 92.

Bismuth C. 1992. Haben Antidote eine Zukunft ? Das Beispiel des 4-Methylpyrazolons und des Hydroxocobalamins. *Ther Umschau* 49:118-123.

Bismuth C, Cantineau JP, Pontal P, Baud F, Garnier R, Poulos L, Bolo A. 1984a. Intoxication cyanhydrique. Primauté du traitement symptomatique: Vingt-cinq observations. *Presse Méd* 13:2493-2497.

Bismuth C, Cantineau JP, Pontal P, Baud F, Garnier R, Poulos L, Bolo A. 1984b. Priorité de l'oxygénation dans l'intoxication cyanhydrique: À propos de 25 cas. *J Toxicol Méd* 4:107-121.

Bismuth C, Baud FJ, Djeghout H, Astier A, Aubriot D. 1987. Cyanide poisoning from propionitrile exposure. *J Emerg Med* 5:191-195.

Bismuth C, Baud FJ, Pontal PG. 1988. Hydroxocobalamin in chronic cyanide poisoning. *J Toxicol Clin Exper* 8:35-38.

Blake G. 1839. Observations on the physiological effects of various agents introduced into the circulation, as indicated by the haemodynamometer. *Edinburgh Med Surg J* 51:330-334.

Boggild MD, Peck RW, Tomson CRV. 1990. Acetonitrile ingestion: Delayed onset of cyanide poisoning due to concurrent ingestion of acetone. *Postgrad Med J* 66:40-41.

Bogusz M, Moroz J, Karski J, Gierz J, Regieli A, Witkoska R, Golabek A. 1979. Blood cyanide and thiocyanate concentrations after administration of sodium nitroprusside as hypotensive agent in neurosurgery. *Clin Chem* 25:60-63.

Bolt HM, Lewalter J. 1994. Acrylnitril. In Greim H, Lehnert G, eds, *Biologische Arbeitsstofftoleranzwerte (BAT-Werte) und Expositionsäquivalente für krebserzeugende Arbeitsstoffe (EKA), Arbeitsmedizinisch-toxikologische Begründungen (Biological exposure values for occupational toxicants and carcinogens, critical data evaluation for BAT and BKA values)*, 7th ed. VCH-Verlagsgesellschaft, Weinheim, Germany pp 183-192 [German; English translation].

Bolt HM, Roos PH, Their R. 2002. The cytochrome P-450 isoenzyme CYP2E1 in the biological processing of industrial chemicals: consequences for occupational and environmental medicine. *Int Arch Occup Environ Health* 76:174-185.

Bonbled F. 2001. *Lecture pathologie médico-légale*. Université Catholique de Louvain, Brussels, Belgium.

Bond EJ. 1962. Effect of oxygen on cyanide poisoning in insects. *Nature* 193:1002-1003.

Bonnichsen R, Maehly AC. 1966. Poisoning by volatile compounds III: Hydrocyanid acid. *J Forens Sci* 11:516-528.

Bonsall JL. 1984. Survival without sequelae following exposure to 500 mg/m³ of hydrogen cyanide. *Hum Toxicol* 3:57-60.

Borak J. 1995. Pharmacologic mechanism of antidotes in cyanide and nitrile poisoning. *J Occup Environ Med* 37:793-794.

Borak J. 2009. CPTC international cyanide antidote database (I-CAD). Yale University Occupational Medicine, Jonathan Borak and Company. Cyanide Poisoning Treatment Coalition, Indianapolis, Indiana, USA. www.firesmoke.org/.

Borgohain R, Singh AK, Radhakrishna H, Rao VC, Mohandas S. 1995. Case report: Delayed onset generalised dystonia after cyanide poisoning. *Clin Neurol Neurosurg* 97:213-215.

Borowitz JL, Kanthasamy AG, Mitchell PJ, Isom GE. 1993. Use of PC12 cells as a neurotoxicological screen: Characterization of anticyanide compounds. *Fundam Appl Toxicol* 20:133-140.

Borron SW. 2006. Efficacy of hydroxocobalamin for the treatment of acute cyanide poisoning in adult beagle dogs. *Clin Toxicol* 44:5-15.

Borron SW, Arias JC. 2008. Carbon monoxide-induced status epilepticus in an adult. *J Burn Care Res* 29:559 [Comment on Brown et al, 2007].

Borron SW, Baud FJ. 1996. Acute cyanide poisoning: Clinical spectrum, diagnosis, and treatment. Review UDC 615.917.099.08.Arh hig rada toksikol [Archiv za Higienu Rada i Toksikologiju] 47:307-322.

Borron SW, Baud FJ. 1998. Toxicity, Cyanide. <http://www.emedecine.com>

Borron SW, Baud FJ. 2008. Reply to: "Is hydroxocobalamin safe and effective for smoke inhalation? Searching for guidance in the haze". *Ann Emerg Med* 51:109-110.

Borron SW, Baud FJ. 2008a. In reply [to Wing et al, 2007]. *Ann Emerg Med* 50:108-109.

Borron SW, Baud FJ. 2008b. To the Editor [Reply to Erdman, 2007a,b]. *Ann Emerg Med* 50:109-110.

Borron SW, Baud FJ. 2012. Antidotes for acute cyanide poisoning. *Curr Pharm Biotechnol* 13(10):1940-1948.

Borron SW, Lee J. Toxicity of cyanide, at www.emedicine.com/emerg/topic118.htm.

Borron SW, Baud FJ, Meric P, Seylaz J, Astier A. 1996. Elevation of blood cyanide values after sodium nitrite administration: Implications regarding the mechanism of detoxification. *Toxicol Lett* 88S1:52.

Borron SW, Lapostolle F, Bismuth C, Baud FJ. 1997. Hydroxycobalamin as antidote in suicidal pure cyanide ingestion: Experience in 6 cases. *EAPCCT Scientific Meeting Abstracts*, p 75.

Borron SW, Vicaut E, Ruttimann M, Baud FJ. 2001. Biological tolerance of hydroxocobalamin in fire victims intoxicated by cyanide. *Clin Toxicol* 39:292-293.

Borron SW, Mégarbane B, Baud FJ. 2004. Hydroxocobalamin is an effective antidote in severe acute cyanide poisoning in man. *Int J Toxicol* 23:399-400.

Borron SW, Stonerook M, Reid F. 2006. Efficacy of hydroxocobalamin for the treatment of acute cyanide poisoning in adult beagle dogs. *Clin Toxicol (Phila)* 44S1:5-15.

Borron SW, Baud FJ, Mégarbane B, Bismuth C. 2007a. Hydroxocobalamin for severe acute cyanide poisoning by ingestion or inhalation. *Am J Emerg Med* 25:551-558.

Borron SW, Baud FJ, Barriot P, Imbert M, Bismuth C. 2007b. Prospective study of hydroxocobalamin for acute cyanide poisoning in smoke inhalation. *Ann Emerg Med* 49:794-801, 801.e1-801-e2.

Borron SW, Arias JC, Bauer CR, Sanchez M, Fernandez M, Jung IK. 2009. Hemodynamics after intraosseous administration of hydroxocobalamin or normal saline in a goat model. *Am J Emerg Med* 27:1065–1071.

Borsdorf H, Will W, Zober A. 1991. Chronische Blausäure- und Cyanid-Vergiftungen durch Galvanikarbeiten? Leserfrage, Antwort. *Arbeitsmed Sozialmed Präventivmed* 26:12-16.

Borsdorf H. 1989. Measures for the prevention and treatment of cyanide poisoning in industry. Conseil Européen des Fédérations de l'Industrie Chimique, Bruxelles, IInd European HCN Forum, Medical Day, Nizza, 31.05.1989.

Borsdorf H. 1994. Blausäurevergiftung. Anästhesie, Notfallmedizin, Intensivbehandlung, Schmerzbehandlung. Wissenschaftliche Informationen Fresenius-Stiftung 22:180-182

Botti P. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 4 cyanide cases]. Personal communication. Centro Antiveneni Firenze, Azienda Ospedaliero Universitaria Careggi, Firenze, Italy.

Bouachour G, Harry P, Varache N, Alqtlier PH. 1990. Hyperbaric oxygen in massive cyanide poisoning. Case report with hemodynamic and oxygen consumption data. Undersea Biomed Res 17Suppl:130.

Bourrelrier J, Paulet G. 1971. Intoxication cyanhydrique consécutive à des brûlures graves par cyanure de sodium fondu. Presse Med 79:1013-1014.

Bovykin BA. 1979. [Use of certain complex compounds of bioelements in experimental and clinical medicine]. Farmatsiia 28:62-6.

BP Chemicals. 2002. Material Safety Data Sheet: High Purity Acetonitrile. Naperville, IL, USA.

Braico KT, Humbert JR, Terplan KL, Lehotay JM. 1979. Laetrile intoxication: Report of a fatal case. N Eng J Med 300:238-240.

Brandstein M, Glass P, Castellano A, Mezzacappa C. 1964. The potential danger of inhalation of hydrogen cyanide when using the autoanalyzer. Tech Bull Regist Med Technol 34:122-124.

Braverman B, Ivankovich AD, Lavine LM, Klowden AJ. 1980. Failure of cyanide to antagonize sodium nitroprusside in vivo in dogs. Anesth Analg 59:672-577.

Breen PH, Isserles SA, Westley J, Roizen MF, Taitelman UZ. 1995a. Combined carbon monoxide and cyanide poisoning: A place for treatment? Anesth Analg 80:671-677.

Breen PH, Isserles SA, Westley J, Roizen MF, Taitelman ZU. 1995b. Effect of oxygen and sodium thiosulfate during combined carbon monoxide and cyanide poisoning. Toxicol Appl Pharmacol 134:229-234.

Breen PH, Isserles SA, Tabac E, Roizen MF, Taitelman ZU. 1996. Protective effect of stroma-free methemoglobin during cyanide poisoning in dogs. Anesthesiology 85:558-564.

Brennan DF. 1988. Cyanide poisoning and the anion gap. Letter to the editor [with reply by Krieg and Saxena]. Ann Emerg Med 17:298-299.

- Brenner M, Kim JG, Lee J, Mahon SB, Lemor D, Ahdout R, Boss GR, Blackledge W, Jann L, Nagasawa HT, Patterson SE. 2010c. Sulfanogen sodium treatment in a rabbit model of sub-lethal cyanide toxicity. *Toxicol Appl Pharmacol* 248:269-276.
- Breton D, Jouvet P, De Blic J, Delacourt C, Hubert P. 1993. Toxicité des fumées d'incendie - A propos de deux observations pédiatriques. *Arch Fr Pediatr* 50 :43-45.
- Brettel HF, Hirsch HH, Raudonat HW. 1968. Beobachtungen bei einer mit Kobaltchelate behandelten percutanen Cyanidvergiftung. *Hefte Unfallheilkd* 94:301-303.
- Brian MJ. 1990. Cyanide poisoning in children in Goroka. *P N G Med J* 33:151-153.
- Bridges WR. 1958. Sodium cyanide as a fish poison. Special Scientific Report - Fisheries 253. US Fish and Wildlife Service, Washington DC, USA, pp 1-11.
- Brieger H, Rieders F, Hodes WA. 1952. Acrylonitrile: Spectrophotometric determination, acute toxicity, and mechanism of action. *AMA Arch Ind Hyg Occ Med* 6:128-140.
- Bright JE, Marrs TC. 1982. A comparison of the methemoglobin-inducing activity of moderate oral doses of 4 dimethylaminophenol and p aminophenone. *Toxicol Lett* 13:81-86.
- Bright JE, Marrs TC. 1982. A model for the induction of moderate levels of methaemoglobinaemia in man using 4-diemthylaminophenol. *Arch Toxicol* 50:57-64.
- Bright JE, Marrs TC. 1988. Pharmacokinetics of intravenous potassium cyanide. *Human Toxicology* 7:183-186.
- Bright JE. 1987. A prophylaxis for cyanide poisoning. In Ballantyne B, Marrs TC, eds, *Clinical and experimental toxicology of cyanides*, Chapter 15. John Wright, Bristol, UK, pp 359-382.
- Brink NG, Kuehl FA, Folkers K. 1950. Vitamin B12: The identification of vitamin B12 as a cyano-cobalt coordination complex. *Science* 112:354.
- Brivet F, Delfraissy JF, Duche M, Bertrand P, Dormont J. 1983. Acute cyanide poisoning: Recovery with non-specific supportive therapy. *Intensive Care Med* 9:33-35.
- Broderick KE, Balasubramanian M, Chan A, Potluri P, Feala J, Belke DD, McCulloch A, Sharma VS, Pilz RB, Bigby TD, Boss GR. 2007. The cobalamin precursor cobinamide detoxifies nitroprusside-generated cyanide. *Exp Biol Med* 232:789-798.
- Broderick KE, Potluri P, Zhuang S, Scheffler IE, Sharma VS, Pilz RB, Boss GR. 2006. Cyanide detoxification by the cobalamin precursor cobinamide. *Exp Biol Med* 231:641-649.

- Bromley J, Hughes BGM, Leong DCS, Buckley NA. 2005. Life-threatening interaction between complementary medicines: Cyanide toxicity following ingestion of amygdalin and vitamin C. *Ann Pharmacother* 39:1566-1569.
- Brouard A, Blaisot B, Bismuth C. 1987. Hydroxocobalamine in cyanide poisoning. *J Toxicol Clin Exp* 7:155-168.
- Brown PW, Buckels JA, Jain AB, McMaster P. 1987. Successful cadaveric renal transplantation from a donor who died of cyanide poisoning. *Brit Med J (Clin Res Ed)* 294 (6583):1325.
- Brueske PJ. 1997. ED management of cyanide poisoning. *J Emerg Nurs* 23:569-573.
- Bryson DD. 1978. Cyanide poisoning. *Lancet* 1:92.
- Bryson DD. 1987. Acute industrial cyanide intoxication and its treatment. In Ballantyne B, Marrs TC, eds, *Clinical and experimental toxicology of cyanides*, Chapter 14. John Wright, Bristol, UK, pp 348-358.
- BUA (Beratergremium für umweltrelevante Altstoffe). 1994. Acrylnitril. BUA-Stoffbericht 142, Hirzel Wissenschaftliche Verlagsgemeinschaft, Stuttgart, Germany. ECETOC library.
- Buchanan IS, Dhamee MS, Griffiths FE, Yeoman WB. 1976. Abnormal fundal appearances in a case of poisoning by a cyanide capsule. *Med Sci Law* 16:29-32.
- Bucht H. 1949. On the tubular secretion of thiosulfate and creatinine under the influence of caronamide. *Scand J Clin Lab Invest* 1:270-276.
- Buchter A, Peter H, Appel HE, Bolt HM. 1981. Grundlagen der Toxizität von Acrylnitril und der Behandlung akuter Acrylnitril-Vergiftungen. *Verhandlungen der Deutschen Gesellschaft für Arbeitsmedizin, 21. Jahrestagung in Berlin* 13.- Gentner Verlag Stuttgart 205-208.
- Buchter A, Peter H, Bolt HM. 1984. N-Acetyl-Cystein als Antidot bei akzidenteller Acrylnitril-Intoxikation. *Int Arch Occup Environ Health* 53:311-319.
- Buchter A, Peter H. 1984. Clinical toxicology of acrylonitrile. *G Ital Med Lav* 8:83-86.
- Buchwald A. 1989. Cyanide toxicity. *Ann Emerg Med* 18:176.
- Bulos BA, Thomas BJ, Shukla SP, Sacktor B. 1984. Regulation of pyruvate oxidation in blowfly flight muscle mitochondria: requirement for ADP. *Arch Biochem Biophys* 234:382-393.
- Bunik VI, Romash OG, Gomazkova VS. 1990. Inactivation of alpha-ketoglutarate dehydrogenase during enzyme-catalyzed reaction. *Biochem Int* 22:967-976.

Burrows GE. 1979. Methylene blue or tolonium chloride antagonism of sodium nitrite induced methemoglobinemia. *J Vet Clin Med* 2:81-86.

Burrows GE. 1981. Cyanide intoxication in sheep: Therapeutics. *Vet Hum Toxicol* 23:22-28.

Burrows GE, Way JL. 1976. Antagonism of cyanide toxicity by phenoxybenzamine. *Fed Proc* 35:533.

Burrows GE, Way JL. 1977. Cyanide intoxication in sheep: Therapeutic value of oxygen or cobalt. *Am J Vet Res* 38:223-227.

Burrows GE, Way JL. 1979. Cyanide intoxication in sheep: Enhancement of efficacy of sodium nitrite, sodium thiosulfate, and cobaltous chloride. *Am J Vet Res* 40 613-617.

Burrows GE, Williams C, Davis A, Way JL. 1977. Antagonism of nitrite toxicity by methylene blue or tolonium chloride. *Proc West Pharmacol Soc* 20:439-443. 4.7.1.

Busch H. 2010. ECETOC questionnaire on cyanide and nitrile poisoning [response for 144 cases]. BASF, Ludwigshafen, Germany.

Butuc A. 1969. Aspecte noi in tratamentul intoxicatiilor cu acrilonitril. *Materiale Plastice* 6:106-108.

Cai J, Benz, FW, Campian EC. 2002. Acrylonitrile irreversibly inactivates glyceraldehydes-3-phosphate dehydrogenase by alkylating the catalytically active cysteine 149. *Chem Biol Interact* 140:279-291.

Calder IC, Funder CC, Green CR, Ham K, Tange JD. 1971. Comparative nephrotoxicity of aspirin and phenacetin derivatives. *Br Med J* 4:518-521 [Nephrotoxicity of aminophenol].

California Bureau of Communicable Disease Control. 1972. Cyanide poisoning from the ingestion of apricot kernels. *California Morbidity* 34:September 1.

California Bureau of Communicable Disease Control. 1975. Cyanide poisoning from apricot kernels. *California Morbidity* 51:December 26.

Callaghan JM, Halton DM. 1988. Solutions A and B: Cyanide antidote of folklore myth? *J Soc Occ Med* 38:65-68.

Campbell A. 2000. Hospital management of poisoning in victims suffering from smoke inhalation. *Emerg Nurse* 8:12-16.

- Campbell A, Jones AL. 2001. Cyanide poisoning managed with hydroxocobalmin in the UK. *Clin Toxicol* 39:294.
- Campian EC, Benz FW. 2008. The acute lethality of acrylonitrile is not due to brain metabolic arrest. *Toxicology* 253:104-109.
- Canfield DV, Chaturvedi AK, Dubowski KM. 2005. Carboxyhemoglobin and blood cyanide concentrations in relation to aviation accidents. *Aviat Space Environ Med* 76(10):978-980.
- Cannava A, Chiarlo B. 1951. Sulla possibilita di liberazione dell'anione cianidrico da taluni solfocianati aromatici [Possibility of liberation of hydrocyanic anion by various aromatic sulfocyanates]. *Boll Soc Ital Biol Sper* 1773-1774.
- Caplan YH. 1982. Pathology and pathophysiology of the systemic toxicants carbon monoxide and cyanide. In Cowley RA, Trump BF, eds, *Pathophysiology of shock, anoxia, and ischemia*. Williams and Wilkins 270-279.
- Cappellini FD, Fiorelli G. 2008. Glucose-6-phosphate dehydrogenase deficiency. *Lancet* 371:64-74.
- Caravati EM, Litovitz TL. 1988. Pediatric cyanide intoxication and death from an acetonitrile-containing cosmetic. *J Am Med Assoc* 260:3470-3473.
- Cardozo RH, Edelmann IS. 1952. The volume of distribution of sodium thiosulfate as a measure of the extracellular fluid space. *J Clin Invest* 31:280-290.
- Carella F, Pia Grassi M, Savoiardo M, Contri P, Rapuzzi B, Mangoni A. 1988. Dystonic-Parkinsonian syndrome after cyanide poisoning: clinical and MRI findings. *J Neurol Neurosurg Psychiatry* 51:1345-1348.
- Carter JH, McLafferty MA, Goldman P. 1980. Role of the gastrointestinal microflora in amygdalin (laetrile)-induced cyanide toxicity. *Biochem Pharmacol* 29:301-304.
- Casey P. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 4 cyanide and 3 nitrile cases]. Personal communication. National Poisons Information Centre, Beaumont Hospital, Dublin, Ireland
- Cereda, M. P. and Mattos, M.C.Y. 1996. Linamarin: The toxic compound of Cassava. *J. Venom. Anim. Toxins* [online]. vol.2, n.1, pp. 06-12 .
- Cassel G, Karlsson L, Sellstrom A. 1991. On the inhibition of glutamic acid decarboxylase and gamma-aminobutyric acid transaminase by sodium cyanide. *Pharmacol Toxicol* 69:238-241.

Cescon DW, Juurlink DN. 2009. Discoloration of skin and urine after treatment with hydroxocobalamin for cyanide poisoning. *CMAJ* 180(2):251.

Cetnarowski AB, Conti DR. 1986. Nitroprusside toxicity and low-dose infusion. *Ann Int Med* 104:895-896.

Chai SC, Bruyere JR, Maroney MJ. 2006. Probes of the catalytic site of cysteine dioxygenase. *J Biol Chem* 281:15774-15779.

Chan A, Balasubramanian M, Blackledge W, Mohammad OM, Alvarez L, Boss GR, Bogby TD. 2010. Cobinamide is superior to other treatments in a mouse model of cyanide poisoning. *Clin Toxicol*, 48:709-717

Chan A, Crankshaw DL, Montei A, Patterson SE, Nagasawa HT, Briggs JE, Kozocas JA, Mahon SB, Brenner M, Pilz RB, Bigby TD, Boss GR. 2011. The combination of cobinamide and sulfanegen is highly effective in mouse models of cyanide poisoning. *Clin Toxicol*, 49:366-373

Chanas,B., H.Wang, B.I.Ghanayem (2003): Differential metabolism of acrylonitrile to cyanide is responsible for the greater sensitivity of male vs. female mice: role of CYP2E1 and epoxide hydrolases. *Toxicol Appl Pharmacol*, 193:293-202.

Chang CJ, Garrett MG. 1987. Effect of nutrients and metabolic inhibitors on the growth of spiroplasmas. *Isr J Med Sci* 23:671-675.

Chappey O, Sandouk P, Barron SW, Baud FJ. YEAR?. Kinetics of cobalt after administration of hydroxocobalamin as a cyanide antidote. *EAPCCT Abstracts* ? [Abstract].

Charnock, E.L., J.J.Meehan: Postburn respiratory injury in children. *Ped Clin North America*, 27(1980), 661-676.

Chaturvedi AK, Sanders DC, Endecott BR, Ritter RM. 1995. Exposures to carbon monoxide, hydrogen cyanide and their mixtures: interrelationship between gas exposure concentration, time to incapacitation, carboxyhemoglobin and blood cyanide in rats. *J Appl Toxicol* 15:357-363.

Chaumont AJ, Weil E. 1960. Intoxications mortelles à l'atelier. *Arch Mal Prof* 21:43.

Chen KK, Rose CL. 1952. Nitrite and thiosulfate therapy in cyanide poisoning. *J Am Med Assoc* 149:113-119.

Chen KK, Rose CL. 1956. Treatment of acute cyanide poisoning. *J Am Med Assoc* 162:1154-1155.

Chen KK, Rose CL, Clowes GHA. 1933a. Amyl nitrite and cyanide poisoning. *J Am Med Assoc* 100:1920-1922.

Chen KK, Rose CL, Clowes GHA. 1933b. Potentiation of antidotal action of sodium tetrathionate and sodium nitrite in cyanide poisoning. *Proc Soc Exp Biol Med* 31:252-253 [Experimental study in dogs].

Chen KK, Rose CL, Clowes GHA. 1934. Comparative values of several antidotes in cyanide poisoning. *Am J Med Sci* 188: 767-781.

Chen KK, Rose CL, Clowes GHA. 1944. The modern treatment of cyanide poisoning. *J Indiana State Med Assoc* 37:344-350.

Chen Y, Chen C, Jin S, Zhou L. 1999. The diagnosis and treatment of acute acrylonitrile poisoning: A clinical study of 144 cases. *J Occup Health* 41:172-176.

Cheok SS. 1978. Acute cassava poisoning in children in Sarawak. *Trop Doct* 8:99-101.

Cherian MA, Richmond I. 2000. Fatal methane and cyanide poisoning as a result of handling industrial fish: A case report and review of the literature. *J Clin Pathol* 53:794-795.

Cheung AT, Cruz-Shiavone GE, Meng QC, Pochettino A, Augoustides JA, Bavaria JE, Ochroch EA. 2007. Cardiopulmonary bypass, hemolysis, and nitroprusside-induced cyanide production. *Anesth Analg* 29-33.

Chin RG, Calderon Y. 2000. Acute cyanide poisoning: A case report. *J Emerg Med* 18:441-445.

Chisholm IA, Bronte-Stewart J, Foulds WS. 1967. Hydroxocobalamin versus cyanocobalamin in the treatment of tobacco amblyopia. *Lancet* 2:450-451.

Clark A. 1936. Report on effects of certain poisons contained in food plants of West Africa upon the health of the native races. *J Trop Med Hyg* 39:269-276.

Clark CJ, Campbell D, Reid WH. 1981. Blood carboxyhaemoglobin and cyanide levels in fire survivors. *Lancet* 1:1332-1335.

Clark JM, Lambertsen CJ. 1971. Rate of development of pulmonary toxicity in man during O₂ breathing at 2.0 Ata. *J Appl Physiol* 30:739-752.

Clemedson CJ, Hultman HE, Sörbo B. 1954. The antidote effect of some sulfur compounds and rhodanese in experimental cyanide poisoning. *Acta Physiol Scand* 32:245-251.

Cliff J, Coutinho J. 1995. Acute intoxication from newly-introduced cassava during drought in Mozambique. *Trop Doct* 25:193.

Cock JH. 1982. Cassava: a basic energy source in the tropics. *Science* 218:755-762.

Coentrão L, Neves A, Moura D. 2010a. Hydroxocobalamin treatment of acute cyanide poisoning with a jewellery-cleaning solution. *BMJ Case Rep.* 2010 Jul 15; 2010.

Coentrão L, Moura D. 2010b. Acute cyanide poisoning among jewelry and textile industry workers. *Am J Emerg Med* 29(1):78-81.

Coentrão L and Moura D. 2011. Acute cyanide poisoning among jewelry and textile industry workers. *Am J Emerg Med* 29(1):4.

Cohen MA, Guzzardi LJ. 1984. A letter on the treatment of cyanide poisoning. *Vet Hum Toxicol* 26:503-504.

Cole PV, Vesey CJ. 1987. Sodium thiosulfate decreases blood cyanide concentrations after the infusion of sodium nitroprusside. *Br J Anaesth* 59:531-535.

Coleman MD, Kuhns MJ. 1999. Bioactivation of the cyanide antidote 4-aminopropiophenone (4-PAPP) by human and rat hepatic microsomal enzymes: effect of inhibitors. *Environ Toxicol Pharmacol* 7:75-80.

Colle R. 1970. [Chronic hydrocyanic poisoning]. *Maroc Med* 50:750-757.

Comroe JH, Dripps RD, Dumke PR, Deming M. 1945. Oxygen toxicity: The effect of inhalation of high concentrations of oxygen for twenty-four hours on normal men at sea level and at a simulated altitude of 18,000 feet. *J Am Med Assoc* 128:710-717.

Cope C. 1961. The importance of oxygen in the treatment of cyanide poisoning. *JAMA* 175:1061-1064.

Cope C, Abramowitz S. 1960. Respiratory responses to i.v. sodium cyanide, a function of the oxygen-cyanide relationship. *Am J Vet Res* 81:321-329.

Corbett D, Li J, Benz FW, Nerland DE. 1996. Dose dependence of covalent binding of acrylonitrile to tissue protein and globin in rats. *Fundam Appl Toxicol* 36:149-156.

Cottrell JE, Casthely P, Brodie JD, Patel K, Klein A, Turndorf H. 1978. Mechanism and prevention of tachyphylaxis and cyanide toxicosis after nitroprusside-induced hypotension. *Surg Forum* 29:308-310.

Cottrell JE, Casthely P, Brodie JD, Patel K, Klein A, Turndorf H. 1978. Prevention of nitroprusside-induced cyanide toxicity with hydroxocobalamin. *N Eng J Med* 298:809-811.

Cottrell JE, Patel K, Casthely P, Klein A, Turndorf H. 1989. Nitroprusside tachyphylaxis without acidosis. *Anesthesiology*. Oct;71(4):605-7.

Council of Europe. 2007. European pharmacopoeia, 6th ed. European Directorate for the Quality of Medicines (EDQM). Council of Europe, Strasbourg, France, pp 2588-2589, 2913.

Crane CR, Sanders DC, Endecott BR. 1989. Times-to-incapacitation for rats exposed to carbon monoxide alone, to hydrogen cyanide alone, and to mixtures of carbon monoxide and hydrogen cyanide. DOT/FAA/AM-89. Civil Aeromedical Institute, Federal Aviation Administration, Oklahoma City, Oklahoma 73125, USA.

Crompton M, Palmieri F, Capano M, Quagliariello E. 1974. The transport of sulphate and sulphite in rat liver mitochondria. *Biochem J* 142:127-137.

Crompton M, Palmieri F, Capano M, Quagliariello E. 1974. The transport of thiosulfate in rat liver mitochondria. *FEBS Lett* 46:247-250.

Csiký P. 1955. [Cyanide poisoning (hydrocyanic acid) consecutive to consumption of fruit pit.]. *Orv Hetil* 96:1055-1057 [Hungarian]. FB 1950-1966 CN cas Clinique.

Cummings T. 1996. Medical management of cyanide poisoning. Prepared for HCN forum, ICI Acrylics.

Cummings TF. 2004. The treatment of cyanide poisoning. *J Occup Med* 54:82-85.

Curry AS. 1963. Cyanide poisoning. *Acta Pharmacol Toxicol (Copenh)* 20:291-294.

Curry AS, Price DE, Rutter ER. 1967. The production of cyanide in *post mortem* material. *Acta Pharmacol Toxicol* 25:339-344.

Curry SC, Connor DA, Raschke RA. 1994. Effect of the cyanide antidote hydroxocobalamin on commonly ordered serum chemistry studies. *Ann Emerg Med*. 24:65-67.

Czekay G, Bauer K. 1993. Identification of the thyrotropin-releasing-hormone-degrading ectoenzyme as a metallopeptidase. *Biochem J* 290:921-926.

Dahl AR. 1989. The cyanide-metabolizing enzyme rhodanese in rat nasal respiratory and olfactory mucosa. *Toxicol Lett* 45:199-205.

- Dahl, AR, Waruszewski BA. 1989. Metabolism of organonitriles to cyanide by rat nasal tissue enzymes. *Xenobiotica* 19:1201-1205.
- Dalhamn, T; Edfors, ML, Rylander, R. (1968a) Mouth adsorption of various compounds in cigarette smoke. *Arch Environ Health* 16:831-835.
- Dalhamn, T; Edfors, ML; Rylander, R. (1968b) Retention of cigarette smoke components in human beings. *Arch Environ Health* 17:746-748.
- Dall JLC, Hannah WM. 1964. Oxygen therapy in cyanide poisoning. *Brit Med J* 2:333-334.
- Dally S, Gaultier M. 1976. Anaphylactic shock caused by hydroxocobalamin. *Nouv Presse Méd* 5:1917.
- Dalvi RR, Sawant SG, Terse PS. 1990. Efficacy of α -ketoglutaric acid as an effective antidote in cyanide poisoning in dogs. *Vet Res Comm* 14:411-414.
- Danel V, Serve F, Barret L, Vincent F. 1986. [Cyanide poisoning. Role of hyperbaric oxygen therapy]. [Article in French]. *Presse Med* 15(2):79.
- Darby PW, Wilson J. 1967. Cyanide, smoling, and tobacco amblyopia: Observations on the cyanide content of tobacco smoke. *Br J Ophtalmol* 51:255-257.
- Darmer KI, MacEwen JD, Smith PW. 1972. Short-term animal exposure to carbon monoxide (CO) and hydrogen cyanide (HCN) singly and in combination. *Proceedings of the 3rd International Conference on Environmental Toxicology, Report AMRL-TR-72-130, paper no.22.*
- Dart RC. 2006. Hydroxocobalamin for acute cyanide poisoning: New data from preclinical and clinical studies; new results from the prehospital emergency setting. *Clin Toxicol* 44:1-3.
- Dart RC, Seifert SA, Boyer LV, Clark RF, Hall E, McKinney P, McNally J, Kitchens CS, Curry SC, Bogdan GM, Ward SB, Porter RS. 2001. A randomized multicenter trial of crotalinae polyvalent immune fab (ovine) antivenom for the treatment for crotaline snakebite in the United States. *Arch Intern Med* 161:2030-2036.
- Daunderer M. 1979. Klinische Erfahrungen mit dem Antidot 4-DMAP, einem Methämoglobinbildner zur Behandlung von Vergiftungen mit Blausäure und ihren Salzen, von Schwefelwasserstoff und Salzen der Stickstoffwasserstoffsäure. *Habilitation thesis (Dissertation for appointment as Reader). Technical University of Munich, Germany.*
- Daunderer M. 1979. Tödliche Vergiftung mit Blausäure durch Schwelbrände. *Fortschr Med* 97:1401-1405.

Daunderer M. 1981. Vergiftungstherapie - Antidote (Schluß): Folge 10: Das Antidot 4-DMAP. Fortschr Med 99:1590-1597.

Daunderer M. 1984. Vergiftung mit Brandgasen. Arbeitsmedizin aktuell, Gustav Fischer Verlag, Stuttgart 11.1:69-72.

Daunderer M. 1986 Schnelle Hilfe bei Schwefelwasserstoff-Intoxikationen. Dt Ärztebl 82:31 32

Daunderer M. 1986. Das Antidot Dimethylaminophenol. Krankenhauspharmazie 7:369-373.

Daunderer M, Theml H, Weger N. 1974. Therapy of cyanide poisoning using 4-dimethylaminophenol (4-DMAP). Report on a case of poisoning in man. Med Klin 69(40):1626-1631.

Davies DW, Greiss L, Kadar D, Steward DJ. 1975a. Sodium nitroprusside in children: Observations on metabolism during normal and abnormal responses. Can Anaesth Soc J 22:553-560.

Davies DW, Kadar D, Steward DJ, Munro IR. 1975b. A sudden death associated with the use of sodium nitroprusside for induction of hypotension during anesthesia. Can Anaesth Soc J 22:547-551.

Davies JWL. 1986. Toxic chemicals versus lung tissue – an aspect of inhalation injury revisited. JBCR 7:213-222.

Davis FM, Ewer T. 1988. Acute cyanide poisoning: Case report of the use of hyperbaric oxygen. J Hyper Med 3:103-106.

Davison V. 1969. Cyanide poisoning: Kelocyanor – a new treatment. Occup Health (Lond) 21:306-308.

Dawood MY. 1969. Acute tapioca poisoning in a child. J Singapore Paed Soc 11:154-158.

De Busk RF, Seidl LG. 1969. Attempted suicide by cyanide, a report of two cases. California Medicine 110(5):394-396.

De Francisci G, Magalini SI. 1981. [Progress in the treatment of acute cyanide poisoning]. Recenti Prog Med 71:41-7.

De Forest RE, Potthoff CJ. 1951. Back pressure-arm lift method for administering artificial respiration recommended. JAMA 147:1454-1455.

De la Coussaye JE, Houeto P, Sandouk P, Levillain P, Sassine A, Riou B. 1994. Pharmacokinetics of hydroxocobalamin in dogs. *J Neurosurg Anesthesiol* 6:111-115.

De Lorenzo RA. 1999. Cyanide: the deadly terror weapon that every EMS provider must know about. *JEMS* 24:54-58, 60-61, 64-65.

Decker RH, Dicken CH. 1967. Biochemical observations on leucine aminopeptidase-II of human skin. *J Invest Dermatol* 48:128-132.

DeFrancesco L, Werntz D, Scheffler IE. 1975. Conditionally lethal mutations in Chinese hamster cells. Characterization of a cell line with a possible defect in the Krebs cycle. *J Cell Physiol* 85:293-305.

Degussa. 1999. Safety data sheet (93/112/EC), hydrocyanic acid 98/99% stabilized. Degussa, Hanau, Germany.

Degussa. 2000. Cyanides. Antidotes, information for the medical practitioner. Degussa, Frankfurt am Main, Germany.

Degussa. 2000a. Safety data sheet (93/112/EC), CyPlus(R) sodium cyanide, Bricks 98/99%. Degussa, Hanau, Germany.

Degussa. 2000b. Safety data sheet (93/112/EC), CyPlus(R) potassium cyanide, Bricks 98/99%. Degussa, Hanau, Germany.

Degussa-Röhm. 2007. Arbeitsanweisung, Spezielle Behandlungsrichtlinien CAS-Nr.74-90-8 Rev.-Nr:1 Cyanwasserstoff (HCN).-Werksärztliche Abt., 8 pp.

Degussa-Röhm. 2007. Arbeitsanweisung, Spezielle Behandlungsrichtlinien CAS-Nr/ Rev.-Nr: 107-13-1 Acrylnitril (A.).-Werksärztliche Abt., 7 pp.

Degussa-Röhm. 2007. Arbeitsanweisung, Spezielle Behandlungsrichtlinien CAS-Nr.75-86-5 Rev.-Nr: 1-Acetoncyanhydrin (A).-Werksärztliche Abt., 8 pp.

Delaney K, Cyanide. 1996. In Tintinalli JE, Ruiz E, Krome RL, eds. *Emergency medicine: A comprehensive study guide*, 4th ed. McGraw-Hill, New York, USA, pp 827-831.

Delange F, Bourdoux P, Colinet P, Courtouis P, Hermart P et al. 1983. Nutritional factors involved in the goitrogenic action of cassava. In Delange F, Ahluwalia R, eds, *Cassava toxicity and thyroid: Research and public health issues*. IDRC Monograph 207e. International Development Research Center, Ottawa, Ontario, Canada, pp 17-27.

- Delange F. 1999. Invited commentary: What do we call a goiter? *Eur J Endocrinol* 140:486-488.
- Delga J, Hazard J. 1963. [A new administration form of vitamin B12: hydroxycobalamin.]. *Presse Med* 71:177-180 [French].
- Delga J, Mizoule J, Veverka B, Bon R. 1961a. Recherches sur le traitement de l'intoxication cyanhydrique par l'hydroxocobalamine. *Ann Pharm Fr* 19:740-752.
- Delga J, Mizoule J, Veverka B. 1961b. [The use of hydroxocobalamine (vitamin B-12a) in the treatment of hydrocyanic acid poisoning.]. *Bull Mens Soc Med Mil Fr* 55:340-342 [French].
- Delga J, Mizoule J, Veverka B. 1961c. Sur la valeur de l'hydroxocobalamine dans le traitement de l'intoxication cyanhydrique. *C R Séances Soc Biol Fil* 155:1016-1019.
- Delga J, Motin J, Bouletreau P, Rouzioux JM. 1970. [Current problems of hydrocyanic poisoning: Biological, clinical and therapeutic aspects]. *Thérapeutique* 46:887-892.
- Delhumeau G, Cruz-Mendoza AM, Gomez Lojero C. 1994. Protection of cytochrome c oxidase against cyanide inhibition by pyruvate and α -ketoglutarate: Effect of aeration *in vitro*. *Toxicol Appl Pharmacol* 126:345-351.
- Demedts P, Wauters A, Franck F, Neels H. 1995. Monitoring of cyanocobalamin and hydroxocobalamin during treatment of cyanide intoxication. *Lancet* 346:1706 [Comment on Houeto et al, 1995a].
- Denninghoff K, Walter FG, Langa AJ, He Y, Chipman RA. 2008. Spectrophotometry of hydroxocobalamin and hemoglobin reveals production of an unanticipated methemoglobin variant. *Clin Toxicol (Phila)* 46:545-50.
- Dennis DL, Fletcher WS. 1966. Toxicity of sodium thiosulfate (NCS-45624), a nitrogen mustard antagonist, in dog. *Cancer Chemother Rep* 50:255-257.
- Dequidt J, Furon D, Wattel F, Haguenoer JM, Scherpereel P, Gosselein B, Ginestet A. 1974. Les intoxications par l'acétonitrile à propos d'un cas mortel. *Eur J Toxicol Environ Hyg* 7:91-97.
- DiNapoli J, Hall AH, Drake R, Rumack BH. 1989. Cyanide and arsenic poisoning by intravenous injection. *Ann Emerg Med* 18:308-311.
- Dodds C, McKnight C. 1985. Cyanide toxicity after immersion and the hazards of dicobalt edetate. *Br Med J (Clin Res Ed)* 291:785-786.

Doherty PA, Ferm VH, Smith RP. 1982. Congenital malformations induced by infusion of sodium cyanide in the golden hamster. *Toxicol Appl Pharmacol* 64:456-464.

Doherty PA, Smith RP, Ferm VH. 1982. Tetramethyl substitution on succinonitrile confers pentylenetetrazole-like activity and blocks cyanide release in mice. *J Pharmacol Exp Ther* 223:635-641.

Domalski CA, Kolb LC, Hines EA. 1953. Deleterious reactions secondary to thiocyanate therapy of hypertension. *Proc Mayo Clin* 28:272-280 [Other poisoning].

Done AK. 1961. Cyanide antidotes. In *Clinical pharmacology of systemic antidotes*. *Clin Pharmacol Ther* 2:765-768.

Donoghue AM. 2003. Alternative methods of administering amyl nitrite to victims of cyanide poisoning. *Occup Environ Med* 60:147.

Douce R, Moore AL, Neuburger M. 1977. Isolation and oxidative properties of intact mitochondria isolated from spinach leaves. *Plant Physiol* 60:625-628.

Dougherty C. 1990. L'intoxication cyanhydrique. Renseignements fournis par l'exploration hémodynamique. Thèse Médecine, Angers, France.

Douze JMC, van Heijst ANP. 1973. Herstel van een patient mit kaliumcyanide vergiftiging gevolgd door nitrite intoxicatie [Recovery of a patient with potassium cyanide poisoning followed by nitrite intoxication]. *Ned Tijdschr Geneesk* 177:1409-1412.

Dr. Frank Köhler Chemie GmbH (Hrsg.) *Antidota*. 19 S.

Drew RH. 1983. The use of hydroxocobalamin in the prophylaxis and treatment of nitroprusside-induced cyanide toxicity. *Vet Hum Toxicol* 25:342-345.

Dreykluft T, Göbel U, Taveira O, Nürnberg M, Luft FC, Kettritz R. 2004. Blue acid blues. *Nephrol Dial Transplant* 19:2668-2671. doi:10.1093/ndt/gfh243

Drinker P. 1932. Hydrocyanic acid gas poisoning by absorption through the skin. *J Ind Hyg* 14:1-2.

DrugBank. 2013. <http://www.drugbank.ca/drugs/DB01612> Amyl nitrite.

Drugs.com. 2012. Sodium nitrite (systemic). Drugsite Trust, Auckland, New Zealand. www.drugs.com

Dudley HC, Neal PA. 1942. Toxicology of acrylonitrile (vinyl cyanide). I: A study of the acute toxicity. *J Ind Hyg Toxicol* 24:27-36.

Dueñas-Laita A, Nogué-Xarau S. 2000. Intoxicacion por el humo de los incendios: Tratamiento antidotico a base de vitaminas. *Med Clin (Barc)* 6:658-660.

Dulaney MD, Brumley M, Willis JT, Hume AS. 1991. Protection against cyanide toxicity by oral alpha-ketoglutaric acid. *Vet Hum Toxicol* 33:571-575.

Dunea G. 1983. Death over the counter. *Br Med J (Clin Res Ed)* 286:211-212.

Dwyer MM, Morris CL. 1993. Toxicity of sodium nitroprusside. *Connecticut Med* 57:489-492.

Easterbrook J, Lu C, Sakai Y, Li AP. 2001. Effects of organic solvents on the activities of cytochrome P450 isoforms, UDP-dependent glucuronyl transferase, and phenol sulfotransferase in human hepatocytes. *Drug Metab Dispos* 29:141-144.

European Chemicals Bureau (2004). European Union Risk Assessment Report acrylonitrile. European Union Risk Assessment Report acrylonitrile. Report no.: EUR 20857 EN.

ECETOC. 2007. Cyanides of hydrogen, sodium and potassium, and acetone cyanohydrin (CAS No. 74-90-8, 143-33-9, 151-50-8 and 75-86-5). Joint Assessment of Commodity Chemicals No. 53. European Centre for Ecotoxicology and Toxicology of Chemicals, Brussels, Belgium.

Eckert KG, Elbers R, Eyer P. 1989. Depletion of mitochondrial coenzyme a glutathione by 4-dimethylaminophenol and formation of mixed thioethers. *Biochem Pharmacol* 38:3253-3259.

Eckert KG, Eyer P. 1986. Formation and transport of xenobiotic glutathione-S-conjugates in red cells. *Biochem Pharmacol* 35:325-329.

Eckstein M, Maniscalco PM. 2006. Focus on smoke inhalation – the most common cause of acute cyanide poisoning. *Prehosp Disaster Med* 21:S49-55.

Editorial. 1977. Which antidote for cyanide? *Lancet* 2:1167.

Edwards AC, Thomas ID. 1978. Cyanide poisoning. *Lancet* 1:92-93, 2.14, 7.14.

Eghbal MA, Pennefather PS, O'Brien PJ. 2004. H₂S cytotoxicity mechanism involves reactive oxygen species formation and mitochondrial depolarisation. *Toxicology* 203:69-76.

Ehrlicher H. 1971. Gewerbliche Vergiftungen und ihre Antidote. Gemeinschaftstagung der Werksärzte aus Holland, Österreich, der Schweiz und der Bundesrepublik Deutschland, Mainz, 20.-24.06.1970, Schriftenreihe Arbeitsmedizin, Sozialmedizin, Arbeitshygiene Band 41, Gentner Verlag Stuttgart, 41-51. DP.

Eiró Gonsalves P, Grinberg M, de Souza e Silva OR. 1956. Intoxicação por cianeto (mandioca brava) em pediatria. *Revista Hosp. Clin Univ S Paulo* 11:265-271.

Elbers R, Eyer P, Kampffmeyer H, Soboll S. 1981. Organ toxicity and metabolic pathway of 4-dimethylaminophenol. *Adv Exp Med Biol* 136 A:419-429.

Elbers R, Kampffmeyer HG, Rabes H. 1980. Effects and metabolic pathway of 4-dimethylaminophenol during kidney perfusion. *Xenobiotic* 10:621-632.

Elbers R, Soboll S, Kampffmeyer G. 1980. Alterations in cellular intermediary metabolism by 4-dimethylaminophenol in the isolated perfused rat liver and the implications for 4-dimethylaminophenol toxicity. *Biochem Pharmacol* 29:1747-1753.

Elcombe CR. 2006. Relevance of animal studies to human toxicology. *Clin Toxicol* 44:401-402.

Elliot R. 1997. Cyanide poisoning: new recommendations on first aid treatment. Health and Safety Executive, Sudbury, Suffolk, United Kingdom.

EMA. 2003. Guidance document on the use of medicinal products for the treatment of patients exposed to terrorist attacks with chemical agents. Report EMA/CPMP/1255/03. Scientific Committee, Committee for Proprietary Medicinal Products. European Agency for the Evaluation of Medicinal Products, London, UK.

EMA. 2006. Guidance document on the use of medicinal products for the treatment of patients exposed to terrorist attacks with chemical agents: Blood agents (cyanides/cyanogens). European Agency for the Evaluation of Medicinal Products, London, UK.

Engstrom GW, Reinhardt TA, Horst RL. 1986. 25-Hydroxyvitamin D3-23-hydroxylase, a renal enzyme in several animal species. *Arch Biochem Biophys* 250:86-93.

Erickson TB, Thompson TM, Lu JJ. 2007. The approach to the patient with an unknown overdose. *Emerg Med Clin North Am* 25:249-281.

Espinoza OB, Perez M, Ramirez MS. 1992. Bitter cassava poisoning in eight children: a case report. *Vet Human Toxicol* 34:65.

Essers AJA, Alsen P, Rosling H. 1992. Insufficient processing of cassava induced acute intoxications and the paralytic disease Konzo in a rural area of Mozambique. *Ecol Food Nutr* 27:17-27.

Estler CJ. 1966. [The effect of cobalt gluconate and of the dicobalt salt of ethylenediaminetetraacetic acid (Co-2 EDTA) on the brain metabolism of normal and cyanide-poisoned mice]. *Naunyn Schmiedebergs Arch Exp Pathol Pharmacol* 252:305-313.

European Pharmacopoeia. 1980. Sodium thiosulphate, Dicobalt edetate. In 2nd ed. Maisonneuve, Sainte-Ruffine, France.

Evans CL. 1964. Cobalt compounds as antidotes for hydrocyanid acid. *Brit J Pharmacol* 23:455-475.

Evonik. 2009. Cyanid, Information für den Vertragsarzt. Evonik Degussa, Wesseling, Germany.

Evonik. 2009. Cyanides, information for the medical practitioner. Evonik Degussa, Wesseling, Germany.

Expert Group on the management of chemical casualties caused by terrorist activity. 2003. Treatment of poisoning by selected chemical compounds. Department of Health.

Eybl V, Sýkora J, Köcher Zd. 1959. Der Einfluß von EDTA auf die Auswirkungen von Kobalt *in vivo*. *Naunyn-Schmiedebergs Arch Exp Pathol Pharmacol* 236:199-201.

Eyer P. 1983. The red cell as a sensitive target for activated toxic arylamines. *Arch Toxicol Suppl* 6:3-12.

Eyer P. 1994. Gasförmige Verbindungen – Cyanverbindungen. In Marquardt H, Schaefer S, ed, *Lehrbuch der Toxikologie*. Wissenschaftsverlag Mannheim, Leipzig, Wien, Zuerich pp 555-563.

Eyer P, Gaber H. 1978. Biotransformation of 4-dimethylaminophenol in the dog. *Biochem Pharmacol* 27:2215-2221.

Eyer P, Kampffmeyer HG. 1978. Biotransformation of 4-dimethylaminophenol in the isolated perfused rat liver and in the rat. *Biochem Pharmacol* 27:2223-2228.

Eyer P, Lengfelder E. 1984. Radical formation during autoxidation of 4-dimethylaminophenol and some properties of the reaction products. *Biochem Pharmacol* 33:1005-1013.

Eyer P, Kiese M, Lipowsky G, Weger N. 1973. Metabolism of 4-dimethylaminophenol. *Arch Exp Path Pharmacol* 277:R29.

Eyer P, Kiese M, Lipowsky G, Weger N. 1974. Reactions of 4-dimethylaminophenol with haemoglobin, and autoxidation of 4-dimethylaminophenol. *Chem Biol Interact* 8:41-59.

Eyer P, Hertle H, Kiese M, Klein G. 1975. Kinetics of ferrihaemoglobin formation by some reducing agents and the role of hydrogen peroxide. *Mol Pharmacol* 11:326-334.

Eyer P, Kawan A, Ladstetter B. 1987. Formation of cyanide after i.v. administration of the oxime HI 6 to dogs. *Arch Toxicol* 61:63-69.

Eyer P, Lierheimer E, Strosar M. 1983. Site and mechanism of covalent binding of 4-dimethylaminophenol to human hemoglobin, and its implications to the functional properties. *Mol Pharmacol* 24:282-290.

FAO. 1990. Toxic substances and antinutritional factors, cassava toxicity. In *Roots, tubers, plantains and bananas in human nutrition*, chapter 7. Food and Agriculture Organization of the UN, Rome, Italy [www.fao.org/docrep/T0207E/T0207E08.htm#Cassava%20toxicity]

Faraone G. 1955. L'intossicazione cianidrica negli addetti alle fumigazioni anticoccidiche delle piante [Hydrocyanic acid poisoning in antiscall fumigators]. *Minerva Medicoleg* 75:224-234.

Farooqui MYH, Mumtaz MM. 1991. Toxicology of methacrylonitrile. *Toxicology* 65:239-250.

Favarel-Garrigues JC, Gabinski C, Buil J. 1982. Intoxications cyanhydriques aigües (à propos d'une observation). *Concours Méd* 10:4430-4435.

Favier C. 1992. Essai thérapeutique de fortes doses d'hydroxocobalamine chez les victimes d'indendies. [Therapeutical trial of high doses of hydroxocobalamine in fire victims.] Dissertation, Nice University, Sophia Antipolis, France.

Favier C, Baud F, Richter F, Imbert M, Julien H. 1993. Clinical tolerance of high doses of hydroxocobalamin (HOCO) in fire cyanide poisoning victims. *Anesthesiology* 79:A318.

Favier C, Baud F, Julien H, Bismuth C. 1995. Les différentes formes d'intoxication cyanhydrique et leur traitement. In Baud F, ed, *Réanimation des intoxications aiguës*. Masson, Paris, France, pp 204-213.

Feihl F, Domenighetti G, Perret C. 1982. Intoxication massive au cyanure avec évolution favorable: Étude hémodynamique. *Schweiz Med Wochenschr* 112:1280-1282.

Feldman JM, Feldman MD. 1990. Sequelae of attempted suicide by cyanide ingestion: a case report. *Int J Psychiatry Med* 20:173-179.

Felgenhauer N, Zilker T. 1999. Cyanidintoxikation, Orphan Europe.

Fennel TR, Kedderis GL, Sumner SCJ. 1991. Urinary metabolites of 1,2,3-acrylonitrile in rats and mice detected by ¹³C nuclear magnetic resonance spectroscopy. *Chem Res Toxicol* 4:678-687.

Fernández MC, Teter CA, Borron SW. 2008. Reversal of atrial fibrillation in cyanide poisoning. *Clin Toxicol* 40:622.

Ferralloro G. 1930. The toxicity of hydrocyanic acid in relation to the partial pressure of oxygen. (trans. Title). *Sperimentale*, 84: 115-143.

Ferrari LA, Arado MG, Giannuzzi L, Mastrantonio G, Guatelli MA. 2001. Hydrogen cyanide and carbon monoxide in blood of convicted dead in a polyurethane combustion: A proposition for the data analysis. *Forensic Sci Int* 121:140-143.

Ferraro A. 1933. Experimental toxic encephalomyelopathy – Diffuse sclerosis following subcutaneous injections of potassium cyanide. *Psychiatric Quaterly* 7:267-283.

Finelli PF. 1981. Changes in the basal ganglia following cyanide poisoning. *J Comput Assist Tomography* 5:755-756.

Fiori A, Marigo M. 1963. Tödliche Cyanidvergiftung bei einem Kind. *Arch Toxikol* 20:21-27. FB 1950-1966 CN cas clinique

Flury F, Heubner W. 1919. Über Wirkung und Entgiftung eingeatmeter Blausäure. *Bioch Z* 95:249-256.

Flury F, Zernik F. 1935. Cyanverbindungen. Blausäure. *Abderalden's Handb Biol Arb* 4:1340.

Flury F. Cyanwasserstoff (Blausäure, Ameisnesäurenitril). In *Schädliche Gase* pp 400-409.

Forsyth JC, Becker CE, Osterloh J, Benowitz NL, Rumack BH, Hall AH. 1992. Hydroxocobalamin as a cyanide antidote: Safety, efficacy and pharmacokinetics in heavy smokers. *Vet Hum Toxicol* 34:338.

Forsyth JC, Mueller PD, Becker CE, Osterloh J, Benowitz NL, Rumack BH, Hall AH. 1993. Hydroxocobalamin as a cyanide antidote: Safety, efficacy and pharmacokinetics in heavily smoking normal volunteers. *Clin Toxicol* 31:277-294.

Fortin JL, Giocanti JP, Ruttimann M, Kowalski JJ. 2006. Prehospital administration of hydroxocobalamin for smoke inhalation-associated cyanide poisoning: 8 years of experience in the Paris fire brigade. *Clin Toxicol (Phila)* 44:37-44.

Foulks J, Brazeau P, Koelle ES, Gilman A. 1952. Renal secretion of thiosulfate in the dog. *Am J Physiol* 168:77-85.

Francone MP, Mariani FH. 1951. El BAL y su empleo en el tratamiento de la intoxicación por el ácido cianhídrico [BAL treatment of hydrocyanic acid poisoning.]. *Sem Med (Buenos Aires)* 58:47-50 [Spanish]. FB 1950-1966 CN cas Clinique.

Frank SJ, Tindale N, Everall N. 2002. Decomposition of low levels of acetone cyanohydrin in water. Unpublished report ACRY/Wilton/R+T/2002/04. Lucite International R+T, Wilton Centre, Cleveland, UK.

Frankenberg L, Sörbo B. 1975. Effect of cyanide antidotes on the metabolic conversion of cyanide to thiocyanate. *Arch Toxicol* 33: 81-89.

Frankenberg L. 1982. Effect of cyanide on the distribution of ⁶⁰Co in mice. *Archives of Toxicology* 51:131-137.

Freeman AG. 1972. Food and cyanide. *Br Med J* 3:49.

Freeman AG. 1988. Optic neuropathy and chronic cyanide intoxication: a review. *JRSM*, 81:103-106.

Freeman, J.J. and Hayes, E.P. 1985. Acetone potentiation of acute acetonitrile toxicity in rats. *J. Toxicol. Environ. Health* 15:609-621.

Freeman, J.J. and Hayes, E.P. 1987a. The metabolism of acetonitrile to cyanide by isolated rat hepatocytes. *Fundam. Appl. Toxicol.* 8:263-371.

Freeman, J.J. and Hayes, E.P. 1987b. Evidence for the microsomal metabolism of glycolonitrile. *Biochem. Pharmacol.* 36:184-187.

Freeman, J.J. and Hayes, E.P. 1988. Microsomal metabolism of acetonitrile to cyanide. Effects of acetone and other compounds. *Biochem. Pharmacol.* 37: 1153-1159.

Freierman DE, Cederbaum AI. 1989. Role of cytochrome P-450 IIE1 and catalase in the oxidation of acetonitrile to cyanide. *Chem. Res. Toxicol.* 2:359-366.

Friedberg KD. 1968. Antidote bei Blausäurevergiftungen. *Arch Toxicol*, 24:41-48.,

Friedberg KD, Grützmaker J. 1968. Die Blausäurevergiftung und ihre Behandlung. Theoretische Grundlagen, experimentelle Erfahrungen und praktische Folgerungen. *Wehrmed* 6:61-71.

Friedberg KD, Schwarzkopf HA. 1969. Blausäure-Exhalation bei der Cynidvergiftung [Exhalation of hydrocyanic acid in cyanide poisoning]. Arch Toxikol 24:235-248.

Friedberg KD, Shukla UR. 1975. The efficiency of aquocobalamine as an antidote in cyanide poisoning when given alone or combined with sodium thiosulfate. Arch Toxicol 33:103-113.

Friedberg KD, Grützmaier J, Lendle L. 1965. Aquocobalamin (Vitamin B12a) als spezifisches Blausäureantidot. [with English summary]. Arch Int Pharmacodyn Ther 154:327-350.

Friedberg KD, Grützmaier J, Lendle L. 1966. Die Bedeutung der Wirkungsgeschwindigkeit von Antidoten bei der Behandlung der Blausäurevergiftung. Arch Toxicol 22:176-191.

Friedel HP. 2000. Antidote durch Nichtärzte.

Friedel HP, Jacobs K. 1990. 4-DMAP als Antidot bei Cyanid- und Schwefelwasserstoffvergiftungen. Ständige Konferenz der Werksärzte der Chemischen Industrie, Wacker-Chemie Burghausen, Germany.

Friederich JA, Butterworth JF. 1995. Sodium nitroprusside: Twenty years and counting. Anesth Analg 81:152-162.

Froldi R, Cingolani M, Cacaci C. 2001. A case of suicide by ingestion of sodium nitroprusside. J Forensic Sci 46:1504-1506.

Froneman JC. 1975. A first experience in the use of Kelocyanor. Proc Mine Med Off Assoc SA 54:43.

Gaillard C. 1991. Compte rendu d'hospitalisation de Monsieur P.P – 33 ans. Personal communication. Centre Hospitalier de Sens, Sens, France.

Gajendragad MR, Gopalakrishna S, Ravikumar SB. 1992. Pathology of the brain in acute hydrocyanic acid poisoning in sheep. Indian Vet J 69:206-210 [Case report on sheep poisoned by eating a plant containing cyanogenic glycosides].

Gambaro V, Arnoldi S, Casagni E, Dell'acqua L, Pecoraro C, Froldi R. 2007. Blood cyanide determination in two cases of fatal intoxication: Comparison between headspace gas chromatography and a spectrophotometric method. J Forensic Sci 52:1401-1404.

Garvin CF. 1939. The fatal toxic manifestations of the thiocyanates. J Am Med Assoc 112:1125-1127 [Other poisoning].

Gaultier M, Le Breton R. 1952. A propos d'une intoxication par cyanure alcalin. Ann Med Leg Criminol Police Sci Toxicol 32:385-386.

Geiger JC, Gray JP. 1935. Cyanide poisoning: Additional note on its treatment with intravenous methylene blue solutions. *Cal West Med*, 43:339-342.

Geiger JC. 1933. Methylene blue as antidote for cyanide and carbon monoxide poisoning. *J Am Med Assoc* 100:59.

Geldmacher-Von Mallinckrodt M, van Heijst ANP. 1987. Analytik und Therapie bei Blausäurevergiftung. *Journal of Clinical Chemistry and Clinical Biochemistry* 25:608-609.

Geller RJ, Ekins BR, Iknoian RC. 1991. Cyanide toxicity from acetonitrile-containing false nail remover. *Am. J. Emerg. Med.* 9:268-270.

Gerostamoulos D, Beyer J, Wong K, Wort C, Drummer OH. 2011. Carbon monoxide concentrations in the 2009 Victorian Bushfire disaster victims. *Forensic Sci Int* 205:69-72.

Gershengorn MC, Thaw C. 1982. TRH mobilizes membrane calcium in thyrotropic cells as monitored by chlortetracycline. *Am J Physiol* 243:298-304.

Gerstenberg E. 1957. Zur Sauerstoffbehandlung der Cyanidvergiftung. *Dtsch Med Wochenschr* 82:923-924. FB 1950-1966 CN cas clinique

Gerth K, Ehring T, Braendle M, Schelling P. 2006. Nitric oxide scavenging by hydroxocobalamin may account for its hemodynamic profile. *Clin Toxicol* 44:29-36.

Gettler AO, Baine JO. 1938. The toxicology of cyanide. *Am J Med Sci* 195:182-198.

Ghanayem BI, Sumner SCJ, Fennel TR, Moore TA, Chanas B, Gonzalez F. 1999. Role of cytochrome P450 2E1 in the metabolism of acrylamide and acrylonitrile in mice. *Chem Res Toxicol* 12:1110-1116.

Ghanayem, B.I., M.Y. Farooqui, O. Elshabrawy, M.M. Mumtaz, A.E. Ahmed (1991): Assessment of the acute acrylonitrile-induced neurotoxicity in rats. *Neurotoxicol Teratol*, 13:499-502

Ghiringhelli L. 1955. Tossicità del nitrile adipico. I: Avvelenamento acuto e meccanismo d'azione. *Med Lav* 46:221-228.

Gibson J. 1975. Uptake of C4 dicarboxylates and pyruvate by *Rhodopseudomonas spheroides*. *J Bacteriol* 123:471-480.

Gilman A, Philips FS, Koelle ES. 1946. The renal clearance of thiosulfate with observations on its volume distribution. *Am J Physiol* 146:348-357.

Gladtko E. 1966. [The thiosulfate space in the child.] Supplements to the Archiv für Kinderheilkunde. Ferdinand Enke, Stuttgart, Germany [German].

Gleadow RM. 1999. Resource allocation in cyanogenic Eucalyptus cladocalyx. Thesis University of Melbourne, Melbourne, Australia.

Glende EA Jr. 1972. Carbon tetrachloride-induced protection against carbon tetrachloride toxicity: The role of the liver microsomal drug-metabolizing system. Biochem. Pharmacol. 21: 1697-1702.

Gobbato F, Galzigna L, Bobbio G. 1967. Sulla capacita' della metemoglobina, della idrossocobalamina e dell'edetato di cobalto a fissare l'acido cianidrico [with English summary]. Med Lav 58:376-84.

Goldfrank LR. 1994. In Sullivan JB, Krieger GR, eds, Goldfrank's toxicologic emergencies, 5th ed. Williams and Wilkins, Baltimore, Philadelphia, Hong Kong, London, Munich, Sydney, Tokyo, pp 1215-1229.

Gonzales J, Sabatini S. 1989. Cyanide poisoning: pathophysiology and current approaches to therapy. Int J Artif Organs 12:347-355.

Goodhart GL. 1994. Patient treated with antidote kit and hyperbaric oxygen survives cyanide poisoning. South Med J 87:814-816.

Goodman LS, Gilman A. 1956. The pharmacological basis of therapeutics, 2nd ed. The Macmillan Company, New York, NY, USA.

Goodman LS, Gilman A, eds. 1975. The pharmacological basis of therapeutics, 5th ed. Macmillan Publishing Co, New York, NY, USA.

Gordh T, Norberg B. 1947. Studies on oxygen treatment in connection with experimental hydrocyanic poisoning. Acta Physiol Scand 13:26-34.

Gossel TA Bricker JD. 1994. Principles of Clinical Toxicology. 3rd ed. 117-119. Raven Press, Ltd., New York, NY.

Gourlain H, Caliez C, Laforge M, Buneaux F, Levillain P. 1994. Study of the mechanisms involved in hydroxocobalamin interference with determination of some biochemical parameters. Ann Biol Clin 52:121-124.

Gourlain H, Buneaux F, Levillain P. 1996. Mesure du CO et de la COHb dans le sang: Interférences de l'hydroxocobalamine et du bleu de méthylène en CO-oxymétrie. Rev Fr Lab 282:144-148.

Gracia R, Pharm D, Greene S. 2004. Cyanide Poisoning and its treatment. *Pharmacotherapy* 10:1359-1365.

Graham DL, Laman D, Theodore J, Robin ED. 1977. Acute cyanide poisoning complicated by lactic acidosis and pulmonary edema. *Arch Intern Med* 137:1051-1055.

Graham JDP. 1963. Hydroxycobalamin as an antidote to acrylonitrile. *Toxicol Appl Pharmacol* 7:367-372.

Grahl R. 1972. Toxikologie und Wirkungsweise von Acrylnitril. *Zbl Arbeitsmed*, 12:369-378.

Grant WM. 1986. Toxicology of the eye, 3rd ed. Charles C. Thomas, Springfield, Illinois, USA.

Gras C, Fritz KW, Staffensky RH; Grotte R, Kirchner E. 1991. Die akzidentelle Schwefelwasserstoffvergiftung. *Notarzt* 149-151

Grassl SM. 2002. Facilitated diffusion of urate in avian brush-border membrane vesicles. *Am J Physiol Cell Physiol* 283:C1155-1162.

Green DE, Williamson S. 1937. Pyruvic and oxaloacetic cyanohydrins. *Biochem.J* 31:617-618

Greene NM. 1961. Effect of epinephrine on lactate, pyruvate, and excess lactate production in normal human subjects. *J Lab Clin Med* 58:682-686.

Greenfield S, Claus GW. 1969. Isocitrate dehydrogenase and glutamate synthesis in *Acetobacter suboxydans*. *J Bacteriol* 100:1264-1270.

Grunske F. 1949. Ventox und Ventox-Vergiftung. *Dtsch Med Wschr* 74:1081-1083.

Guest A, Jackson JR, James SP. 1982. Toxicity of benzyl cyanide in the rat. *Toxicol Lett* 10:265-72.

Gunder AE, Brahamov AA, Weisenberg E, Gratner S, Shpern S. 1969. [Cyanide poisoning in a child due to the consumption of apricot pits]. *Harefuah* 76:536-538 [Hebrew; English translation].

Gut I, Kopecký J, Nerudová J. 1981. Relationship between acrylonitrile biotransformation, pharmacokinetics, and acute toxicity. A short review. *G Ital Med Lav* 3:131-136.

Gut I, Nerudová J, Kopecký J, Holeček V. 1975. Acrylonitrile biotransformation in rats, mice, and Chinese hamsters as influenced by the route of administration and by Phenobarbital, SKF 525-A, cysteine, dimercaprol, or thiosulfate. *Arch Toxicol* 33:151-161.

Haasnoot K, van Vught AJ, Meulenbelt J, Bergman LR. 1989. Acute blauwzuurvergiftiging bij een zuigeling. *Ned Tijdschr Geneesk* 133:1753-1756.

Hackstadt T, Williams JC. 1983. pH dependence of the *Coxiella burnetii* glutamate transport system. *J Bacteriol* 154:598-603.

Hager HHJ. 1977. *Hagers Handbuch der pharmazeutischen Praxis*, 4th ed. Springer, Berlin, Germany.

Hagers Enzyklopädie der Arzneistoffe und Drogen -6 Auflage, Stuttgart Wissenschaftliche Verlagsgesellschaft 2007 88

Haguenoer JM, Dequidt J, Jacquemont MC. 1975a. Intoxications expérimentales par l'acétonitrile. 1ère note: Intoxications aiguës par voie intrapéritonéale. *J Eur Toxicol* 8:94-101.

Haguenoer JM, Dequidt J, Jacquemont MC. 1975b. Intoxications expérimentales par l'acétonitrile. 2ème note: Intoxications aiguës par voie pulmonaire. *J Eur Toxicol* 8:102-106.

Haiharakrishnan J, Satpue RM, Bhattacharya R. 2010. Cyanide-induced changes in the levels of neurotransmitters in discrete brain regions and their response to oral treatment with α -Ketoglutarate. *Indian J Exp Biol* 48:731-736.

Hall AH. 1986. Cyanide poisoning: Dealing with unexpected menace. *Emergency Med* 18:919-205.

Hall AH, Rumack BH. 1986. Clinical toxicology of cyanide. *Ann Emerg Med* 15:1067-1074.

Hall AH, Rumack BH. 1987. Hydroxycobalamin/sodium thiosulfate as a cyanide antidote. *J Emerg Med* 5:115-121.

Hall AH, Linden CH, Kulig KW, Rumack BH. 1986. Cyanide poisoning from laetrile ingestion: Role of nitrite therapy. *Pediatrics* 78:269-272.

Hall AH, Dautre WH, Ludden T, Kulig KW, Rumack BH. 1987a. Nitrite/thiosulfate treated acute cyanide poisoning: Estimated kinetics after antidote. *Journal of toxicology. Clin Toxicol* 25:121-133.

Hall AH, Rumack BH, Schaffer MI, Linden CH. 1987b. Clinical toxicology of cyanide: North American clinical experiences. In Ballantyne B, Marrs TC, eds, *Clinical and experimental toxicology of cyanides*, Chapter 12. John Wright, Bristol UK, pp 312-333.

Hall AH, Kulig KW, Rumack BH. 1989a. Suspected cyanide poisoning in smoke inhalation: Complications of sodium nitrite therapy. *J Toxicol Clin Exp* 9:3-9.

Hall AH, Kulig KW, Rumack BH. 1989b. Toxic smoke inhalation. *Am J Emerg Med* 7:121-122.

Hall AH, Dart R, Bogdan G. 2007. Sodium thiosulfate or hydroxocobalamin for the empiric treatment of cyanide poisoning? *Ann Emerg Med* 49:806-813.

Hall AH, Sainers J, Baud F. 2009. Which cyanide antidote? *Crit Rev Toxicol* 39:541-552.

Hall VA, Guest JM. 1992. Sodium nitroprusside-induced cyanide intoxication and prevention with sodium thiosulfate prophylaxis. *Am J Crit Care* 1:19-25.

Halpern J, Guastalla G, Bercaw J. 1972. Some aspects of the chemistry of cobalt (I) cyanide and related complexes. *Coord Chem Rev* 8:167-184.

Halström F, Möller KN. 1945. The content of cyanide in human organs from cases of poisoning with cyanide taken by mouth. With a contribution to the toxicology of cyanides. *Acta Pharmacol* 1:18-28.

Hames J, Warin RP. 1971. Sensitivity to cyanocobalamin and hydroxocobalamin. *Br Med J* 2:262.

Hantson P, Baud FJ, Garnier R. 1996. Toxic gases. In Descotes J, ed, *Human Toxicology*. Elsevier Science BV, pp 661-669.

Hantson P, N'Geye P, Laforge M, Clemessy JL, Baud F. 1996. Suicide attempt by ingestion of potassium ferricyanide. *J Toxicol Clin Toxicol* 34:471-473.

Hantson P, Squifflet JP, Vanormelingen P, Mahieu P. 1999. Organ transplantation after fatal cyanide poisoning. *Clin Transplant* 13:72-73.

Hantson P. 2000. Lecture « Physiopathologie des intoxications », Univ.Cathol.de Louvain, Brussels, 4/2000.

Hantson Ph, Benaissa L, Baud F. 1999. Intoxication par les fumées d'incendie. *La Presse Médicale* 28:1949-1954.

Hariharakrishnan J, Anand T, Satpute RM, Jayaraj R, Prasad GBKS, Bhattacharya R. 2009. Activity and gene expression profile of certain antioxidant enzymes in different organs of rats after subacute cyanide exposure: Effect of alpha-ketoglutarate. *Drug and Chemical Toxicology* 32:268-276.

Harloff FP, Sauer O, Weilemann LS. 1995. Kritische Analyse von Vergiftungen mit Blausäure und cyanhaltigen Substanzen anhand 182 Beratungen der Mainzer Giftinformationszentrale. *Intensivmedizin und Notfallmedizin* 6:499-500

Harry P, Bouachour G, Clément A, Guilleux-Dumont AM, Alquier P. 1985. Renseignements fournis par l'exploration hémodynamique au cours d'une intoxication cyanhydrique. Personal communication. Centre Anti-Poison et Réanimation Médicale, Angers, France.

Hartung R. 1985. Cyanides and nitriles. In Clayton GD and Clayton FE, eds. *Patty's Industrial hygiene and toxicology*, 3rd ed., vol 2C. John Wiley & Sons, New York, USA, 4845-49001.

Hashimoto K, Kanai R. 1965. Studies on the toxicology of acrylonitrile: Metabolism, mode of action and therapy. *Ind. Health* 3:30-46.

Hatch RC, Laflamme DP, Jain AV. 1990. Effects of various known and potential cyanide antagonists and a glutathione depletor on acute toxicity of cyanide in mice. *Vet Hum Toxicol* 32:9-16.

Hausmann D, Pfeifer G, Mück R. 1982. Vergiftungen im Notarztdienst: auf Mandelgeruch achten. *Notfallmedizin* 8:2-12.

Haustein J, Gabsch HC, Lohs K. 1969. Die Cyanidentgiftung durch DL-Glycerinaldehyd. *Pharmazie* 24:482-483.

Hawkins SF, Groff WA, Johnson RP, Froehlich HL, Kaminskis A, Stemler FW. 1981. Comparison of the in vivo formation of methemoglobin by 4-dimethylaminophenol and sodium nitrite in the *Cynomolgus* monkey. *Fed Proc* 40:718.

HSE. 1997. Cyanide poisoning. New recommendations on first aid treatment. Health and Safety Executive, UK.

Healy JC. 1931. Therapeutics and toxicology of sulfocyanates. *New Engl J Med* 205:5481-5583 [Other poisoning, by thiocyanate].

Heinemeyer G. 1989. Cyanidantidote. *Notfallmedizin* 15:709-711.

Heintz B, Bock TA, Kierdorf H, Sieberth HG. 1990. Cyanid-Intoxikation: Behandlung mit Hyperoxygenation und Natriumthiosulfat. *Dtsch Med Wochenschr* 115:1100-1103..

Henry J, Volans G. 1984. ABC of poisoning: Immediate measures outside hospital. *Br Med J (Clin Res Ed)* 289:39-40.

Herbert V, Sullivan LW. 1964. Activity of coenzyme B12 in man. *Ann NY Acad Sci* 112:855-870.

Herbert V. 1979. Laetrile: the cult of cyanide: Promoting poison for profit. *Am J Clin Nutr* 32:1121-1158.

Hermann LW. 1886. Ueber die Wirkung des Nitroprussidnatriums. *Arch Ges Physiol* 39:419-424.

Herold G. 2011. Innere Medizin. Köln, Germany. www.herold-innere-medizin.de

Hewick DS, Butler AR, Glidewell C, McIntosh AS. 1987. Sodium nitroprusside: Pharmacological aspects of its interaction with hydroxocobalamin and thiosulphate. *J Pharm Pharmacol* 39:113-117.

Hill HE. 1942. A contribution to the toxicology of sodium nitroprusside 1. The decomposition and determination of sodium nitroprusside. *Aust Chem Inst J Proc* 9:89-93.

Hill Top Research. 1997. Delayed Contact Hypersensitivity Study with Acetonitrile (HPLC Grade) in Guinea Pigs (Buehler Technique).

Hillman B, Bardhan K, Bain J. 1974. The use of dicobalt edetate (Kelocyanor) in cyanide poisoning. *Postgrad Med J* 50:171-174.

Himwich WA, Saunders JP. 1948. Enzymatic conversion of cyanide to thiocyanate. *Am J Physiol* 153:348-354.

Hinsberg O, Treupel G. 1984. Über die physiologische Wirkung des p-Amidophenols und einiger Derivate desselben. *Naunyn-Schmiedebergs Arch Exp Pathol Pharmacol* 33:216-250. 4.7.1

Hoang The Dan P, Bretszajn A, Pourriat JL, Lapandry C, Cupa M. 1982. Un cas d'intoxication au cyanure. *Anesth Analg (Paris)* 38:161-162.

Hoang The Dan P, Bretszajn A, Pourriat JL, Lapandry C, Cupa M. 1981. Un cas d'intoxication aigue au cyanure. *Anesth Anal Rean* 38:161-162.

James M, Kreye VA, Pill J. 1978. Effect of sodium nitroprusside alone and in combination with sodium thiosulfate on the acid-base balance, and on thiocyanate and iron plasma levels in the rabbit. *Klin Wochenschr* 56 1:147-152.

- Höbel M, Engeser P, Nemeth L, Pill J. 1980. The antidote effect of thiosulphate and hydroxocobalamin in formation of nitroprusside intoxication of rabbits. *Arch Toxicol* 46:207-213.
- Hodgkins PS, Schwarcz R. 1998. Interference with cellular energy metabolism reduces kynurenic acid formation in rat brain slices: reversal by lactate and pyruvate. *Eur J Neurosci* 10:1986-1994.
- Holland DJ. 1983. Cyanide poisoning: an uncommon encounter. *Journal of Emergency Nursing* 9:138-140 [General overview, no specific case].
- Holland MA, Kozlowski LM. 1986. Therapy reviews: clinical features and management of cyanide poisoning. *Clinical Pharmacy* 5:737-741 [General overview, no specific case].
- Hoppu K. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 0 cases]. Personal communication. Poison Information Centre, Helsinki University Central Hospital, Helsinki, Finland.
- Horwitz ET. 1979. Of love and laetrile: Medical decision making in a child's best interests. *Am J Law Med* 5:271-294.
- Houeto P, Buneaux F, Galliot-Guilley M, Baud FJ, Levillain P. 1994. Determination of hydroxocobalamin and cyanocobalamin by derivative spectrophotometry in cyanide poisoning. *J Anal Toxicol* 18:154-158.
- Houeto P, Hoffman JR, Imbert M, Levillain P, Baud FJ. 1995a. Relation of blood cyanide to plasma cyanocobalamin concentration after a fixed dose of hydroxocobalamin in cyanide poisoning. *Lancet* 346:605-608. WS thesis.
- Houeto P, Hoffman JR, Imbert M, Levillain P, Baud FJ. 1995b. Authors' reply: Monitoring of cyanocobalamin and hydroxocobalamin during treatment of cyanide intoxication. *Lancet* 346:1706-1707. WS thesis.
- Houeto P, Borron SW, Sandouk P, Imbert M, Levillain P, Baud FJ. 1996. Pharmacokinetics of hydroxocobalamin in smoke inhalation victims. *Clin Toxicol* 34:397-404.
- Howell SB, Pfeifle CL, Wung WE, Olshen RA, Lucas WE, Yon JL, Green M. 1982. Intraperitoneal cisplatin with systemic thiosulfate protection. *Ann Intern Med* 97:845-851.
- Howlett WP, Brubaker GR, Mlingi N, Rosling H. 1990. Konzo, an epidemic upper motor neuron disease studied in Tanzania. *Brain* 113:223-235.

- Huang J, Niknahad H, Khan S, O'Brien PJ. 1998. Hepatocyte-catalysed detoxification of cyanide by L- and D-cysteine. *Biochem Pharmacol* 55:1983-1990.
- Humbert JR, Tress JH, Braico KT. 1977. Fatal cyanide poisoning: Accidental ingestion of amygdalin. *JAMA* 238:482.
- Hume AS, Moore SJ, Hume AT. 1996. Effects of α -ketoglutaric acid on the distribution of cyanide and acidosis associated with cyanide intoxication. *Toxicologist* 3:98.
- Hume AS, Mozingo JR, McIntyre B, Ho IK. 1995. Antidotal efficacy of alpha-ketoglutaric acid and sodium thiosulfate in cyanide poisoning. *ClinToxicol* 33:721-724.
- Humphrey SH, Nash DA. 1978. Lactic acidosis complicating sodium nitroprusside therapy. *Ann Int Med*, 88:58-59.
- Hung D, Hsu S, Chen Y. 2005. Cyanide intoxication in an acrylonitrile chemical accident. *Clin Toxicol* 43:751.
- Hung DZ, Tsan YT, Yu YJ, Hsu CL. 2009. Cyanide poisoning in Taiwan. *Lancet* 374:1212 [Erratum in: *Lancet* 374:1820.
- Hung YM, Hung SY, Olson KR, Chou KJ, Lin SL, Chung HM, Tung CN, Chang JC. 2007. Yam bean seed poisoning mimicking cyanide intoxication. *Intern Med J* 37:130-132.
- Hunt R. 1905. The influence of thyroid feeding upon poisoning by acetonitrile. *J. Biol. Chem* 33-44.
- Ignarro LJ, Harbison RG, Wood KS, Kadowitz PJ. 1986. Dissimilarities between methylene blue and cyanide on relaxation and cyclic GMP formation in endothelium-intact intrapulmonary artery caused by nitrogen oxide-containing vasodilators and acetylcholine. *J Pharmacol Exp Ther* 26:30-36 [Does not address cyanide mechanism of action but interactions with sodium nitroprusside].
- ILO. 2000a. Sodium nitrite ICSC 1120, October 2000. In International chemical safety cards. International Labour Organization, International Programme on Chemical Safety and the European Commission. International Occupational Safety and Health Information Centre (CIS), Geneva, Switzerland..
www.ilo.org/public/english/protection/safework/cis/products/icsc/dtasht/index.htm

ILO. 2000b. Isoamyl nitrite ICSC 1012, November 1998. In International chemical safety cards. International Labour Organization, International Programme on Chemical Safety and the European Commission. International Occupational Safety and Health Information Centre (CIS), Geneva, Switzerland.

www.ilo.org/public/english/protection/safework/cis/products/icsc/dtasht/index.htm

ILO. 2004. Oxygen ICSC 0138, October 1999. In International chemical safety cards. International Labour Organization, International Programme on Chemical Safety and the European Commission. International Occupational Safety and Health Information Centre (CIS), Geneva, Switzerland.

www.ilo.org/public/english/protection/safework/cis/products/icsc/dtasht/index.htm

ILO. 2008. Sodium thiosulfate ICSC 1138, April 2006. In International chemical safety cards. International Labour Organization, International Programme on Chemical Safety and the European Commission. International Occupational Safety and Health Information Centre (CIS), Geneva, Switzerland.

www.ilo.org/public/english/protection/safework/cis/products/icsc/dtasht/index.htm

Imbert M, Lambert E, Michel A, Julien H. 1993. L'hydroxocobalamine: antidote de choix des intoxications cyanhydriques au cours des feux d'habitation. *Urgences* 5:202-209.

Ioanid N, Bors G. 1961. Contribution chimico-judiciaire a l'étude des empoisonnements par l'acide cyanhydrique. *Ann Méd Lég* 41:275-279.

IPCS. 1983. Acrylonitrile. Environmental Health Criteria 28. International Programme on Chemical Safety, World Health Organisation, Geneva, Switzerland.

[www.inchem.org/documents/ehc/ehc/ehc28.htm].

IPCS. 1993a. Antidotes for poisoning by cyanide. In Meredith TJ, Jacobsen D, Haines JA, Berger JC, van Heijst ANP, eds, International Programme on Chemical Safety, Commission of the European Communities, Evaluation of antidotes series 2. EUR 14280 EN. Cambridge University Press, Cambridge, UK. [www.inchem.org/documents/antidote/antidote/ant02.htm]

IPCS. 1993b. Acetonitrile. Environmental Health Criteria 154. International Programme on Chemical Safety, World Health Organisation, Geneva, Switzerland.

[www.inchem.org/documents/ehc/ehc/ehc154.htm]

IPCS. 2004. Concise International Chemical Assessment Document, hydrogen cyanide and cyanides, human health aspects. International Programme on Chemical Safety, World Health Organization, Geneva, Switzerland.

IPCS. 2009. IPCS/CEC evaluation of antidotes series. Volume 2: Antidotes for poisoning by cyanide. International Programme on Chemical Safety, World Health Organization, Geneva, Switzerland.

Isom GE, Way JL. 1972. Cyanide intoxication: protection with cobaltous chloride. *Toxicol Appl Pharmacol* 24:449-456.

Isom GE, Way JL. 1974. Effect of oxygen on cyanide intoxication. VI. Reactivation of cyanide-inhibited glucose metabolism. *J Pharmacol Exp Ther* 189:235-243.

Isom GE, Way JL. 1984. Effects of oxygen on the antagonism of cyanide intoxication: Cytochrome oxidase, *in vitro*. *Toxicol Appl Pharmacol* 74:57-62.

Isom GE, Liu DWH, Way JL. 1975. Effect of sublethal doses of cyanide on glucose catabolism. *Biochem Pharmacol* 24:871-875.

Isom GE, Burrows GE, Way JL. 1982. Effect of oxygen on the antagonism of cyanide intoxication: Cytochrome oxidase, *in vivo*. *Toxicol Appl Pharmacol* 65:250-256.

Isom GE, Borowitz JL. 1995. Cyanide-induced neurotoxicity: mechanism of action. Presented at the Wenner-Gren International Symposium on Cyanide Neuro, Stockholm, Sweden.

Isom GE, Borowitz JL. 1995. Modification of cyanide toxicodynamics: mechanistic based antidote development. *Toxicol Lett* 82/83:795-799.

IUBMB. 1999. Enzyme nomenclature, recommendations 1992. Supplements 1-5, 1994-1999. Nomenclature Committee of the International Union of Biochemistry and Molecular Biology. Academic Press, San Diego, California, USA. *Eur J Biochem* 223:1-52; 32:1-6; 237:1-5; 250:1-6; 264:610-650. [www.chem.qmul.ac.uk/iubmb/enzyme/]

IUPAC. 1969. Dissociation constants of inorganic acids and bases in aqueous solutions. Oxford, United Kingdom, International Union of Pure and Applied Chemistry.

Ivankovich AD, Braverman B, Kanuru RP, Heyman HJ, Paulissian R. 1980. Cyanide antidotes and methods of their administration in dogs: A comparative study. *Anesthesiology* 52:210-216.

Ivankovich AD, Braverman B, Stephens TS, Shulmann M, Heyman HJ. 1983. Sodium thiosulfate disposition in humans: Relation to sodium nitroprusside toxicity. *Anesthesiology* 58:11-17.

Ivanov JP. 1959. [The effect of elevated oxygen pressure on animals poisoned with potassium cyanide]. *Pharmacol Toxicol (USSR)* 22:476-479 [Russian]. IPCS 2.14

Izraeli S, Israeli A, Danon Y. 1988. [Pharmacological treatment of cyanide poisoning]. Harefuah 114:338-42.

Jack RD. 1974. Toxicity of sodium nitroprusside. Br J Anaesth 46:952.

Jacobs K. 1971. Erfahrungsbericht über Anwendung von Kelocyanor bei Blausäurevergiftung. Zentralbl Arbeitsmed 21:372-374.

Jacobs K. 1984. Erfahrungsbericht über Anwendung von 4-DMAP (Dimethylaminophenol) bei einer schweren Blausäure-Vergiftung, Konsequenzen für die Praxis. Zentralbl Arbeitsmed 34:274-277.

Jacobs K. 1990. Cyanid-Intoxikation. Dtsch med Wschr 115:1895.

Jaeger A, Tempe JD, Porte A, Stoeckel L, Mantz JM. 1977. Acute voluntary intoxication by acetonitrile. Acta Pharmacol Toxicol (Copenh) 41 Suppl 2:340.

Jakubowski M, Linhart I, Pielas G, Kopecký J. 1987. 2-Cyanoethylmercapturic acid (CEMA) in the urine as a possible indicator of exposure to acrylonitrile. Br J Ind Med 44:834-840.

Jancso P, Szinicz L, Eyer P. 1981. Biotransformation of 4-dimethylaminophenol in man. Arch Toxicol 47:39-45.

Jandorf B, Bodansky O. 1946. Therapeutic and prophylactic effect of methemoglobinemia in inhalation poisoning by hydrogen cyanide and cyanogen chlorid. J Ind Hyg Toxicol 28:125-132.

Jeannin J, Berrod R, Lemaire, Bazin. 1961. Intoxication grave survenue chez une enfant après ingestion de noyaux de pêches. Lyon Méd 93:785-793.

Jerretin S. 1963. Un succès thérapeutique dans la réanimation d'un sujet intoxiqué au cyanure de potassium. Zdraw Vestn 32:3-4.

Jervis G. 1937. Sulle alterazioni mieliniche da cianuro. Rivista di Patologia Nevosa e Mentale 50, (Sept.-Oct.): 410-419.

Jiang S, Liu Z, Zhuang X. 1998. Effect of procaine hydrochloride on cyanide intoxication and its effect on neuronal calcium in mice. Toxicol Appl Pharmacol 150:32-36.

Johanning RJ, Zaske DE, Tschida SJ, Johnson SV, Hoey LL, Vance-Bryan K. 1995. A retrospective study of sodium nitroprusside use and assessment of the potential risk of cyanide poisoning. Pharmacotherapy 15:773-777.

Johannsen FR, Levinskas GJ. 1986. Relationship between toxicity and structure of aliphatic nitriles. Fundam Appl Toxicol 7:690-697.

Johnson RP, Mellors JW. 1988. Arteriolization of venous blood gases: A clue to the diagnosis of cyanide poisoning. *Am J Emerg Med* 6:401-404.

Johnson WR, Hale RW, Clough SC, Chen PH. 1973a. Chemistry of the conversion of nitrite nitrogen to smoke products. *Nature* 243:223-225 [Cited by IARC, 1986].

Johnson WS, Hall AH, Rumack BH. 1989. Cyanide poisoning successfully treated without 'therapeutic methemoglobin levels'. *Am J Emerg Med* 7:437-440.

Johnson-Flanagan AM, Spencer MS. 1981. The effect of rotenone on respiration in pea cotyledon mitochondria. *Plant Physiol* 68:1211-1217.

Jones AW, Löfgren A, Eklund A, Grundin R. 1992. Two fatalities from ingestion of acetonitrile: Limited specificity of analysis by headspace gas chromatography. *J. Anal. Toxicol* 16:104-106.

Jouglar J, Nava G, Botta A, Michel-Manuel C, Poyen B, Mattei JL. 1974. A propos d'une intoxication aigue par le cyanure de potassium traitée par l'hydroxocobalamine. *Marseille Med* 111:617-624.

Jouglard J, Fagot G, Deguigne B, Arlaud JA. 1971. L'intoxication cyanhydrique aiguë et son traitement d'urgence. *Marseille Médical* 9:1-5.

Jouglard J, Nava G, Botta A, Michel-Manuel C, Poyen B, Mattei JL. 1974. A propos d'une intoxication aiguë par le cyanure de potassium traitée par l'hydroxocobalamine. *Marseille Méd* 12:617-624.

Jourdan C, Convert J, Tell L, Wasylkiewicz Y, Artru F. 1993. Evolution de la pression intracrânienne (PIC) et des lésions cérébrales au cours d'une intoxication cyanhydrique (CN) aiguë grave. Paper presented at XXXIème Congrès de la Société de Toxicologie Clinique.

Kaczka EA, Wolf DE, Kuehl FA, Folkers K. 1950. Vitamin B12: Reactions of cyanocobalamin and related compounds. *Science* 112:354-355.

Kalyanaraman UP, Kalyanaraman K, Cullinan SA, McLean JM. 1983. Neuromyopathy of cyanide intoxication due to 'laetrile' (amygdalin): A clinicopathologic study. *Cancer* 51:2126-2133.

Kampe S, Iffland R, Korenkov M, Diefenbach C. 2000. Survival from a lethal blood concentration of cyanide with associated alcohol intoxication. *Anaesthesia* 55:1189-1191.

Kanthasamy AG, Ardelt B, Malave A, Mills EM, Powley TL, Borowitz JL, Isom GE. 1997. Reactive oxygen species generated by cyanide mediate toxicity in rat pheochromocytoma cells. *Toxicol Lett* 93:47-54.

Kaplan JA, Finlayson DC, Woodward S. 1980. Vasodilator therapy after cardiac surgery: A review of the efficacy and toxicity of nitroglycerin and nitroprusside. *Can Anaesth Soc J* 27:254-259.

Kaplan JA. 1981. In reply: Cyanide antagonist. *Can Anaesth Soc J* 28:290 [Comment on Vesey and Cole, 1981].

Kasamo K, Okuhata Y, Satoh R, Ikeda M, Takahashi S, Kamata R, Nogami Y, Kojima T. 1993. Chronological changes of MRI findings on striatal damage after acute cyanide intoxication: pathogenesis of the damage and its selectivity, and prevention for neurological sequelae: a case report. *Eur Arch Psychiatry Clin Neurosci* 243:71-74.

Katzman GM, Penney DG. 1993. Electrocardiographic responses to carbon-monoxide and cyanide in the conscious rat. *Toxicol Lett* 69:139-153.

Kawalek JC, Andrews AW. 1980. The effect of solvents on drug metabolism *in vitro*. *Drug Metab Dispos* 8:380-384.

Kealy P. 2009. Carbon Monoxide Toxicity. *J Burn Care Res* 30:146-155.

Kedderis GL, Batra R. 1993. Species differences in the hydrolysis of 2-cyanoethylene oxide, the epoxide metabolite of acrylonitrile. *Carcinogenesis* 14:685-689.

Kedderis GL. 1990. The role of the mixed-function oxidase system in the toxication and detoxication of chemicals: Relationship to chemical interactions. In Goldstein RS, Hewitt WR, Hook JB, eds, *Toxic Interactions*. Academic Press, San Diego, CA, USA, pp 31-60.

Keim ME. 2006. Terrorism involving cyanide: the prospect of improving preparedness in the prehospital setting. *Prehosp Disaster Med* 21:s56-60.

Kerger H, Dodidou P, Passani-Kruppa D, Grüttner J, Birmelin M, Volz A, Waschke KF. 2005. Excessive methaemoglobinaemia and multi-organ failure following 4-DMAP antidote therapy. *Resuscitation* 66:231-235.

Kessler DL Hines LE. 1948. Hazards of thiocyanate therapy in hypertension. *J Am Med Assoc* 138:549-551 [Other poisoning].

Khandekar JD, Edelman H. 1979. Studies of amygdalin(laetril) toxicity in rodents. *JAMA* 242:169-171.

Kiese M, Szinicz L, Thiel N, Weger N. 1975. Ferrihemoglobin and kidney lesions in rats produced by 4-aminophenol or 4-dimethylaminophenol. *Arch Toxicol* 34:337-340.

Kiese M, Weger N. 1965. The treatment of experimental cyanide poisoning by hemoglobin formation. *Archiv für Toxikologie* 21:89-100.

Kiese M, Weger N. 1969. Formation of ferrihaemoglobin with aminophenols in the human for the treatment of cyanide poisoning. *Eur J Pharmacol* 7:97-105.

Kimura ET, Ebert DM, Dodge PW. 1971. Acute toxicity and limits of solvent residue for sixteen organic solvents. *Toxicol. Appl. Pharmacol* 19: 699-704.

KimYH., Foo M, Terry RD. 1982. Cyanide encephalopathy following therapy with sodium nitroprusside: Report of a case. *Arch Pathol Lab Med*, 106:392-393.

Kirk M, Kulig K, Rumack BH. 1989. Methemoglobin and cyanide kinetics in smoke inhalation. *Vet Hum Toxicol* 31:353 [Abstract].

Kirk MA, Gerace R, Kulig KW. 1993. Cyanide and methemoglobin kinetics in smoke inhalation victims treated with the cyanide antidote kit. *Ann Emerg Med* 22:1413-1418.

Kirk MA, Gerace R, Kulig KW. 1994. In reply: Use of sodium nitrite needs further investigation. *Ann Emerg Med*, 24:540-541.

Kirk-Othmer. 1969. *Encyclopedia of chemical technology*, 2nd ed. New York, John Wiley & Sons, Vol. 20.

Kirk-Othmer. 1978. *Kirk-Othmer Encyclopedia of Chemical Technology*. 3rd Edition. John Wiley and Sons, NY.

Kirk-Othmer. 1993. Cyanides. In *Kirk-Othmer Encyclopedia of Chemical Technology*, 4th ed, vol 7. John Wiley and Sons, New York, USA, 1-18.

Kläui H, Russi E, Baumann PC. 1984. Cyanid-Intoxikation. *Schweiz med Wschr* 114:983-989.

Klimek J, Boguslawski W, Zelewski L. 1979. The relationship between energy generation and cholesterol side-chain cleavage reaction in the mitochondria from human term placenta. *Biochim Biophys Acta* 587:362-372.

Klimmek R, Fladerer H, Szinicz L, Weger N, Kiese M. 1979a. Effects of 4-dimethylaminophenol and Co₂EDTA on circulation, respiration, and blood homeostasis in dogs. *Arch Toxicol* 42:75-84.

Klimmek R, Fladerer H, Weger N. 1979b. Circulation, respiration, and blood homeostasis in cyanide-poisoned dogs after treatment with 4-dimethylaminophenol or cobalt compounds. *Arch Toxicol* 43:121-133.

Klimmek R, Krettek C, Szinicz L, Eyer P, Weger N. 1983. Effects and biotransformation of 4-dimethylaminophenol in man and dog. *Arch Toxicol* 53:275-288.

Klimmek R, Krettek C, Werner HW. 1988. Ferrihaemoglobin formation by amyl nitrite and sodium nitrite in different species *in vivo* and *in vitro*. *Arch Toxicol* 62:152-160.

Klimmek R, Krettek C. 1988. Effects of amyl nitrite on circulation, respiration and blood homeostasis in cyanide poisoning. *Arch Toxicol* 62:161-166.

Klimmek R, Roddewig C, Fladerer H, Krettek C, Weger N. 1983. Effects of 4-dimethylaminophenol, Co₂EDTA, or NaNO₂ on cerebral blood flow and sinus blood homeostasis of dogs in connection with acute cyanide poisoning. *Toxicology* 6:143-154.

Klimmek R, Roddewig C, Weger N. 1978. The effect of 4-dimethylaminophenol on acid-base balance, lactate, pyruvate, local cerebral blood flow and brain temperature of dogs after acute cyanide poisoning. Abstracts of the 19th spring meeting March 14–17, 1978, Mainz, Germany. Naunyn-Schmiedeberg's *Arch Pharmacol* 302 R14.

Klimmek R, Roddewig C, Weger N. 1981. Effects of 4-dimethylaminophenol on blood flow and blood gases in the brain. *Res Exp Med (Berl)* 179:141-151.

Klimmek R, Szinicz L, Weger N. 1993. Blausäure und Chlorcyan. In Klimmek R, Szinicz L, Weger N, eds. *Chemische Gifte und Kampfstoffe – Wirkung und Therapie*. Hippokrates, Stuttgart, Germany, 91-102

Klöver E, Wenderoth H. 1965. Die Blausäurevergiftung durch bittere Mandeln in psychologischer und therapeutischer Sicht. *Med Klin* 60:213-216. FB 1950-1966 CN cas Clinique

Knapp W, Hammer L. 1959. [Poisoning of drinking water by cyanide-contaminated waste water.]. *Arch Hyg Bakteriol* 143:501-516 [German]. FB 1950-1966 CN cas Clinique

Köhler Chemie, Electronic Phrama-medical Encyclopedia Online, 2013

Kong A, Shen A, Burrows G, Sylvester D, Ison GE, Way JL. 1983. Effect of chlorpromazine on cyanide intoxication. *Toxicol Appl Pharmacol* 71:407-413.4.7.1

Kontokolias JS; Schulz F; Hildebrandt E; Sautter R. 1993. Die Vergiftung durch Rauchgase - eine Übersicht. *Rettungsdienst* 16:697-705 DP

Kornfeld S, Benziman M, Milner Y. 1978. Regulatory properties of the alpha-ketoglutarate dehydrogenase complex of *Acetobacter xylinum*. In situ studies and localization of the allosteric response in the E1 component. *J Biol Chem* 253:5678-5684.

Koschel MJ. 2006. Management of the cyanide-poisoned patient. *J Emerg Nurs* 32:S19-26. Review. No abstract available.

Krapez JR, Vesey CJ, Adams L, Cole PV. 1981. Effects of cyanide antidotes used with sodium nitroprusside infusions: Sodium thiosulfate and hydroxocobalamin given prophylactically to dogs. *Br J Anaesth* 53:793-804.

Kravzov J, Ríos C, Altagracia M, Jiménez D. 1994. Comparison of cobalt edetate and α -ketoglutarate as antidotes against cyanide intoxication in mice. *Proc West Pharmacol Soc* 37:43-44.

Krefft S. 1955. Vergiftungen durch Acetoncyanhydrin bei Mensch und Tier: Experimentelle Untersuchungen zur percutanen Toxizität des Acetoncyanhydrins. *Arch Gewerbepath Gewerbehyg* 14:110-116.

Krieg A, Saxena K. 1987. Cyanide poisoning from metal cleaning solutions. *Ann Emerg Med* 16:582-584.

Krishna L; Katoch RC. 1987. Investigation of 'mysterious' disease in livestock: hydrocyanic acid poisoning. *Pediatrics* 80:359-363.

Krivanek ND, McLaughlin M, Sprout WL, Trochimowicz HJ. 1984. Studies of possible acrylonitrile poisoning. *Texas Med* 80:6.

Kruszyna H, Kruszyna R, Smith RP. 1985. Cyanide and sulfide interact with nitrogenous compounds to influence the relaxation of various smooth muscles. *Proc Soc Exp Biol Med* 179:44-49.

Kruszyna R, Kruszyna H, Smith RP. 1982. Comparison of hydroxylamine, 4-dimethylaminophenol and nitrite protection against cyanide poisoning in mice. *Arch Toxicol* 49:191-202.

Kulig K. 1991. Cyanide antidotes and fire toxicology. *New Engl J Med* 325:1801-1802.

Kulling P, Personne M. 1997. Fortydligande om antidotbehandling vid cyanidförgiftning. *Läkartidningen* 46:4212.

Kulling P, Persson H. 1997. Förgiftningar. Behandlingsanvisningar. In *Läkemedelsboken 97/98*. Apoteksbolaget AB, Stockholm, Sweden, pp 929-956.

Kunz DA, Chen JL, Pan G. 1998. Accumulation of alpha-keto acids as essential components in cyanide assimilation by *Pseudomonas fluorescens* NCIMB 11764. *Appl Environ Microbiol* 64:4452-4459.

- Kupferschmidt H. 2006. Nitrite (Amylnitrit, Natriumnitrit). Antidot-Monographie: Amylnitrit/Natriumnitrit. Schweizerisches Toxikologisches Informationszentrum, Zürich, Switzerland.
- Küpper U. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 9 cyanide cases 1977-2008]. Personal communication. Evonik Degussa, Wesseling, Germany.
- Kurt TL, Day LC, Reed WG, Gandy W. 1991. Cyanide poisoning from glue-on nail remover. *Am J Emerg Med* 9:271-272.
- Kurt TL, Day LC, Reed WG, Gandy W. 1991. Cyanide poisoning from glue-on nail remover. *Am. J. Emerg. Med.* 9:271-272.
- Lam KK, Lau FL. 2000. An incident of hydrogen cyanide poisoning. *Am J Emerg Med* 18:172-175.
- Lambert W, Meyer E et al. 1995. Cyanide and sodium azide intoxication. *Ann Emerg Med* 26:392.
- Lang S. 1895. Studien über Entgiftungstherapie I: Über Entgiftung der Blausäure. *Naunyn-Schmiedebergs Arch Exp Pathol Pharmacol* 36:75-99.
- Lang J, Stintzy F. 1960. Un cas d'intoxication lente à l'acide cyanhydrique par l'acétone-cyanhydrine. *Arch Mal Prof* 21:652-657.
- Laroche-Navarro L. Département Médical, KELOCYANOR.
- Lascelles J, Calder KM. 1985. Participation of cytochromes in some oxidation-reduction systems in *Campylobacter fetus*. *J Bacteriol* 164:401-409.
- Lasch EE, El Shawa R. 1981. Multiple cases of cyanide poisoning by apricot kernels in children from Gaza. *Pediatrics* 68:5-7.
- Lau FL. 2000. Emergency management of poisoning in Hong Kong. *Hong Kong Med J* 6:288-292.
- Lawson-Smith P, Jansen EC, Hilsted L, Hydegaard O. 2010. Effect of hyperbaric oxygen therapy on whole blood cyanide concentrations in carbon monoxide intoxicated patients from fire accidents. *Scand J Trauma Resusc Emerg Med* 18:32-36.
- Lazarus-Barlow P, Norman GM. 1941. Fatal cases of poisoning with sodium nitroprussid. *Br Med J* 2:407.

Lee CG, Webber TD. 1983. The mutagenicity of a cyanide antidote, dimethylaminophenol, in Chinese hamster cells. *Toxicol Lett* 16:85-88.

Lee J, Mukai D, Kreuter K, Mahon S, Tromberg B, Brenner M. 2007. Potential interference by hydroxocobalamin on cooximetry hemoglobin measurements during cyanide and smoke inhalation treatments. *Ann Emerg Med* 49:802-805

Lee M, Berger HW, Givre HL, Jayamanne DS. 1982. Near fatal laetrile intoxication: Complete recovery with supportive treatment. *Mount Sinai J Med* 40:305-307.

Lee-Jones M, Bennett MA, Sherwell JM. 1970. Cyanide self-poisoning. *Br Med J* 4:780-781.

Leng G, Lewalter J. 2001. Polymorphism of glutathione s-transferases and susceptibility of acrylonitrile and dimethylsulfate in cases of intoxication. *Toxicol Lett* 134:209-217.

Leor R, Michaeli J, Brezis M, Stessman J. 1986. Laetrile intoxication and hepatic necrosis: a possible association. *South Med J* 79:259-260.

Leeuwenkamp OR, van Bennekom WP, van der Mark EJ, Bult A. 1984. Nitroprusside, antihypertensive drug and analytical reagent. Review of (photo)stability, pharmacology and analytical properties. *Pharm Weekbl Sci* 6(4):129-140.

Levillain P. 1997. L'intoxication cyanhydrique: ses aspects fondamentaux et cliniques. Deuxième Journée de Toxicologie de l'Hôpital Fernand Widai, Paris, France.

Levin BC, Gurman JL, Paabo M, Baier L, Procell L, Newball HH. 1986. Acute inhalation toxicity of hydrogen cyanide. *The Toxicologist* 6:59.

Levine S. 1959. Oxygen in the therapy of cyanide poisoning. *JAMA* 170(13):1585.

Lewalter J. 1991. Empfehlungen zur arbeitsmedizinischen Prävention expositions- und dispositionsbedingter Arbeitsstoffbelastungen. *Verh.Dtsch.Ges.Arbeitsmed* 31:135-139.

Liebowitz D, Schwartz H. 1948. Cyanide poisoning – report of a case with recovery. *Am J Clin Pathol* 18:965-970.

Liegner KB, Beck EM, Rosenberg A. 1981. Laetrile-induced agranulocytosis. *JAMA* 246:2841-2842.

Linakis JG, Lacouture pg, Woolf A. 1991. Monitoring cyanide and thiocyanate concentrations during infusion of sodium nitroprusside in children. *Pediatr Cardio*, 12:214-218.

Linnell JC. 1987. The role of cobalamin in cyanide detoxification. In Ballantyne B, Marrs T, eds. *Clinical and experimental toxicology of cyanides*. John Wright, Bristol, England, UK, pp 427-439.

Lipczak H, Jensen K. 1998. Cyanide poisoning during exposure to smoke from fires. (Article in Danish) *Cyanidforgiftning ved udsættelse for brandrog*. *Ugeskr Laeger* 160:6369-6371.

Lipha Santé. 1996. L'intoxication par les cyanures et son traitement. Cyanokit 2,5 g hydroxocobalamine. Lipha Santé, Lyon, France. Lipha, Pharmaceuticals Ltd, Dicobalt Edetate (Kelocyanor).

Litovitz TL. 1987. The use of oxygen in the treatment of acute cyanide poisoning. In Ballantyne B, Marrs TC, *Clinical and Experimental Toxicology of Cyanides*. John Wright, Bristol, UK.

Litovitz TL, Larkin RF, Myers RA. 1983. Cyanide poisoning treated with hyperbaric oxygen. *Am J Emerg Med* 1:94-101.

Lloyd NL. 1957. A new kit for sodium nitrite-thiosulphate therapy in the treatment of acute cyanide poisoning. *Br J Ind Med* 14:137. FB 1950-1966 CN cas clinique

Logan BK. 1996. Improved postmortem detection of carbon monoxide and cyanide. A summary of a research study. Toxicology Laboratory, University of Washington DC. National Institute of Justice Research Preview. US Department of Justice, Washington, DC, USA.

Lohs K, Taufmann M, Kühn G, Krüger S. 1974. Weitere Untersuchungen zur Cyanidentgiftung durch DL-Glycerinaldehyd. *Pharmazie* 29:292.

Lorz H. 1950. Über perkutane Vergiftung mit Akrylnitril (Ventox). *Dtsch Med Wschr* 75:1987-1088.

Losek JD, Rock AL, Boldt RR. 1991. Cyanide poisoning from a cosmetic nail remover. *Pediatrics* 88:337-340.

Löser E. 1994. Organische Verbindungen: Cyanverbindungen. In Gloxhuber C, ed, *Toxikologie*, 5th ed. Georg Thieme Verlag Stuttgart, New York, USA pp 183-192.

Lowry WT, Juarez L, Petty CS, Roberts B. 1985. Studies of toxic gas production during actual structural fires in the Dallas area. *J Forensic Sci* 30:59-72 [Experimental study].

Ludwig E, Eyer P. 1994. Structural requirements for the ferrihemoglobin-forming activity of glutathione s-conjugates of 4-dimethylaminophenol *Environ Health Perspect* 6:133-136

Ludwig E, Eyer P. 1995. Oxidation versus addition reactions of glutathione during the interactions with quinoid thioethers of 4-(dimethylamino)phenol. *Chem Res Toxicol* 8:302-309

Ludwig E, Eyer P. 1995. Reactivity of glutathione adducts of 4-(dimethylamino)phenol. Formation of a highly reactive cyclization product. *Chem Res Toxicol* 8:310-315

Ludwig E, Eyer P. 1995. Reactivity of glutathione adducts of 4-(dimethylamino)phenol. Involvement of reactive oxygen species during the interaction with oxyhemoglobin. *Chem Res Toxicol* 8:363-368

Lundquist P, Kagedal B, Nilsson L. 1995. An improved method for determination of thiocyanate in plasma and urine. *Eur J Clin Chem Clin Biochem* 33:343-349.

Lundquist P, Rosling H, Sörbo B. 1988. The origin of hydrogen cyanide in breath. *Arch Toxicol* 61:270-274.

Lundquist P. 1992. Determination of cyanide and thiocyanate in humans. PhD thesis. Faculty of Medicine. Linköping University, Linköping, Sweden.

Lurie I. 1953. ACTH treatment in a case of cyanide poisoning. *Acta Med Orient* 12:87-88.

Lutier F, Dusoleil P, De Montgros J. 1971. Action de l'hydroxocobalamine à dose massive dans l'intoxication aiguë au cyanure: A propos d'un cas. *Arch Mal Prof* 32:683-686.

Mace SE, Barata IA, Cravero JP, Dalsey WC, Godwin SA, Kennedy RM, Malley KC, Moss RL, Sacchetti AD, Warden CR, Wears RL; American College of Emergency Physicians. 2004. Clinical policy: Evidence-based approach to pharmacologic agents used in pediatric sedation and analgesia in the emergency department. *Ann Emerg Med* 44:342-377.

MacRae WR, Owen M. 1974. Severe metabolic acidosis following hypotension induced with sodium nitroprusside. *Br J Anaesth*, 46:795-797.

Maechler P, Antinozzi PA, Wollheim CB. 2000. Modulation of glutamate generation in mitochondria affects hormone secretion in INS-1E beta cells. *IUBMB Life* 50:27-31.

Mannaioni G, Vannacci A, Marzocca C, Zorn AM, Peruzzi S, Moroni F. 2002. Acute cyanide intoxication treated with a combination of hydroxocobalamine, sodium nitrite, and sodium thiosulfate. *J Toxicol Clin Toxicol* 40:181-183.

MarburyTC, Sheppard JE, Gibbons K, Lee CSC. 1982. Combined antidotal and hemodialysis treatments for nitroprusside-induced cyanide toxicity. *J Toxicol Clin Toxicol* 19:475-482.

Marrs TC, Bright JE, Swanston DW. 1982. The effect of prior treatment with 4-dimethylaminophenol (DMAP) on animals experimentally poisoned with hydrogen cyanide. *Arch Toxicol* 51:247-253.

Marrs TC, Bright JE, Woodman AC. 1987. Species differences in methaemoglobin production after addition of 4 dimethylaminophenol, a cyanide antidote, to blood in vitro: A comparative study. *Comp Biochem Physiol B* 86:141-148. ,

Marrs TC, Bright JE. 1987. Effect on blood and plasma cyanide levels and on methaemoglobin levels of cyanide administered with and without previous protection using PAPP. *Hum Toxicol* 6:139-145.

Marrs TC, Scawin J, Swanston DW. 1984. The acute intravenous and oral toxicity in mice, rats and guinea-pigs of 4-dimethylaminophenol (DMAP) and its effects on haematological variables. *Toxicology* 31:165-173

Marrs TC. 1987. Dicobalt edetate (Kelocyanor).

Marrs TC. 1987. The choice of cyanide antidotes. In Ballantyne B, Marrs T, eds. *Clinical and experimental toxicology of cyanides*. John Wright, Bristol, England, UK 383-401.

Marrs TC. 1988. Antidotal treatment of acute cyanide poisoning. *Adverse Drug React Acute Poisoning Rev* 7:179-206.

Martin C. 1987. Antidotes: certains agissent par compétition avec le toxique au niveau du récepteur. *Panorama du Médecin*.

Martin K, Shaw TI. 1970. The production of adenosine triphosphate in perfused giant axons of *Loligo*. *J Physiol* 208:171-185.

Martín-Bermúdez R, Maestre-Romero A, Goñi-Belzunegui MV, Bautista-Lorite A, Arenas-Cabrera C. 1997. Venous blood ateriolization and multiple organ failure after cyanide poisoning. *Intensive Care Med* 23:1286-1289.

Martindale. 1989a. Sodium nitrite. In Reynolds JEF, ed, *The Extra Pharmacopoeia*, 29th ed. Pharmaceutical Press, London, UK, p 855.

Martindale. 1989b. Dicobalt edetate. In Reynolds JEF, ed, *The extra pharmacopoeia*, 29th ed. Pharmaceutical Press, London, UK, p 839.

Martindale. 2008. *The complete drug reference (Micromedex® Healthcare Series Databases)*. Thomson Micromedex (a part of Thomson Reuters), New York, USA, pp 1974-2008.

Martinelli CJ, Barko IR, O'Toole K, Bayer M. 2008. "Vitamin" B17 toxicity treated with hydroxocobalamin. *Clin Toxicol* 46:631 [Abstract].

Mascarenhas BR, Geller AC, Goodman AI. 1969. Cyanide poisoning: medical emergency. *NY State J Med* 60:1782-1784.

Mathangi DC, Namasivayam A. 2000. Neurochemical and behavioural correlates in cassava-induced neurotoxicity in rats. *Neurotox Res*, 2:29-35

Mathangi DC, Shyamala R, Vijayashree R, Rao KR, Ruckmani A, Vijayaraghavan R, Bhattacharya R. 2011. Effect of alpha-ketoglutarate on neurobehavioral, neurochemical and oxidative changes caused by sub-chronic cyanide poisoning in rats. *Neurochem Res* 36:540-548

Mathes GV, von Clarmann M. 1977. The index-line as a new aid in the clinical and forensic toxicology. *Acta Pharmacol Toxicol (Copenhagen)*, 41:488-497.

Matijak-Schaper M, Alarie Y. 1982. Toxicity of carbon monoxide, hydrogen cyanide and low oxygen. *J Combust Toxicol* 9:21-61 [Cited by US-NAC, 2000a].

Matthews DM, Wilson J. 1970. Cobalamins and cyanide metabolism in neurological diseases. In Arnstein HRV, Wrighton RJ, eds. *The cobalamins: A Glaxo Symposium*. Churchill Livingstone, Edinburgh, UK 115-136.

Maxwell DM, 1978. Increased cyanide values in a laetrile user. *Can Med Assoc J* 119:18.

McAnalley BH, Gardiner TH, Garriott JC. 1980. Cyanide concentrations in blood after amygdalin (Laetrile) administration in rats. *Vet Hum Toxicol* 22:400-402.

McCormack, JG, Bromidge ES, Dawes NJ. 1988. Characterization of the effects of Ca²⁺ on the intramitochondrial Ca²⁺-sensitive dehydrogenases within intact rat-kidney mitochondria. *Biochim Biophys Acta* 934:282-292.

McEvoy, Gerald K et al. 2008. AHFS drug information 2008. American hospital formulary service - American Society of Health-System Pharmacists, Bethesda, MD, USA, p 1800.

McGaughey C. 1951. Hydrocyanic acid poisoning in nutrias caused by cassava (*Manihot utilissima* Pohl). *Br Vet J* 107:279-280.

McGuinn WD, Baxter L, Petrikovics I, Cannon EP, Way JL. 1994. Antagonism of the lethal effects of cyanide by a synthetic water-soluble cobalt(III) porphyrin compound. *Fundam Appl Toxicol* 23:76-80.

McKiernan MJ. 1980. Emergency treatment of cyanide poisoning. *Lancet* 2:86.

- McLaughlin M, Krivanek ND, Trochimowicz HJ, Sprout WL. 1985. Studies of possible acrylonitrile poisoning antidote. *Texas Med* 81:6.
- Mégarbane B, Borron SW, Baud F. 2002. Hydroxocobalamin versus conventional treatment in cyanide poisoning. *Clin Toxicol* 40:314-315.
- Mégarbane B, Delahaye A, Goldgran-Tolédano D, Baud FJ. 2003. Antidotal treatment of cyanide poisoning. *J Chin Med Assoc* 66:193-203.
- Mehta C. 1995. Antidotal effect of sodium thiosulfate in mice exposed to acrylonitrile. *Res Comm Mol Pathol Pharmacol* 87:155-165.
- Mehta R, Shangari N, O'Brien PJ. 2008. Preventing cell death induced by carbonyl stress, oxidative stress or mitochondrial toxins with vitamin B anti-AGE agents. *Mol Nutr Food Res* 52:379-385.
- Meigs RA, Sheean LA. 1977. Mitochondria from human term placenta. III: The role of respiration and energy generation in progesterone biosynthesis. *Biochim Biophys Acta* 489:225-235.
- Mellino M, Phillips DF. 1980. Severe lactic acidosis in a case of nitroprusside resistance. *Cleveland Clinic Quarterly* 47:119-122.
- Melloni C, Corticelli AS, Di Nino GF, Mazzetti P, Torti D. 1983. Un caso di avvelenamento da cianuri [A case of cyanide poisoning]. *Minerva Anestesiol* 49:527-530.
- Mengel K, Krämer W, Isert B, Friedberg KD. 1989. Thiosulphate and hydroxocobalamin prophylaxis in progressive cyanide poisoning in guinea-pigs. *Toxicology* 54:335-342.
- Merck. 2006. Cyanokit (hydroxocobalamin for injection) 5 g for intravenous infusion. Merck Santé, Semoy, France.
- Merck. 2007. Cyanide disease backgrounder: Acute cyanide poisoning in the United States – The scope of the problem.
- Merck Serono. 2009. Cyanokit 2.5 g in practice. Merck Santé, Lyon, France.
- Merck Serono. 2010. New Presentation of Cyanokit approved in Europe. Merck Serono, Geneva, Switzerland.
- Mercker H, Bastian G. 1959. Kobaltverbindungen zur Entgiftung von Blausäure. *Naunyn-Schmiedeberg Arch Exp Pathol Pharmacol* 236:449-458.

Mercker H, Lochner W, Gerstenberg E. 1957. Arterielle und venöse Sauerstoffsättigungen bei Cyanidvergiftung. Naunyn Schmiedebergs Arch Exp Pathol Pharmacol 232:289-290.

Mercker H, Lochner W, Gerstenberg E. 1958. Untersuchungen über das Verhalten des Kreislaufes, der Atmung und der Sauerstoffsättigung des Blutes bei Cyanidvergiftung. Naunyn Schmiedebergs Arch Exp Pathol Pharmacol 232(2):459-469.

Mercker H; Roser F. 1957. Über Kreislauf- und Stoffwechselreaktionen bei der spezifischen Behandlung der Blausäurevergiftung. Arch exper Pathol Pharmacol 230:125-141.

Meredith TJ. 1993. Antidotes for poisoning by cyanide. In IPCS, Antidotes for poisoning by cyanide. Meredith TJ, Jacobsen D, Haines JA, Berger J-C, van Heijst ANP, eds, International Programme on Chemical Safety, Commission of the European Communities, Evaluation of antidotes series 2. EUR 14280 EN. Cambridge University Press, Cambridge, UK [www.inchem.org/pages/antidote.html], pp. 1-26.

Merkel JR, Sipos T. 1971. Marine bacterial proteases. I: Characterization of an endopeptidase produced by *Vibrio B-30*. Arch Biochem Biophys 145:126-36.

Merrifield AJ, Blundell MD. 1974. Toxicity of sodium nitroprusside. Br J Anaesth 46:324.

Messing B 1991. Extrapryridal disturbances after cyanide poisoning (first MRI-investigations of the al transm, 33, 141-147.

Meyer GW, Hart GB, Strauss MB. 1991. Hyperbaric oxygen therapy for acute smoke inhalation injuries. Postgrad Med 89(1):221-223.

Michaelis HC, Clemens C, Kijewski H, Neurath H, Eggert A. 1991. Acetonitrile serum concentrations and cyanide blood levels in a case of suicidal oral acetonitrile ingestion. J Toxicol Clin Toxicol 29:447-458.

Michenfelder JD, Tinker JH. 1977. Cyanide toxicity and thiosulfate protection during chronic administration of sodium nitroprusside in the dog: Correlation with a human case. Anesthesiology 47:441-448.

Milazzo S, Lejeune S, Ernst E. 2007. Laetrile for cancer: a systematic review of the evidence. Support Care Cancer 15:583-595.

Miller MH, Toops TC. 1951. Acute cyanide poisoning; recovery with sodium thiosulfate therapy. J Indiana State Med Assoc 44:1164.

Mirkin D B. 1989. Signs and symptoms of cyanide and acrylonitrile poisoning.

Mittal G, Singh T, Kumar N, Bhatnagar A, Tripathi RP, Tulsawani R, Vijaraghavan R, Bhattacharya R. 2010. Radiolabeling and dose fixation study of oral alpha-ketoglutarate as a cyanide antidote in healthy human volunteers. *Clin Toxicol* 48:509-515.

Mizoule J. 1966. Étude de l'action antidote de l'hydroxocobalamine à l'égard de l'intoxication cyanhydrique. PhD thesis. Université de Paris, Paris, France.

Mjörndal T, Norberg B. 1993. Management of vitamin B12 deficiency in outpatient care. *Scand J Prim Health Care* 11:1-2.

Mlingi N, Poulter NH, Rosling H. 1992. An outbreak of acute intoxications from consumption of insufficiently processed cassava in Tanzania. *Nutr Res* 12:677-687.

Moertel CG, Fleming TR, Rubin J, Kvols LK, Sarna G, Koch R, Currie VE, Young CW, Jones SE, Davignon JP. 1982. A clinical trial of amygdalin (laetrile) in the treatment of human cancer. *N Engl J Med* 306:201-206.

Moeschlin S. 1980. Cyanwasserstoff (Blausäure, HCN). In *Klinik und Therapie der Vergiftungen*, 6th ed. Thieme, Stuttgart, Germany, pp 252-257.

Mofenson HC, Greensher J, Horowitz R, Berlin CM. 1971. Treatment of cyanide poisoning. *Pediatrics* 47:1093.

Moffett BS, Price JF. 2008. Evaluation of sodiumnitroprusside toxicity in pediatric cardiac surgical patients. *Ann Pharmacother* 42:1600-1604.

Mohler SR. 1975. Air crash survival: Injuries and evacuation toxic hazards. *Aviat Space Environ Med* 46:86-88.

Moldovanyi IS, Denes G. 1968. Mechanism of the action and of the allosteric inhibition of 3-deoxy-d-arabino-heptulosonate 7-phosphate synthase (tyrosine-sensitive) of *Escherichia coli* W. *Acta Biochim Biophys Acad Sci Hung* 3:259-73.

Montero Pérez FJM. 1998. La hidroxocobalamina en el tratamiento de la intoxicación conjunta por monóxido de carbono y cianuro. *Med Clin (Barc)* 110:558-559.

Montgomery RD. 1965. The medical significance of cyanogen in plant foodstuffs. *Am J Clin Nutr* 17:103-113.

Montoliu J, Botey A, Pons JM, Revert L. 1979. Fatal hypotension in normal-dose nitroprusside therapy. *Am Heart J* 97:541-542.

Moore DGF. 1934. Manioc (Kassava) as a native food in Nigeria. *W Afr Med J* 7:97-98 [With notes by Oluwole IL *ibidem* p 98].

Moore DGF. 1934. Retrobulbar neuritis and partial optic atrophy as sequelae of avitaminosis. *Ann Trop Med Parasit* 28:295-303.

Moore SJ, Ho IK, Hume AS. 1991. Severe hypoxia by concomitant intoxication with sublethal doses of carbon monoxide and cyanide. *Toxicol Appl Pharmacol* 109:412-420.

Moore SJ, Norris JC, Ho IK, Hume AS. 1986. The efficacy of α -ketoglutaric acid in the antagonism of cyanide intoxication. *Toxicol Appl Pharmacol* 82:40-44.

Moore SJ, Norris JC, Walsh DA, Hume AS. 1987. Antidotal use of methemoglobin forming cyanide antagonists in concurrent carbon monoxide/cyanide intoxication. *J Pharmacol Exp Ther* 242:70-73 [Experimental study in mice].

Moreau F, Romani R. 1982. Malate oxidation and cyanide-insensitive respiration in avocado mitochondria during the climacteric cycle. *Plant Physiol* 70:1385-1390.

Morse DL, Boros L, Findley PA. 1979. More on cyanide poisoning from laetrile. *N Engl J Med* 301:892.

Moss M, Khalil N, Gray J. 1981. Deliberate self-poisoning with laetrile. *Can Med Assoc J* 125:1126-1128.

Moss RH, Jackson CF, Seiberlich J. 1951. Toxicity of carbon monoxide and hydrogen cyanide gas mixtures. A preliminary report. *Arch Ind Hyg Occup Med* 4:53-64.

Mota MM. 1933. [On a case of intoxication by potassium cyanide treated with sodium nitrite with survival.] *Rev Med Rosario* 23:670-674 [Spanish].

Motin J, Bouletreau P, Rouzioux JM. 1970. Intoxication cyanhydrique grave traitée avec succès par hydroxocobalamine. *J Méd Strasbourg* 1:717-722.

Moureau P, Heusghem C, Warin J, Delree H. 1953. Intoxication massive par l'acide cyanhydrique dans une installation industrielle de lavage du benzol. *Acta Med Leg Soc (Liège)* 6:257-262.

Mousa HM, Davis RH. 1991. Alternative sulphur donors for detoxification of cyanide in the chicken. *Comp Biochem Physiol* 99:309-315.

MPI Research. 1997a. A dermal irritation study of acetonitrile in rabbits.

- MPI Research. 1997b. An Eye Irritation Study of Acetonitrile in Rabbits.
- MPI Research. 1997c. An Acute Dermal Toxicity Study of Acetonitrile in Rabbits.
- MPI Research. 1998a. An Acute Oral Toxicity Study of Acetonitrile in Mice.
- MPI Research. 1998b. An Acute Inhalation Toxicity Study of Acetonitrile in Mice.
- Mueller M, Borland C. 1997. Delayed cyanide poisoning following acetonitrile ingestion. *Postgrad Med J* 73:299-300.
- Muraki K, Inoue Y, Ohta I, Kondo K, Matayoshi Y, Kamei T. 2001. Massive rhabdomyolysis and acute renal failure after acetonitrile exposure. *Intern Med* 40:936-939.
- Mushett CW, Kelley KL, Boxer GE, Richards JC. 1952. Antidotal efficacy of vitamin B12a (hydroxocobalamin) in experimental cyanide poisoning. *Proc Soc Exp Biol Med* 81:234-237.
- Musil J. 1966. Der Einfluss einer chronischen Natriumnitrit-Intoxikation auf Ratten. *Acta Biol Med Ger* 16:388-394 [Experimental study].
- Musshoff F, Schmidt P, Daldrup T, Madea B. 2002. Cyanide fatalities: Case studies of four suicides and one homicide. *Am J Forensic Med Pathol* 23:315-320.
- Mustea I, Bara A, Petrescu I, Revesz L. 1978. Effect of anoxic radiosensitizers on cellular and mitochondrial oxygen consumption and respiration control ratio. *Br J Cancer Suppl* 3:159-162.
- Mutlu GM, Leiking JB, Oh K, Factor P. 2002. An unresponsive biochemistry professor in the bathtub. *Chest* 122:1073-1076.
- N N. 1977. Blausäure - Antidote bei akuten Vergiftungen. *Fortschr Med* 95:2409-2412.
- N N. 1990. Rauchgasvergiftung - ein Behandlungsschema. *Notfallmedizin* 16:546-548.
- Nadler JE, Green H, Rosenbaum A. 1945. Intravenous injection of methylene blue in man with reference to its toxic symptoms and effect on the electrocardiogram. *Am J Med Sci* 188:15-20.4.7.1
- Nagahara N, Li Q, Sawada N. 2003. Do antidotes for acute cyanide poisoning act on mercaptopyruvate sulfurtransferase to facilitate detoxification? *Curr Drug Targets Immune Endocr Metabol Disord* 3:198-204.
- Nagasawa HT, Goon DJW, Crankshaw DL, Vince R, Patterson SE. 2007. Novel, orally effective cyanide antidotes. *J Med Chem* 50:6462-6464.

Nagler J, Provoost RA, Parizel G. 1978. Hydrogen cyanide poisoning: Treatment with cobalt EDTA. *J Occup Med* 20:414-416.

Nagorzanski R. 1966. Pathogenesis, symptoms, first aid and treatment of acute poisoning with hydrogen cyanide and acrylonitrile. *Medycyna Pracy* 17:5.

Nagorzanski R. 1966. Patogeneza, objawy, pierwsza pomoc i leczenie ostrych zatruc cyjanowodorem i akrylonitrylem [Pathogenesis, symptoms, first aid and treatment of acute poisoning with hydrogen cyanide and acrylonitrile] . *Medycyna Pracy* 17:466-473 [Polish].

Nakatani T, Kosugi Y, Mori A, Tajimi K, Kobayashi K. 1993. Changes in the parameters of oxygen metabolism in a clinical course recovering from potassium cyanide. *Am J Emerg Med* 11:213-217.

Narayanaswami K, Jand S, Kotangale RS, Nath P, Bisaria VS. 1981. Changes in cyanide as a function of storage time in autopsy material by UV spectrometry. *Indian J Med Res*, 73:291-295.

Naughton M. 1974. Acute cyanide poisoning. *Anaesth Intens Care* 2:351-356.

Nemec MD, Kirkpatrick DT, Sherman J, Van Miller JP, Pershing ML & Strother DE. 2008. Two-Generation Reproductive Toxicity Study of Inhaled Acrylonitrile Vapors in Crl: CD(SD) Rats. *International Journal of Toxicology*, 27: 11-29. Testing laboratory: WIL Research Laboratories, LLC, Ohio.

Nerland DE, Benz FW, Babiuk C. 1989. Effects of cysteine isomers and derivatives on acute acrylonitrile toxicity. *Drug Metab Rev* 20:233-246.

Nerland DE, Benz FW, Feldhoff RC, Pierce WM. 1999. Identification of the predominate site of covalent binding of acrylonitrile to rat hemoglobin. 1999 SOT Annual Meeting [Abstract]. .

Nerland DE, Cai J, Benz FW. 2002. Selective covalent binding of acrylonitrile to Cys 186 in rat liver carbonic anhydrase III in vivo. *Chem. Res. Toxicol.* 16:583-589.

Nessim SJ, Richardson RM., 2006. Dialysis for thiocyanate intoxication: a case report and review of the literature. *ASAIO J.*52(4):479-81.

Nicholson PJ, Ferguson-Smith J, Pemberton MA, Campbell A, Edwards JN Ferner RE. 1994. Time to discontinue the use of solutions A and B as a cyanide 'antidote'. *J Occup Med* 44:125-128.

Nicoletta G, Cascelli M, Marchesini L, Tesoro S. 2007. A probable case of nitroprusside intoxication. *Minerva Anestesiol*, 73:471-473.

- Niknahad H, Ghelichkhani E. 2002. Antagonism of cyanide poisoning by dihydroxyacetone. *Toxicol Lett* 132:95-100.
- Niknahad H, Khan S, O'Brien PJ. 1995. Oxygen dependence of hepatocyte susceptibility to mitochondrial respiratory inhibitors. *Biochem Pharmacol* 50:1859-1865.
- Niknahad H, Khan S, Sood C, O'Brien PJ. 1994. Prevention of cyanide-induced cytotoxicity by nutrients in isolated rat hepatocytes. *Toxicol Appl Pharmacol*, 128:271-279
- Niknahad H, O'Brien PJ. 1996. Antidotal effect of dihydroxyacetone against cyanide toxicity in vivo. *Toxicol Appl Pharmacol*, 138:186-191.
- NIOSH. 1986. Register of toxic effects of chemical substances. National Institute for Occupational Safety and Health, Washington, DC, USA.
- Njoh J. 1990. Tropical ataxic neuropathy in Liberians. *Trop Geogr Med*, 42:92-94
- Nolting A, Uhl W, Gallemann D, Hecht S, Kovar A. 2007. Changes in blood pressure after administration of hydroxocobalamin: relationship to changes in plasma cobalamins-(III) concentrations in healthy volunteers. *Clin Toxicol* 45:615.
- Norcross NC. 1938. Intracerebral blood flow: An experimental study. *Arch Neurol Psychiatry (Chicago)* 40:291-299 [Amyl nitrite increases cerebrospinal fluid pressure].
- Norris JC, Moore SJ, Hume AS. 1986. Synergistic lethality induced by the combination of carbon monoxide and cyanide. *Toxicology* 40:121-129.
- Norris JC, Utley WA, Hume AS. 1990. Mechanism of antagonizing cyanide-induced lethality by α -ketoglutaric acid. *Toxicology*, 62:275-283
- O'Brien B, Quigg C, Leong T. 2005. Severe cyanide toxicity from 'vitamin supplements'. *Eur J Emerg Med* 12:257-258.
- O'Brien DJ, Walsh DW, Terriff CM, Hall AH. 2011. Empiric management of cyanide toxicity associated with smoke inhalation. *Prehosp Disaster Med* 5:374-382.
- Obungu VH, Kiaina JK, Njogu RM, Olembo N K. 1999. Catabolism of proline by procyclic culture forms of *Trypanosoma congolense*. *Comp Biochem Physiol B Biochem Mol Biol* 123:59-65.
- Obungu VH, Kiaina JK, Olembo NK, Njogu MR. 1999. Pathways of glucose catabolism in procyclic *Trypanosoma congolense*. *Indian J Biochem Biophys* 36:305-311.

OECD. 2005. Sodium nitrite. SIDS initial assessment report for SIAM 20 (Paris, France, April 2005). Organisation for Economic Co-operation and Development, United Nations Environment Programme. UNEP publications, UNEP Chemicals, Geneva, Switzerland.

Offterdinger H, Weger N. 1969. Kreislauf und Atmung bei Blausäurevergiftung und Therapie mit Ferrihämoglobinbildnern und Kobaltverbindungen. *Naunyn-Schmiedeberg's Arch Pharmacol* 264:289.

O'Flaherty EJ, Thomas WC. 1982. The cardiotoxicity of hydrogen cyanide as a component of polymer pyrolysis smokes. *Toxicol Appl Pharmacol* 63:373-381.

Ohkawa H, Ohkawa R, Yamamoto I, Casida JE. 1972. Enzymatic mechanisms and toxicological significance of hydrogen cyanide liberation from various organothiocyanates and organonitriles in mice and houseflies. *Pestic. Biochem. Physiol.* 2:95-112.

Oi, S. 1974. Inhibition of bovine liver rhodanese by alpha-ketoglutarate. *J Biochem (Tokyo)* 76:455-458.

Okafor PN, Okorowkwo CO, Maduagwu EN. 2002. Occupational and dietary exposures of humans to cyanide poisoning from large-scale cassava processing and ingestion of cassava food. *Food Chem Toxicol* 40:1001-1005

Oke OL. 1980. Toxicity of cyanogenic glycosides. *Food Chem* 6:97-109 [Review].

Olusi SO, Oke OL, Odusote A. 1979. Effect of cyanogenic agents on reproduction and neonatal development of rats. *Biology of the Neonate* 36:233-234.

Oluwole OSA, Onabolu AO, Link H, Rosling H. 2000. Persistence of tropical ataxic neuropathy in a Nigerian community. *J Neurol Neurosurg Psychiatry.* 69:96-101.

Ortega JA, Creek JE. 1978. Acute cyanide poisoning following administration of laetrile enemas. *J Pediatr* 93:1059.

OrušeVT, Jovanović S, Bauer S. 1971. Slučaj akutne profesionalne intoksikacije akrilonitrilom (Un cas d'intoxication aiguë professionnelle par l'acrylonitrile) [Case report of acute occupational poisoning by acrylonitrile]. *Arg Hig Rada*, 23:139-141 [Croatian; Frech abstract].

Orusev T, Jovanović S, Bauer S. 1972. A case of acute occupational intoxication by acrylonitrile. *Arh Hig Rada* 23:139-141 [Croatian].

Osuntokun BO, Langman MJ, Wilson J, Aladetoyinbo A. 1970. Controlled trial of hydroxocobalamin and riboflavine in Nigerian ataxic neuropathy. *J Neurol Neurosurg Psychiat* 33:663-666.

Osuntokun BO, Monekosso GL. 1969. Degenerative tropical neuropathy and diet. *Br Med J* 3:178-179.

Osuntokun,B.O., G.L.Monekosso, J.Wilson. 1969. Relationship of a degenerative tropical neuropathy to diet: Report of a field survey. *Br Med J* 1(5643):547-550.

Padmaja G. 1995. Cyanide detoxification in cassava for food and feed use. *Crit Rev Food Sci Nutr* 35:299-339.

Page IH, Corcoran AC, Dustan HP, Koppanyi T. 1955. Cardiovascular actions of sodium nitroprusside in animals and hypertensive patients. *Circulation* 11(2):188-198.

Pahl MV, Vaziri ND. 1982. In-vivo and in-vitro hemodialysis studies of thiocyanate. *J Toxicol Clin Toxicol.* 19(9):965-74.

Palcic B, Skarsgard LD. 1975. Absence of ultrafast processes of repair of single-strand breaks in mammalian DNA. *Int J Radiat Biol Relat Stud Phys Chem Med* 27:121-33.

Pallapies D, Klees J, eds. 1999. Cyanides. Chemical Emergency Medical Guidelines. BASF AG, Ludwigshafen, Germany.

Pamidi PV, DeAbreu M, Kim D, Mansouri S. 2009. Hydroxocobalamin and cyanocobalamin interference on co-oximetry based hemoglobin measurements. *Clin Chim Acta* 401:63-67.

Pardee GK. 1847. Case of poisoning by the wild cherry. In: Lawson LM. ed. *The Western Lancet* vol 6. Robinson and Jones, Cincinnati, Ohio, USA 289-291.

Patel CB, Laboy V, Venus B, Mathru M, Wier D. 1986. Use of sodium nitroprusside in post-coronary bypass surgery: A plea for conservatism. *Chest* 89:663-667.

Paulet G. 1955a. De l'importance relative de la défaillance respiratoire et de la défaillance cardiaque dans l'action létale des cyanures. *Arch Int Physiol Biochim* 63:328-339.

Paulet G. 1955b. Valeur et mécanisme d'action de l'oxygénothérapie dans le traitement de l'intoxication cyanhydrique. *Arch Int Physiol Biochim* 63:340-360. FB 1950-1966 CN cas clinique, IPCS 2.14

Paulet G. 1957. Valeur des sels organiques du cobalt dans le traitement de l'intoxication cyanhydrique. *C R Séances Soc Biol* 151:1932-1935. IPCS 5.14.

Paulet G. 1958. Intoxication cyanhydrique et chélates de cobalt. *J Physiol (Paris)* 50(2):438-442.

- Paulet G. 1960a. L'intoxication cyanhydrique et son traitement. Masson, Paris, France.
- Paulet G. 1960b. Les chélates de cobalt dans le traitement de l'intoxication cyanhydrique [Cobalt chelates in the treatment of hydrocyanic intoxication]. *Pathol Biol (Paris)* 8:256-266.
- Paulet G. 1960. Propriétés toxico-pharmacologiques du Kelocyanor. Ref type: Report.
- Paulet G. 1961. Nouvelles perspectives dans le traitement de l'intoxication cyanhydrique. [New perspectives in the treatment of hydrocyanide intoxication.] *Arch Mal Prof* 22:120-127.
- Paulet G. 1965. Au sujet de traitement de l'intoxication cyanhydrique par les chélates de cobalt. *Urgence médicale et chirurgicale* 2:611-613.
- Paulet G. 1984. État actuel du traitement de l'intoxication cyanhydrique. [Present situation of the treatment of cyanide poisoning.] *Arch Mal Prof* 45:185-192.
- Paulet G, Dassonville J. 1985. Valeur du diméthylaminophénol (DMAP) dans le traitement de l'intoxication cyanhydrique: étude expérimentale. *J Toxicol Clin Exper* 5:105-111.
- Paulet G, Desnos J. 1961. L'Acrylonitrile, toxicité, mécanisme d'action thérapeutique. *Arch Int Pharmacodyn* 131 :54-83.
- Paulet G, Le Bars R. 1959. Réanimation après mort par intoxication cyanhydrique (étude chez le chien). *J Physiologie* 51:547-548.
- Paulet G, Olivier M. 1963. Au sujet de la valeur thérapeutique de l'hydroxocobalamin dans l'intoxication cyanhydrique. *Ann Pharm Fr* 21:133-137.
- Paulet G, Bernard JP, Olivier M. 1962. [Hydroxocobalamin in the treatment of hydrocyanic acid poisoning.]. *C R Séances Soc Biol Fil* 156:1867-1869.
- Paulet G, Chary R, Bocquet P. 1959. De l'antidotisme réciproque entre le cobalt et le groupement CN. *C R Soc Biol* 153:1800-1801.
- PCI. 2010. Acrylonitril and Derivatives. World Supply / Demand Report 2010. PCI Acrylonitrile Ltd, Guildford, Surrey GU2 4BX, England.
- Péclet C, Rousseau M, Ponton G, Rousseau JJ. 1982. Carbon monoxide and cyanide inhalation. *Can Soc Forens Sci J* 15:81-86.
- Peddy SB, Rigby MR, Shaffner DH. 2006. Acute cyanide poisoning. *Pediatr Crit Care Med* 7:79-82.

Peden NR, Taha A, McSorley PD, Bryden GT, Murdoch IB, Anderson JM. 1986. Industrial exposure to hydrogen cyanide: Implications for treatment. *Br Med J (Clin Res Ed)* 293:538.

Pedigo LG. 1888. Antagonism between amyl nitrite and prussic acid. *Trans Med Soc Virginia* 19:124-131, 6.14.

Penninks AH, Verschuren PM, Seinen W. 1983. Di-n-butyltin dichloride uncouples oxidative phosphorylation in rat liver mitochondria. *Toxicol Appl Pharmacol* 70:115-120.

Pentore R, Venneri A, Nichelli P. 1996. Accidental choke cherry poisoning: Early symptoms and neurological sequelae of an unusual case of cyanide intoxication. *Ital J Neurol Sci* 17:233-235.

Perrin A, Gout E, Chambaz EM, Defaye G. 1994. Metabolism of malate in bovine adrenocortical mitochondria studied by ¹³C-NMR spectroscopy. *Eur J Biochem* 223:51-59.

Perrone J, Hoffman RS. 1994. Use of sodium nitrite needs further investigation. *Ann Emerg Med* 24:539-540.

Perschau RA, Modell JH, Bright RW, Shirley PD. 1977. Suspected sodium nitroprusside-induced cyanide intoxication. *Anesth Analg* 56:533-537.

Personne M, Kulling P. 1997. Hydroxocobalamin nytt effektivt motgift. *Laekartidningen*, 94:877-878.

Persson HE. 1984. Aspects on antidote therapy in acute poisoning affecting the nervous system. *Acta Neurol Scand Suppl* 100:203-213.

Persson HE, Sjöberg GK, Haines JA, Pronczuk De Garbino J. 1998. Poisoning severity score. Grading of acute poisoning. *J Toxicol Clin Toxicol* 36:205-213.

Persson S-A, Cassel G, Sellström Å. 1986. Acute cyanide intoxication and central transmitter systems. Third symposium on prophylaxis and treatment of chemical poisoning, Stockholm, Sweden, Apr. 22-24, 1985. *Fundam Appl Toxicol* 5:150-159.

Péry-Man N, Houeto P, Coirault C, Suard I, Perennec J, Riou B, Lecarpentier Y. 1996. Hydroxocobalamin vs cobalt toxicity on rat cardiac and diaphragmatic muscles. *Intensive Care Med* 108-115.

Peter H, Bolt HM. 1985. Effect of antidotes of acute toxicity of methacrylonitrile. *Int Arch Occup Environ Health* 55:175-177.

Peter H, Buchter A. 1984. Clinical toxicology of acrylonitrile. *G. Ital. Med. Lav.* 6:83-86.

Peters CG, Mundy JV, Rayner PR. 1982. Acute cyanide poisoning: The treatment of a suicide attempt. *Anaesthesia* 37:582-586.

Pham JC, Huang DT, McGeorge FT, Rivers EP. 2007. Clarification of cyanide's effect on oxygen transport characteristics in a canine model. *Emerg Med J* 24:152-156.

Phillips CI, Wang MK, van Peborgh PF. 1970. Some observations on the mechanism of tobacco amblyopia and its treatment with sodium thiosulphate. *Trans Ophthalmol Soc UK* 90:809-26.

Pickering WG. 1985. Cyanide toxicity and the hazards of dicobalt edetate. *Br Med J (Clin Res Ed)* 291:1644.

Pijoan M. 1942. Cyanide poisoning from choke cherry seed. *Am J Med Sci* 204:550-553.

Pill J, Engeser P, Höbel M, Kreye VA. 1980. Sodium nitroprusside: Comparison of the antidotal effect of hydroxocobalamin and sodium thiosulfate in rabbits. *Dev Toxicol Environ Sci* 8:423-426.

Piper J. 1975. Use and toxicity of nitroprusside. *New Engl J Med* 292:1081-1082. ordered

Pitt BR, Radford EP, Gurtner GH, Traystman RJ. 1979. Interaction of carbon monoxide and cyanide on cerebral circulation and metabolism. *Arch Environ Health*, 34:354-359.

Plackova S. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 1 cyanide case]. Personal communication. National Toxicological Information Center, Bratislava, Slovak Republic.

Polson CJ, Green MA, Lee MR. 1983. *Clinical toxicology*, 3rd ed., London, Pitman 162-196.

Pontal PG, Bismuth C, Garnier R, Pronczuck de Garbino J. 1982. Therapeutic attitude in cyanide poisoning - retrospective study of 24 non-lethal cases. *Vet Hum Toxicol*, 24:90-94.

Poole CJ, Kind PR. 1986. Deficiency of thiosulphate sulphurtransferase (rhodanese) in Leber's hereditary optic neuropathy. *Br Med J (Clin Res Ed)* 292:1229-1230.

Porter DW., Baskin SI. 1996. The effect of three alpha-keto acids on 3-mercaptopyruvate sulfurtransferase activity. *J Biochem Toxicol* 11:45-50.

Posner MA, Tobey RE, McElroy H. 1976a. Hydroxocobalamin therapy of cyanide intoxication in guinea pigs. *Anesthesiology*. 44:157-160.

Posner MA, Rodkey FL, Tobey RE. 1976b. Nitroprusside-induced cyanide poisoning: Antidotal effect of hydroxocobalamin. *Anesthesiology* 44:330-335.

Posner MA, Rodman GH, Klick JM, Tobey RE. 1977. Cyanide production during nitroprusside therapy in an oliguric patient. *Anesth Analg* 56:729-731.

Potter AL. 1950. The successful treatment of two recent cases of cyanide poisoning. *Br J Ind Med* 7:125-130.

Pouwels HMM, Bos LP, Hooymans PM, Parren AGCM. 1987. Een patient met een industriële blauwzuurintoxicatie. *Ned Tijdschr Geneesk* 131:13-16.

Pozzani UC, Carpenter CP, Palm PE, Weil CS, Nair JH. 1959. An investigation of the mammalian toxicity of acetonitrile. *J Occup Med* 1:634-642.

Prajapati NC, Puri RK, Sarangi MP, Yadav S, Khalil A. 1992. Potassium cyanide poisoning. *Indian Pediatr* 29:903-905.

Prieto I, Pujol I, Santiuste C, Poyo-Guerro R, Diego A. 2005. Acute cyanide poisoning by subcutaneous injection. *Emerg Med J* 22:389-390.

Pronczuk de Garbino J, Bismuth C. 1981. Propositions thérapeutiques actuelles en cas d'intoxication par les cyanure. *Toxicol Eur Res* 3 :69-76.

Propriétés pharmacotoxocologiques du KELOCYANOR. Laroche-Navarron, Laboratoires.

Ref Type: Serial (Book, Monograph)

Puntel RL, Nogueira CW, Rocha JB. 2005. Krebs cycle intermediates modulate thiobarbituric acid reactive species (TBARS) production in rat brain in vitro. *Neurochem Res* 30:225-235.

Puskarczyk E. 2006. Aspects toxicocinétiques, toxicodynamiques et thérapeutiques actuels de l'intoxication aiguë à l'ion cyanure. Thesis, Faculté de Médecine. Université Henri Poincaré, Nancy, France.

Queisser H. 1966. Tödliche Blausäurevergiftungen durch Genuss süßer Cassava (*Manihot aipi*): Bericht über 2 Fälle. *Dtsch Gesundheitsw* 21:726-728.

Rachinger J, Fellner FA, Stieglbauer K, Trenkler J. 2002. MR changes after acute cyanide intoxication. *Am J Neuroradiol* 23:1398-1401.

Racle JP, Chausset R, Dissait F, Fontanella JM. 1976. L'intoxication de cyanhydrique aiguë. *Rev Med Clermont-Ferrand* 5:371-382.

Radeleff. 1970. Cyanides and cyanogenic plants. In *Veterinary toxicology*, 2nd ed. Lea and Febiger, Philadelphia, Pennsylvania, USA 50-58 [Review].

Rainey PM, Roberts WL. 1993. Diagnosis and misdiagnosis of poisoning with the cyanide precursor acetonitrile: nail polish remover or nail glue remover? *Am. J. Emerg. Med.* 11:104-108.

Ram Z, Spiegelman R, Findler G, Hadani M. 1989. Delayed postoperative neurological deterioration from prolonged sodium nitroprusside administration, case report. *J Neurosurg*, 71:605-607.

Raub JA, Benignus VA. 2002. Carbon monoxide and the nervous system. *Neurosci Biobehav Rev* 26:925-940.

Rauscher LA, Hurst JM, Collins GM. 1978. Nitroprusside toxicity in a renal transplant patient. *Anesthesiology* 49:428-430.

Rein H, Ristau O, Kupfer E, Jung F. 1968. [Nitrosohemoglobinemia in sodium nitrite poisoning.] *Folia Haematol* 90:176-181 [German].

Reinke LA, Kauffman FC, Belinsky SA, Thurman RG. 1983. Inhibition of p-nitroanisole O-demethylation in perfused rat liver by potassium cyanide. *Arch Biochem Biophys* 225:313-319.

Renard C, Borron SW, Renaudeau C, Baud FJ. 2005. Sodium thiosulfate for acute cyanide poisoning: study in a rat model. *Ann Pharm Fr* 63:154-161.

Renner G. 1974. [Biochemical reactions of cyanic acid as the basis for therapy of cyanic acid poisoning]. *Dtsch Med Wochenschr* 99:1693-1695.

Reyes-Prieto A, El-Hafidi M, Moreno-Sanchez R, Gonzalez-Halphen D. 2002. Characterization of oxidative phosphorylation in the colorless chlorophyte *Polytomella* sp. Its mitochondrial respiratory chain lacks a plant-like alternative oxidase. *Biochim Biophys Acta* 1554:170-179.

Reynolds JEF, ed. 1989. The extra pharmacopoeia, 29th ed. Pharmaceutical Press, London, UK.

Rindone JP, Sloane EP. 1992. Cyanide toxicity from sodium nitroprusside: Risks and management. *Ann Pharmacotherm* 26:515-519.

Riou B, Baud FJ. 1996. L'hydroxocobalamine: Quand une vitamine devient un antidote. *Médecine thérapeutique* 10:791-797.

Riou B, Lecarpentier Y, Toffis V, Barriot B, Baud FJ, Astier A, Viars P. 1989. Effets *in vitro* de l'hydroxocobalamine après intoxication cyanhydrique. *Ann Fr Anesth Reanim* 8:241.

- Riou B, Baud FJ, Astier A, Barriot P, Lecarpentier Y. 1990. *In vitro* demonstration of the antidotal efficacy of hydroxocobalamin in cyanide poisoning. *J Neurosurg Anaesthesiol* 2:296-304.
- Riou B, Gérard JL, Drieu La Rochelle C, Bourdon R, Berdeaux A, Giudicelli JF. 1991. Hemodynamic effects of hydroxocobalamin in conscious dogs. *Anesthesiology* 74:552-558.
- Riou B, Baud FJ, Renaud C, Bismuth C. 1992. L'hydroxocobalamine. In Baud FJ, Barriot P, Riou B, eds, *Les antidotes*. Masson, Paris, France, pp 175-193.
- Riou B, Berdeaux A, Pussard E, Giudicelli JF. 1993. Comparison of the hemodynamic effects of hydroxocobalamin and cobalt edetate at equipotent cyanide antidotal doses in conscious dogs. *Intensive Care Med* 19:26-32.
- Riudavets MA, Aronica-Pollak P, Troncoso JC. 2005. Pseudolaminar necrosis in cyanide intoxication: a neuropathology case report. *Am J Forensic Med Pathol* 26:189-191.
- Rivera-Ch M, León-Velarde F, Huicho L, Monge CC. 1995. Time-course of the polycythemic response in normoxic and hypoxic mice with high blood oxygen affinity induced by cyanate administration. *J Comp Physiol B* 164:659-662.
- Robin ED, McCauley R. 1992. Nitroprusside-related cyanide poisoning, time (long past due) for urgent, effective interventions. *Chest* 102:1842-1845.
- Robinson A. 1959. A case of cyanide poisoning. *JR Nav Med Serv* 45:93-95.
- Roch-Ramel F, Werner D, Guisan B. 1994. Urate transport in brush-border membrane of human kidney. *Am J Physiol* 266:F797-805.
- Rödelsperger R, Kögel T. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 17 cyanide and 2 nitrile cases]. Personal communication. Evonik Röhm, Darmstadt, Germany.
- Rose CL, Worth RM, Kikuchi K, Chen KK. 1965. Cobalt salts in acute cyanide poisoning. *Proc Soc Exp Biol Med* 120:780-783.
- Rosen PJ, Johnson C, McGehee WG, Beutler E. 1971. Failure of methylene blue treatment in toxic methemoglobinemia. Association with glucose-6-phosphate dehydrogenase deficiency. *Ann Intern Med* 75:83-86 [Mechanism of action against methemoglobonaemia and side effects].
- Rosenberg NL, Myers JA, Martin WR. 1989. Cyanide-induced parkinsonism: Clinical, MRI, and 6-fluorodopa PET studies. *Neurology* 39:142.

Rousselin X, Garnier R. 1985. L'intoxication cyanhydrique: Conduite à tenir en milieu de travail et aspect actuel du traitement de l'intoxication aiguë. Fiche médico-technique TC11, Documents pour le Médecin du Travail 23:3-8.

RTI International. 2006. Cyanide: Understanding the risk, enhancing preparedness. Clin Toxicol 44:47-63.

Ruangkanchanasetr S, Wananukul V, Suwanjutha S. 1999. Cyanide poisoning, 2 cases report and treatment review. J Med Assoc Thai 82 Suppl 1:S162-167.

Ruangkanchanasetr S, Wananukul V, Suwanjutha S. 1999. Cyanide poisoning, 2 cases report and treatment review. J Med Assoc Thai 82 Suppl 1:S162-167.

Rubino JM. 1978. Cyanide toxicity: Report of a case. Connecticut Poison Inf Bull 3:1-6.

Rubino MJ, Davidoff F. 1979. Cyanide poisoning from apricot seeds. JAMA 241:359.

Rustin P, Lance C. 1986. Malate Metabolism in Leaf Mitochondria from the Crassulacean Acid Metabolism Plant Kalanchoe blossfeldiana Poelln. Plant Physiol 81:1039-1043.

Sadoff L, Fuchs K, Hollander J. 1978. Rapid death associated with laetrile ingestion. JAMA 239:1532.

Saincher A, Swirsky N, Tenenbein M. 1994. Cyanide overdose: Survival with fatal blood concentration without antidotal therapy. J Emerg Med 12:555-557.

Salkowski AA, Penney DG. 1994. Cyanide poisoning in animals and humans: A review. Vet Human Toxicol 36:455-466.

Salkowski AA, Penney DG. 1995. Metabolic, cardiovascular, and neurologic aspects of acute cyanide poisoning in the rat. Toxicol Lett 75:19-27.

Samo DG. 1988. Cyanide poisoning and the anion gap. Letter to the editor [with reply by Krieg and Saxena]. Ann Emerg Med 17:298-299.

Sartorelli E. 1966. Intossicazione acute da acrilonitrile. Med Lav 57:184-187.

Sarikaya I, Apaydin H, Topal U, Karaoglan O. 2006. Cyanide-induced parkinsonism and F-18 FDG PET/CT findings. Clin Nucl Med 31(6):363-364.

Sass MD, Caruso CJ, Farhangi M. 1967. TPNH-methemoglobin reductase deficiency: A new red-cell enzyme defect. J Lab Clin Med 70:760-767.

Sato M, Ishizu S, Momotani H. 1975. [Determination of acrylonitrile, cyanide and thiocyanate in blood and urine]. *Jap J Ind Health* 17:99-105 [Japanese; English abstract].

Satpute RM, Hariharakrishnan J, Bhattacharya R. 2008. Alpha-ketoglutarate and N-acetyl cysteine protect PC12 cells from cyanide-induced cytotoxicity and altered energy metabolism. *Neurotoxicology* 29:170-178.

Sauer O. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 88 cyanide and 20 nitrile cases]. Personal communication. Giftinformationszentrum der Länder Rheinland-Pfalz und Hessen, Klinische Toxikologie, Universitätsklinikum, Mainz, Germany.

Sauer SW, Keim ME. 2001. Hydroxocobalamin: Improved public health readiness for cyanide disasters. *Ann Emerg Med* 37:635-641.

Saunders JP, Himwich WA. 1950. Properties of the transulfurase responsible for conversion of cyanide to thiocyanate. *Am J Physiol* 163:404-409.

Sax NI. 1984a. Sodium nitrite. In *Dangerous properties of industrial materials*, 6th ed. Van Nostrand Reinhold, New York, USA pp 2442-2443.

Sax NI. 1984b. Sodium sulphate. In *Dangerous properties of industrial materials*, 6th ed. Van Nostrand Reinhold, New York, USA, pp ??

Sayre JW, Kaymakcalan S. 1964. Cyanide poisoning from apricot seeds among children in Central Turkey. *N Engl J Med* 270:1113-1115.

Scalley RD, Ferguson DR, Piccaro JC, Smart ML, Archie TE. 2002 Treatment of ethylene glycol poisoning. *Am Fam Physician* 66:807-812.

Schäfer SG, Maurer HH. 1993. Erkennen und Behandeln von Vergiftungen: ein Leitfaden für Ärzte, Apotheker und Studenten. BI Wissenschaftsverlag, Mannheim, Germany 102-105.

Scheck T. 1992. Intoxikation mit Kaliumcyanid in suizidaler Absicht. *Notfallmedizin* 18:546-547.

Scheffer JG, Campello AF, Voss DO. 1970. Efeitos farmacológicos da hidroxocobalamina. Antagonismo entre a hidroxocobalamina e cianetos. [Pharmacological effects of hydroxocobalamin. Antagonism between hydroxocobalamin and cyanates]. *Hospital (Rio J)* 78:391-397.

Schmidt ES, Newton GW, Sanders SM, Lewis JP, Conn E. 1978. Laetrile toxicity studies in dogs. *JAM* 239:943-947.

Schneider JF, Westley J. 1969. Metabolic interrelations of sulphur in proteins, thiosulfate, and cystine. *J Biol Chem* 244:5735-5744.

Schnelle R. 2007. Spezielle Vergiftungen: Meist ist Sauerstoff richtig und wichtig. *Rettungsdienst* 30:1048-1053.

Schreiner RD, Morgan HJ. 1989. A near-fatal poisoning. *J Tenn Med Assoc* 82:138.

Schubert J, Brill WA. 1968. Antagonism of experimental cyanide toxicity in relation to the *in vivo* activity of the cytochrome oxydase. *J Pharmacol Exp Ther* 162:352-359.

Schulz V. 1984. Clinical pharmacokinetics on nitroprusside, cyanide, thiosulphate and thiocyanate. *Clin Pharmacokinet* 9:239-251.

Schulz V, Roth B. 1982. Detoxification of cyanide in a new-born child. *Klin Wochenschr* 60:527-528.

Schulz V, Döhring W, Rathsack P. 1978. Thiozyanat-Vergiftung bei der antihypertensiven Therapie mit Natriumnitroprussid. *Klin Wochenschr* 56:355-361.

Schulz V, Bonn R, Kindler J. 1979a. Kinetics of elimination of thiocyanate in 7 healthy subjects and in 8 subjects with renal failure. *Klin Wochenschr* 57:243-247.

Schulz V, Bonn R, Kämmerer H, Kriegel R, Ecker N. 1979b. Counteraction of cyanide poisoning by thiosulfate when administering sodium nitroprusside as a hypotensive treatment. *Klin Wochenschr* 57:905-907.

Schulz V, Gross R, Pasch T, Busse J, Loeschcke G. 1982. Cyanide toxicity of sodium nitroprusside in therapeutic use with and without sodium thiosulphate. *Klin Wochenschr* 60:1393-1400.

Schwanecke R. 1966. Unfallgefahren beim Umgang mit Acrylnitril und Metacrylnitril. *Zbl Arbeitsmed* 16:1-3.

Schwartz C, Morgan, RL, Way LM, Way JL. 1979. Antagonism of cyanide intoxication with sodium pyruvate. *Toxicol Appl Pharmacol* 50:437-441.

Schwarzkopf HA, Friedberg KD. 1971. Zur Beurteilung der Blausäure-Antidote. *Arch Toxicol* 27:111-123.

Scolnick B, Hamel D, Woolf AD. 1993. Successful treatment of life-threatening propionitrile exposure with sodium nitrite/sodium thiosulfate followed by hyperbaric oxygen. *J Occup Med* 35:577-580.

Scott EM, Wright RC. 1980. Genetic polymorphism of rhodanese from human erythrocytes. *Am J Hum Genet* 32:112-114.

Seeger R, Neumann H-G. 1988. Cyanwasserstoff (Blausäure), Cyanide. *DAZ-Giftlexikon. Deutsche Apotheker Zeitung* 38:1924-1930.

Selden BS, Clark RF, Curry SC. 1990. Elimination kinetics of cyanide after acute KCN ingestion. *Vet Hum Toxicol* 32:361.

SERB. 1990a. Kelocyanor, ampoules injectables. Document de synthèse.

SERB. 1990b. Kelocyanor. Résumé des caractéristiques du produit.

Shea M, Koziol JA, Howell SB. 1984. Kinetics of sodium thiosulfate, a cisplatin neutralizer. *Clin Pharmacol Ther* 35:419-425.

Sheehy M, Way JL. 1968. Effect of oxygen on cyanide intoxication III: Mithridate. *J Pharmacol Experim Toxicol* 161:163-168. WS thesis

Shimada T, Yamazaki H, Mimura M, Inui Y, Guengerich FP. 1994. Interindividual variances in human liver cytochrome P-450 enzymes involved in the oxidation of drugs, carcinogens and toxic chemicals: studies with liver microsomes of 30 Japanese and 30 Caucasians. *J Pharmacol Exp Ther* 270:414-423.

Shragg TA, Albertson TE, Fisher CJ. 1982. Cyanide poisoning after bitter almond ingestion. *West J Med* 136:65-69.

Shraideh Z, Baranowski Z, Korohoda W, Wohlfarth-Bottermann KE. 1983. Consequences of impeding in mitochondrial function in *Physarum polycephalum*. I: Reversible effects of anoxia, KCN, and influences of the Ca²⁺ ionophore A-23187. *Eur J Cell Biol* 31:175-186.

Shults MC, Updike SJ, Treichel PM, Treichel IW. 1974. High efficiency oxford type oxygenator based on hemodialysis and transmembrane catalysis of hydrogen peroxide. *Trans Am Soc Artif Intern Organs* 286-91.

Shusterman D, Alexeeff G, Hargis C, Kaplan J, Sato R, Gelb A, Becker C, Benowitz N, Gillen M, Thollaug S, Balmes J. 1996. Predictors of carbon monoxide and hydrogen cyanide exposure in smoke inhalation patients. *Clin Toxicol* 34:61-71.

Shusterman D. 1994. Surrogate laboratory measures of cyanide intoxication. *Ann Emerg Med* 24:537-538.

Shusterman DJ. 1993. Clinical smoke inhalation injury: systemic effects. *State Art Rev Occup Med* 8:469-503.

Silber AM, Rojas RL, Urias U, Colli W, Alves M J. 2006. Biochemical characterization of the glutamate transport in *Trypanosoma cruzi*. *Int J Parasitol* 36:157-163.

Silver EH, Kuttub SH, Hasan T, Hassan M. 1982. Structural considerations in the metabolism of nitriles to cyanide in vivo. *Drug Metab. Dispos.* 10:495-498.

Silvestrini MC, Sarti P, Tegoni M. 1995. Reaction of the *Hansenula anomala* flavocytochrome b2 and cytochrome b2 core with inorganic outer sphere redox compounds. *Biochimie* 77:531-8.

Simpson DP. 1988. Dissociation of acid-base effects on substrate accumulation and on delta pH in dog mitochondria. *Am J Physiol* 254:F863-870.

Singh BM, Coles N, Lewis P, Braithwaite RA, Nattrass M, FitzGerald MG. 1989. The metabolic effects of fatal cyanide poisoning. *Postgrad Med J* 65:923-925.

Sipe EK, Trienski TL, Porter JM. 2001. Cyanide toxicity in the surgical intensive care unit: A case report. *Am Surg* 67:684-686.

Skene WG, Norman JN, Smith G. 1966. Effect of hyperbaric oxygen in cyanide poisoning. *Proceedings of the Third International Congress on Hyperbaric Medicine*. In Brown IW Jr, Cox BG, eds, *Hyperbaric Medicine*, pp 705-710.

Slayman CW, Rees DC, Orchard PP, Slayman CL. 1975. Generation of adenosine triphosphate in cytochrome-deficient mutants of *Neurospora*. *J Biol Chem* 250:396-408.

Smet H. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 17 cases]. Personal communication. Poisons Centre, Hospitaalcentrum van de Basis Koningin Astrid, Brussels, Belgium.

Smith AD. 1964. Cyanide encephalopathy in man? *Lancet* 19:668-670. FB 1950-1966 CN cas clinique

Smith FP, Butler TP, Cohan S, Schein PS. 1977. Laetrile toxicity: A report of two cases. *JAMA* 238:1361.

Smith FP, Butler TP, Cohan S, Schein PS. 1978. Laetrile toxicity: A report of two patients. *Cancer Treat Rep* 62:169-171.

Smith PW, Crane CR, Sanders DC, Abbott JK, Endecott B. 1974. Effects of exposure to carbon monoxide and cyanide. *Flammability Research Center, University of Utah, Salt Lake City, UT*,

USA: Conference proceedings of an International Symposium on the Physiological and Toxicological Aspects of Combustion Products. National Academy of Science.

Smith RP, Kruszyna H. 1974. Nitroprusside produces cyanide poisoning via reaction with hemoglobin. *J Pharmacol Exp Ther* 191:557-563.

Smith RP. 1973. Cyanate and thiocyanate: Acute toxicity. *Proc Soc Exp Biol Med* 142:1041-1044.

Smyth HF, Weil CS, West JS, Carpenter CP. 1969. An exploration of joint toxic action: Twenty-seven industrial chemicals intubated in rats in all possible pairs. *Toxicol Appl Pharmacol* 14: 340-347.

SNF (2005). SNF unpublished. Testing laboratory: SNF. In REACH registration, October 2010.

Society of Dyers and Colourists. 1979. The colour index. Society of Dyers and Colourists, Bradford, UK.

Solomon JJ, Cote IL, Wortman M, Decker K, Segal A. 1984. *In vitro* alkylation of calf thymus DNA by acrylonitrile. Isolation of cyanoethyl adducts of guanine and thymine and carboxylethyl adducts of adenine and cytosine. *Chem Bio Interact* 51:167-190.

Sörbo B. 1972. The pharmacology and toxicology of inorganic sulphur compounds. In Senning A, ed. Sulfur in organic and inorganic chemistry, vol 2. Marcel Dekker, New York, USA, pp 156-158.

Spach PI, Cunningham CC. 1987. Control of state 3 respiration in liver mitochondria from rats subjected to chronic ethanol consumption. *Biochim Biophys Acta* 894:460-467.

Speijers GJA, Krajnc EI, Berkvens JM, van Logten MJ. 1982. Acute oral toxicity of inorganic cobalt compounds in rats. *Food Chem Toxicol* 20:311-314.

SreejaVG, .Nagahara Li Q, Minami M. 2003. New aspects on pathogenesis of konzo: neural cell damage directly caused by linamarin contained in cassava (*Manihot esculenta* Crantz). *Br J Nutr* 90:467-472.

Stamyr K, Vaittinen O, Jaakola J, Guss J, Metsälä M, Halonen L, Johanson G. 2009. Background levels of hydrogen cyanide in exhaled breath of healthy population (91). Presented at EAPCCT congress, May 2009, Stockholm, Sweden.

Starkov AA, Fiskum G, Chinopoulos C, Lorenzo BJ, Browne SE, Patel MS, Beal MF. 2004. Mitochondrial alpha-ketoglutarate dehydrogenase complex generates reactive oxygen species. *J Neurosci* 24:7779-7788.

Staub M, Denes G. 1969. Purification and properties of the 3-deoxy-D-arabino-heptulosonate-7-phosphate synthase (phenylalanine sensitive) of *Escherichia coli* K12. II: Inhibition of activity of the enzyme with phenylalanine and functional group-specific reagents. *Biochim Biophys Acta* 178:599-608.

Steffens W, Kehrig B, Lewalter J. 2001. Acrylnitril-Intoxikationen – Cyanidbildung und Therapie. *Arbeitsmed Sozialmed Umweltmed*, 36:351-355.

Steffens W, Sibbing D, Kiesselbach N, Lewalter J. 1998. Behandlung von Patienten mit Vergiftungen durch aliphatische oder olefinische Nitrile. *Arbeitsmed Sozialmed Umweltmed*, 33:479-484.

Steffens W, Leng G. 2003. Nitrile poisonings – cyanide formation, clinical course and treatment. BSD-DOR Medical Services, Dormagen, Germany.

Steffens W. 2002. Cyanide concentrations from nitriles poisoning and smoke inhalations – levels and therapeutical consequences. PhD thesis. Université Catholique de Louvain, Brussels, Belgium.

Steffens W. 2003a. Is N-acetylcysteine an effective cyanide antidote? BSD-LEV Medical Services, Leverkusen, Germany [Abstract].

Steffens W. 2003b. Is N-acetylcysteine an efficient cyanide antidote? Poster presented at EAPCCT. Bayer Industry Services, Monheim, Germany.

Steffens W, Leng G, Pelster M. 2003. Percutaneous hydrocyanic acid poisoning. *Clinical Toxicology*, 41:483-484.

Steffens W. 2008. ECETOC questionnaire on cyanide and nitrile poisoning [response for 7 cases]. Personal communication. Bayer CropScience, Monheim, Germany.

Steffens W. 2010. Percutaneous hydrocyanic acid poisoning. Abstract 129; Presented at the EAPCCT meeting, Rome 2003.

Steinhaus RK, Baskin SI, Clark JH, Kirby SD. 1990. Formation of methemoglobin and metmyoglobin using 8-aminoquinoline derivatives or sodium nitrite and subsequent reaction with cyanide. *J Appl Toxicol* 10:345-351.

Stemler FW, Groff WA Sr, Kaminskis A, Johnson RP, Froehlich HL, Hawkins SF. 1994. A comparison of the treatment of cyanide poisoning in the cynomolgus monkey with sodium nitrite of 4-Dimethylaminophenol (4-DMAP), with and without sodium thiosulfate. US Army Medical Research Institute of Chemical Defense. Aberdeen Proving Ground, MD 210 10-5425

Stewart R. 1974. Cyanide poisoning. *Clin Toxicol* 7:561-564.

Stickel E. 2008. Behandlung der Zyanidvergiftung mittels 4-DMAP. MD thesis. Technical University, Munich, Germany.

Stocker WG, Thiess AM. 1986. Notfallversorgung von Intoxikationen. Verhandlungen der Deutschen Gesellschaft für Arbeitsmedizin, 26. Jahrestagung in Hamburg 07. Gentner Verlag Stuttgart 209-214.

Stotz et al. 1938. Hogness

Stotz E, Sidwell Jr. A. E., and Hogness T. R. 1938. The rôle of the cytochromes in the action of Indophenol Oxidase. J. Biol. Chem., 124:733-744.

Strickland A, Wang RY, Hoffman RS, Goldfrank LR. 1992. Blood cyanide concentrations after smoke-inhalation. N Engl J Med 326:1362.

Suchard JR, Wallace KL, Gerkin RD. 1998. Acute cyanide toxicity caused by apricot kernel ingestion. Ann Emerg Med 32:742-744.

Sugiura M, Kato R. 1977. Reduction of tertiary amine N-oxides by rat liver mitochondria. J Pharmacol Exp Ther 200:25-32.

Sultana S, Singh T, Ahmad FJ, Bhatnagar A, Mittal G. 2011. Development of nano alpha-ketoglutarate nebulization formulation and its pharmacokinetics and safety evaluation in healthy human volunteers for cyanide poisoning. Environ Toxicol Pharmacol 31:436-442

Sun P, Borowitz JL, Kanthasamy AG, Kane MD, Gunasekar PG, Isom GE. 1995. Antagonism of cyanide toxicity by isosorbide dinitrate: Possible role of nitric-oxide. Toxicology, 104:105-111.

Sunderman FW, Kincaid JF. 1953. Toxicity studies of acetone cyanohydrin and ethylene cyanohydrin. AMA Arch Ind Hyg Occup Med 4:371-376.

Sunshine I, Finkle B. 1964. The necessity for tissue studies in fatal cyanide poisoning. Int Archiv Gewerbepath Gewerbehyg, 20:558-561.

Surgenor DM. 1970. Sodium nitrite USP. In: Osol E, ed. Remington's pharmaceutical sciences, 14th ed. Mack Publishing, Easton, Pennsylvania, USA 846-847.

Sutter ME, Tereshchenko N, Daubert GP. 2009. Hydroxycobalamin: An effective but challenging new antidote. Clin Toxicol 47:702-765.

Sutter M, Tereshchenko N, Rafii R, Daubert GP. 2010. Hemodialysis complications of hydroxocobalamin: A case report. J Med Toxicol 6:165-167.

Swanson JR, Krasselt WG. 1994. An acetonitrile-related death. J. Forensic Sci. 39:271-279.

Sylvester DM, Hayton WL, Morgan RL, Way JL. 1983. Effects of thiosulfate on cyanide pharmacokinetics in dogs. *Toxicol Appl Pharmacol* 69:265-271.

Sylvester DM, Morgan RL, Hayton WL, Way JL. 1980. Effects of sodium thiosulfate on the pharmacokinetics of sodium cyanide. *Fed Proc* 39:1358.

Sylvester DM, Sander C, Hayton WL, Way JL. 1981. Alteration of the pharmacokinetics of sodium cyanide by sodium thiosulfate. *Proc West Pharmacol Soc* 24:135. .

Symington IS, Anderson RA, Thomson I, Oliver JS, Harland WA, Kerr JW. 1978. Cyanide exposure in fires. *Lancet* 2:91-92.

Szczepkowski TW, Skarzynski B, Weber M. 1961. The metabolic state of thiosulfate. *Nature* 189:1007-1008.

Szinicz L, Weger N, Schneiderhan W, Kiese M. 1979. Nephrotoxicity of aminophenols: Effects of 4 dimethylaminophenol on isolated rat kidney tubules. *Arch Toxicol* 42:63-73 [Experimental study].

Taboulet P, Clemessy JL, Freminet A, Baud FJ. 1995. A case of life-threatening lactic-acidosis after smoke-inhalation – interference between beta-adrenergic agents and ethanol. *Intens Care Med* 21:1039-1042.

Tadic V. 1992. The *in vivo* effects of cyanide and its antidotes on rat brain cytochrome oxidase activity. *Toxicology* 76:59-67.

Takahara K, Nakanishi S, Natelson S. 1971. Studies on the reductive cleavage of canavanine and canavaninosuccinic acid. *Arch Biochem Biophys* 145:85-95.

Takano T, Miyazaki Y. 1981. Combined effect of carbon monoxide and potassium cyanide on intracellular oxidation-reduction in rabbit kidney in situ. *Jpn J Hyg* 5:811-815.

Takano T, Miyazaki Y, Nashimoto I, Kobayashi K. 1980. Effect of hyperbaric oxygen on cyanide intoxication: *In situ* changes in intracellular oxidation reduction. *Undersea Biomed Res* 7:191-197.

Tanii H, Hashimoto K. 1984a. Studies on the mechanism of acute toxicity of nitriles in mice. *Arch. Toxicol* 55: 47-54.

Tanii H, Hashimoto K. 1986. Influence of ethanol on the *in vivo* and *in vitro* metabolism of nitriles in mice. *Arch. Toxicol* 58:171-176.

Tanii H, Hashimoto K. 1984b. Structure-toxicity relationship of aliphatic nitriles. *Toxicol. Lett* 22:267-272.

Tarkowski S. 1966. Effect of cobalt on the inhibition of the cytochrome oxidase activity by potassium cyanide. *Med Pr* 17:116-119.

Tartière S, Richter F, Taboulet P, Baud FJ, Laforge M. 1992. Traitement préhospitalier d'une intoxication cyanhydrique sévère par l'hydroxocobalamine. *JEUR* 5:35-37.

Tassan H, Joyon D, Richard T, Lamaison D, Guelon D, Barakeh S. 1990. Intoxication au cyanure de potassium traitée par l'hydroxocobalamine. *Ann Fr Anesth Réanim* 9:383-385.

Ten Eyck RP, Schaerdel AD, Lynett JE, Marks DH, Patrissi GA, Ottinger WE, Stansell MJ. 1984. Stroma-free methemoglobin solution as an antidote for cyanide poisoning: a preliminary study. *Clin Toxicol* 21:343-358.

Ten Eyck RP, Schaerdel AD, Ottinger WE. 1985. Stroma-free methemoglobin solution: an effective antidote for acute cyanide poisoning. *Am J Emergency Med* 3:519-523.

Ten Eyck RP, Schaerdel AD, Ottinger WE. 1985-1986. Comparison of nitrite treatment and stroma-free methemoglobin solution as antidotes for cyanide poisoning in a rat model. *Clin Toxicol* 23:477-487.

TerraBase. 2012. TerraQSAR-LOGPTM computed octanol / water partition coefficients (CLOGPs): Vasodilators [selection]. TerraBase, Hamilton, ON, Canada. www.terrabase-inc.com/

Terzic M, Milosevic M. 1963. Action protectrice de l'éthylène-diaminetétra-acétate-dicobaltique dans l'intoxication cyanée. *Thérapie* 18:55-61 [French].

Their R, Lewalter J, Bolt HM. 2000. Species differences in acrylonitrile metabolism and toxicity between experimental animals and humans based on observation in human accidental poisonings. *Arch Toxicol* 74:184-189.

Thier R, Bolt HM, Lewalter J. 2000. Species differences in acrylonitrile metabolism and toxicity between experimental animals and humans based on observations in human accidental poisonings. *Arch Toxicol* 74:184-189.

Thiess AM, Hey W. 1969. Zur Toxizität des Isobutyronitrils und des α -Hydroxyisobutyronitrils (Acetoncyanhydrin). *Arch Toxicol* 24:271-282.

Thoma R, Zilker T. 1987. Toxische Methämoglobinämie: Toluidinblau als Antidot bei irrtümlicher, hochdosierter Gabe von Dimethyl-p-Aminophenol (4-DMAP). *Notarzt* 3:169-170.

Thomas TA, Brooks JW. 1970. Accidental cyanide poisoning. *Anaesthesia* 25:110-114.

Thomson I, Anderson RA, Harland WA. 1978. The determination of blood cyanide and plasma thiocyanate by gas chromatography — mass spectrometry. In Frigerio A, ed, *Recent developments in mass spectrometry in biochemistry and medicine*. Plenum Press, New York, NY, USA, pp 215-226.

Tinker JH, Michenfelder JD. 1976. Sodium nitroprusside: Pharmacology, toxicology, therapeutics. *Anesthesiology* 45:340-354.

Tomoda I. 1955. Influence of vitamin B12a on the activity of perfused toad heart poisoned by cyanide. *Kumamoto Med J* 8:23-25.

Torremadé Barreda J, Serralladh Orejas M, Valles-Ortega J, Franco Miranda E. 2010. Treatment with hydroxocobalamin for cyanide poisoning: A rare cause of pseudohematuria. *Actas Urol Esp* 34:124-126.

Toth PP, Sumerix KJ, Ferguson-Miller S, Suelter CH. 1990. Respiratory control and ADP:O coupling ratios of isolated chick heart mitochondria. *Arch Biochem Biophys* 276:199-211.

Townsend WA, Boni B. 1975. Cyanide poisoning from ingestion of apricot kernels. *Morbid Mortal Weekly Rep* 24:427.

Tracqui A, Raul JS, Geraut A, Berthelon L, Ludes B. 2002. Determination of blood cyanide by HPLC-MS. *J Anal Toxicol* 26:144-148.

Tradić V. 1992. The in vivo effects of cyanide and its antidotes on rat brain cytochrome oxidase activity. *Toxicology* 76:59-67.

Trapp WG, Lepawsky M. 1983. One hundred percent survival in five life-threatening acute cyanide poisoning victims treated by a therapeutic spectrum including hyperbaric oxygen. *Proceedings of the 1st European Conference on Hyperbaric Medicine Amsterdam, The Netherlands* p 45. Presented at 8th Annual Conference on Clinical Applications of Hyperbaric Oxygen. Long Beach Memorial Hospital Medical Center, Long Beach, California, USA. *IPCS* 2.14.

Trapp WG. 1970. Massive cyanide poisoning with recovery: A boxing-day story. *Can Med Assoc J* 102:517.

Trolli G. 1938. Paraplégie spastique épidémique, “Konzo” des indigènes du Kwango. In: G. Trolli (Ed.): *Résumé des observations réunies, au Kwango, au sujet de deux affections d'origine indéterminée*, Fonds Reine Elisabeth, Brussels 1–36.

Troup CM, Ballantyne B. 1987. Analysis of cyanide in biological fluids and tissues. Chapter 2 in Ballantyne B, Marrs TC, eds, Clinical and experimental toxicology of cyanides. John Wright, Bristol, UK 22-40.

Tsuge K, Kataoka M, Seto Y. 2000. Cyanide and thiocyanate levels in blood and saliva of healthy adult volunteers. J Health Sci 46:343–350.

Tsutaoka BT, Anderson IB, Olson KR. 2003. A case series of dermal acetonitrile exposures. Clin Toxicol 5:643 [Abstract].

Tudella VG, Curti C, Soriani FM, Santos AC, Uyemura SA. 2004. *In situ* evidence of an alternative oxidase and an uncoupling protein in the respiratory chain of *Aspergillus fumigatus*. Int J Biochem Cell Biol 36:162-172.

Tulsawani R, Bhattacharya R. 2006. Effect of α -ketoglutarate on cyanide-induced biochemical alterations in rat brain and liver. Biomed Environ Sci 19:61-66.

Tulsawani RK, Debnath M, Pant SC, Kumar D, Prakash AO, Vijayaraghavan R, Bhattacharya R. 2005. Effect of sub-acute oral cyanide administration in rats: Protective efficacy of alpha-ketoglutarate and sodium thiosulfate. Chem Biol Interact 156:1-12.

Tulsawani R, Kumar D, Bhattacharya R. 2007. Effect of pre-treatment with α -ketoglutarate on cyanide-induced toxicity and alterations in various physiological variables in rodents. Biomed Environ Sci 20:56-63.

Turchen SG, Manoguena AS. 1989. Severe cyanide poisoning following suicidal ingestion of acetonitrile. Vet Hum Toxicol 31:356.

Turchen SG, Manoguerra AS, Whitney C. 1991. Severe cyanide poisoning from the ingestion of acetonitrile-containing cosmetic. Am J Emerg Med 9(3):264-267.

Tylleskär T, Légué FD, Peterson S, Kpizingui E, Stecker P. 1994. Konzo in the Central African Republic. Neurology 44:959-961.

Tyrer FH. 1981. Treatment of cyanide poisoning. J Soc Occup Med 31:65-66.

Udvardy J, Marre E, Farkas GL. 1970. Purification and properties of a phosphodiesterase from Avena leaf tissues. Biochim Biophys Acta 206:392-403.

Uhl W, Nolting A, Golor G, Rost KL, Kovar A. 2006. Safety of hydroxocobalamin in healthy volunteers in a randomized, placebo-controlled study. Clin Toxicol 44:17-28.

Uitti RJ, Rajput AH, Ashenhurst EM, Rozdilsky B. 1985. Cyanide-induced parkinsonism: a clinicopathologic report. *Neurology* 35:921-925.

US EPA. 2004. Estimation Program Interface (EPI) Suite. Ver.3.12. Available from <http://www.epa.gov/oppt/exposure/docs/episuitedi.htm>

US NAC. 2007. Acute exposure guideline levels (AEGLs) for acrylonitrile (CAS Reg. No. 107-13-1), draft July 2007. Unpublished report. National Advisory Committee to develop AEGLs. US Environmental Protection Agency, Washington DC, USA [Review].

US Pharmacopeia. 1985. Amyl nitrite, hydroxocobalamin USP 31 NF 26. (United States Pharmacopoeia: National Formulary). Rockville, Maryland, USA 64-65, 512-513.

US Pharmacopeia. 2008. Oxygen, sodium nitrite, sodium thiosulfate, amyl nitrite, hydroxocobalamin, USP 31 NF 26. (United States Pharmacopoeia: National Formulary). Rockville, Maryland, USA 2886, 3254, 3260, 1423, 2356-2357.

Uyemura SA, Luo S, Moreno SN, Docampo R. 2000. Oxidative phosphorylation, Ca(2+) transport, and fatty acid-induced uncoupling in malaria parasites mitochondria. *J Biol Chem* 275:9709-9715.

van Buuren KJH, Nicholls P, van Gelder BF. 1972. Biochemical and biophysical studies on cytochrome aa3. VI. Reaction of cyanide with oxidized and reduced enzyme. *Biochim Biophys Acta* 256:258-276.

Van der Lande VM. 1972. Observations on the histochemical "aminopeptidase" reaction in the intestine of certain species of leech (Annelida: Hirudinea), with particular reference to *Erpobdella octoculata* (L.). *Comp Biochem Physiol A* 41:813-24.

van Dijk A, Glerum JH, van Heijst CNP, Douze JMC. 1987. Clinical evaluation of the cyanide antagonist 4 DMAP in a lethal cyanide poisoning case. *Vet Hum Toxicol* 29:38-39.

van Dop C, Hutson SM., Lardy HA. 1977. Pyruvate metabolism in bovine epididymal spermatozoa. *J Biol Chem* 252:1303-1308.

van Heijst AN, Douze JM, van Kesteren RG, van Bergen JE, van Dijk A. 1987. Therapeutic problems in cyanide poisoning. *Clin Toxicol* 25:383-398.

van Heijst ANP, Ruse M. 1997. Cyanides poisons information monograph. International Programme on Chemical Safety, Geneva, Switzerland.
www.inchem.org/documents/pims/chemical/pimg003.htm

van Heijst ANP. 1986. N-Acetylcysteine bij vergiftigingen met paracetamol en acrylnitril. Pharm Weekbl 121:663-668.

van Luijt DE. 1963. Casuistische mededelingen: Acrylonitrilvergiftiging met dodelijke afloop. Ned Tijdschr Geneesk 107:2186-2188.

Vernon PA, Dulak LH & Deskin R (1990a). Acute toxicological evaluation of acrylonitrile (1969). Journal of American College of Toxicology (1990):1, 114-115.

Veronneau S, Ribe JK, Sathyavagiswaran L, Lewis I, Muto J. 1992. Lessons learned from the 1991 US Air/SkyWest collision at LAX. 44th Annual Meeting of the American Academy of Forensic Sciences 138.

Vesey CJ. 1979. Blood cyanide. Letter to the editor. Clin Toxicol 14:307-309.

Vesey CJ, Cole PV. 1981. Cyanide antagonist. Can Anaesth Soc J 28:290 [Comment on Kaplan et al, 1980; author reply by Kaplan, 1981].

Vesey CJ, Langford RM. 1998. Stabilization of blood cyanide. J Anal Toxicol 22:176-178.

Vesey CJ, Cole PV, Linnell JC, Wilson J. 1974. Some metabolic effects of sodium nitroprusside in man. Br Med J 2:140-142.

Vesey CJ, Cole PV, Simpson PJ. 1976. Cyanide and thiocyanate concentrations following sodium nitroprusside in man. Br J Anaesth 48:651-660.

Vesey CJ, Krapez JR, Varley JG, Cole PV. 1985. The antidotal action of thiosulfate following acute nitroprusside infusion in dogs. Anesthesiology 62:415-421.

Vest P, Renaudeau C, Tellal S, Ragot C, Renard C. 2002. Interférence de l'hydroxocobalamine sur les résultats des dosages biochimiques usuels. Ann Biol Clin 60 :57-64.

Viana C, Cagnoli H, Cedan J. 1934. Action du nitrite de sodium dans l'intoxication au cyanure. C R Séances Soc Biol Fil 115:1649-1651.

Vick JA, Froehlich HL. 1991. Treatment of cyanide poisoning. Military Medicine 156:330-339.

Vick JA, Von Bredow JD. 1996. Effectiveness of intramuscularly administered cyanide antidotes on methemoglobin formation and survival. J Appl Toxicol, 16:509-516.

Vick JA. 1991. Treatment of Cyanide Poisoning. Military Medicine Vol. 156.

Vick JA., Froehlich HL. 1985. Studies of cyanide poisoning. *Arch Int Pharmacodyn Ther* 273:314-322.

Vincent M, Vincent F, Marka C. 1981. Cyanide and its relationship to nervous suffering. Physiopathological aspects of intoxication. *Clin Toxicol* 18:1519-1527.

Vinogradov AD. 1968. [The effect of alpha-ketoglutarate and ammonia ions on the oxidation of succinate by liver mitochondria]. *Biokhimiia* 33:553-560.

Vock R, Magerl H, Lange O, Betz P, Eisenmenger W, Freisleder A, Graw M, von Meyer L, Mohsenian F, Müller RK, Püschel K, Schmidt V, Schmoldt A. 1999. Handlungsfähigkeit bei tödlichen oralen Intoxikationen mit Cyan-Verbindungen. *Rechtsmedizin* 9:56-61.

Voegtlin C, Johnson JM, Dyer HA. 1926. Biological significance of cystine and glutathione. I: On the mechanism of the cyanide action. *J Pharma Exp Therapy*, 27:467-483.

Vogel RA, Kirkendall WM. 1984. Acrylonitrile (vinyl cyanide) poisoning: A case report. *Texas Med* 80:48-51.

Vogel SN, Sultan TR, Ten Eyck RP. 1981. Cyanide poisoning. *Clin Toxicol* 18:367-383.

Vogel SN. 1987. Lactic acidosis in acute cyanide poisoning. In Ballantyne B, Marrs T, eds. *Clinical and experimental toxicology of cyanides*. John Wright, Bristol, England, UK 451-466.

Von Clarmann M. 1987. 4-Dimethylaminophenol. Communication at the joint meeting of the EAPCC, CEC, IPCS and the National Poison Control Centre in the Netherlands, Utrecht. *IPCS* 8.14

Von Lüpke H, Schmidt K. 1977. Kombinierte Kohlenmonoxid/Cyanwasser-stoff-Vergiftung nach Verschmoren von Kunststoff. *Z Rechtsmed* 79:69-71.

Von Meyer L, Drasch G, Kauert G. 1979. Zur Bedeutung der Bildung von Cyanwasserstoff bei Bränden. *Z Rechtsmed* 84:69-73.

Wacker WE, Haynes H, Druyan R, Fisher W, Coleman J. 1965. Treatment of ethylene glycol poisoning with ethyl alcohol. *J. Am. Med. Assoc.* 194:1231-1233.

Wainio WW, Cooperstein SJ. 1956. Some controversial aspects of the mammalian cytochromes. *Adv Enzymol Relat Subj Biochem* 17:329-392.

Walsh DW, Eckstein M. 2004. Hydrogen cyanide in fire smoke: an underappreciated threat. *Emerg Med Serv* 33:160-163.

- Wananukul W, Kaojarern S. 1992. Acute cyanide poisoning: a case report with toxicokinetic study. *J Med Assoc Thai* 75:304-309.
- Wang H, Chanas, B and Ghanayem BI. 2002. CYP2E1-mediated Oxidation. *Drug Metabolism and Disposition*. Vol. 30, No. 8, 911-917.
- Warburg O. 1927. Über die Wirkung von Kohlenoxyd und Stickoxyd auf Atmung und Gärung. *Biochem. Z.*, 189, 354.
- Watanabe F. 2007. Vitamin B12 sources and bioavailability. *Exp Biol Med (Maywood)* 232:1266-1274.
- Way JL. 1983. Cyanide antagonism. *Fundam Appl Toxicol* 3:383-386.
- Way JL. 1984. Cyanide intoxication and its mechanism of antagonism. *Ann Rev Pharmacol Toxicol* 24:451-481.
- Way JL, Gibbon SL, Sheehy M. 1966. Effect of oxygen in cyanide intoxication. I: Prophylactic protection. *J Pharmacol Exper Ther* 153:381-385.
- Way JL, End E, Sheehy MH, de Miranda P, Feitknecht UF, Bachand R, Gibbon SL, Burrows GE. 1972. Effect of oxygen on cyanide intoxication. IV. Hyperbaric oxygen. *Toxicol Appl Pharmacol* 22:415-421.
- Way JL, Leung P, Cannon E, Morgan R, Tamulinas C, Leong-Way J, Baxter L, Nagi A, Chui C. 1988. The mechanism of cyanide intoxication and its antagonism. *Ciba Found Symp* 140:232-243.
- Weger N. 1968. Aminophenole als Blausäureantidote. *Arch. Toxikol* 24:49-50
- Weger N. 1975. Cyanidvergiftung und Therapie. *Wehrmed. Monatsschrift* 19:6-11.
- Weger NP. 1983. Treatment of cyanide poisoning with 4-dimethylaminophenol (DMAP) – experimental and clinical overview. *Fundam Appl Toxicol*, 3:387-396. s
- Weger NP. 1991. Human intoxications with hydrogen cyanide, hydrogen sulfide, and nitriles treated with DMAP. *Vet Hum Toxicol* 33:366 [Abstract].
- Weng TI, Fang CC, Kin SM, Chen WJ. 2004. Elevated plasma cyanide level after hydroxocobalamin infusion for cyanide poisoning. *Am J Emerg Med* 22:492-493.
- Werner H. 1979. Die Behandlung von Vergiftungen mit Blausäure, ihrer Salze und Derivate mit 4 dimethylaminophenol. PhD thesis. Technical University, Munich, Germany.

Wesson DE, Foley R, Sabatini S, Wharton J, Kapusnik J, Kurtzmann NA. 1985. Treatment of acute cyanide intoxication with hemodialysis. *Am J. Nephrol.* 5:121-126.

Westley J, Adler H, Westley L, Nishida C. 1983. The sulfurtransferases. *Fundam Appl Toxicol* 3:377-382.

Wheatley C. 2007. The return of the Scarlet Pimpernel: Cobalamin in inflammation II – Cobalamins can both selectively promote all three nitric oxide synthases (NOS), particularly iNOS and eNOS, and, as needed, selectively inhibit iNOS and nNOS. *Journal of Nutritional and Environmental Medicine* 16:181-211.

Wherland S, Gray HB. 1976. Metalloprotein electron transfer reactions: analysis of reactivity of horse heart cytochrome c with inorganic complexes. *Proc Natl Acad Sci U S A* 73:2950-4.

White WLB, Arias-Garzon DI, McMahon JM, Sayre RT. 1998. Cyanogenesis in cassava: The role of hydronitrile lyase in root cyanide production. *Plant Physiol*, 116:1219-1225.

Whitworth DG. 2007. Personal Communication.

WHO. 1992. Evaluation of certain food additives and naturally occurring toxicants. Technical Report series 828. Thirty-ninth report of the Joint FAO/WHO Expert Committee on Food Additives. World Health Organization, Geneva, Switzerland 30-31.

WHO. 1993. Cyanogenic glycosides. Toxicological evaluation of certain food additives and naturally occurring toxicants. WHO Food Additives series 30. First draft prepared by Speijers G, National Institute of Public Health and Environmental Protection, Bilthoven, Netherlands. International Programme on Chemical Safety, World Health Organization, Geneva, Switzerland.

Whyte I. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 3 cyanide cases]. Personal communication. Hunter Area Toxicology Service, Newcastle, UK.

Wiedemann K. 1976. Toxizität von Natriumnitroprussid. *Prakt Anästh*, 11:387-396.

Wikimedia Commons. 2006. <http://commons.wikimedia.org/wiki/File:Natriumnitrit.png>.

Wikimedia Commons. 2010.

http://commons.wikimedia.org/wiki/File:Sodium_thiosulfate.svg?uselang=en-gb

Willhite CC, Smith RP. 1981. The role of cyanide liberation in the acute toxicity of aliphatic nitriles. *Toxicol Appl Pharmacol* 59:589-602.

Willhite CC. 1980. Studies on the basis of the toxicity of aliphatic organonitriles. *Diss Abstr Int*, 41:536-B.

Willhite CC. 1981. Inhalation toxicology of acute exposure to aliphatic nitriles. *Clin. Toxicol* 18:991-1003.

Willhite CC. 1983. Developmental toxicology of acetonitrile in the Syrian golden hamster. *Teratology* 27:313-325.

Williams HL, Johnson DJ, McNeil JS, Wright DG. 1990. Studies of cobalamin as a vehicle for the renal excretion of cyanide anion. *J Lab Clin Med* 116:37-44. please

Wilson J. 1965. Leber's hereditary optic atrophy: A possible defect of cyanide metabolism. *Clin Sci* 29:505-515.

Wilson RH, McCormick WE. 1949. Acrylonitrile: Its physiology and toxicology. *Ind Med Surg* 18:243-245.

Wilson RH. 1944. Health hazards encountered in the manufacture of synthetic rubber. *J Am Med Assoc* 124:701-703.

Windholz M, Budavari S, Blumetti RF, ed. 1983. Sodium nitrite. In: *The Merck index: An encyclopedia of chemicals, drugs, and biologicals*, 10th ed. Merck, Rahway, New Jersey, USA, p 8480.

Windholz M, Budavari S, Blumetti RF, ed. 1983. Sodium thiosulphate. In *The Merck index: An encyclopedia of chemicals, drugs, and biologicals*, 10th ed. Merck, Rahway, New Jersey, USA, pp ??.

Wing KS, Cheung CY, Lung LF. 2008. Hydroxocobalamin for acute cyanide poisoning in smoke inhalation [Correspondence on Borron et al, 2007b]. *Ann Emerg Med* 51:108.

Winkelmann J. 1984. Bereitstellung von 4-DMAP als Antidot in H₂S-gefährdeten Betrieben. 21. Ständige Konferenz der Werksärzte der Chemischen Industrie, Mannheim 8:140-143.

Winston JM, Roberts RJ. 1975 Influence of carbon monoxide, hypoxic hypoxia or potassium cyanide pretreatment on acute carbon monoxide and hypoxic hypoxia lethality. *J Pharmacol Exp Ther* 193:713-719.

Winter CA, Mushett CW. 1950. Absence of toxic effects from single injections of crystalline vitamin B₁₂. *J Am Pharm Assoc Am Pharm Assoc* 39:360-361.

- Wirth W. 1935. Experimentelle Studien zur Behandlung der Vergiftung durch Blausäure-Einatmung. *N Schmied Arch Pharmacol* 179:558-602.
- Wiskich JT, Day DA. 1982. Malate Oxidation, Rotenone-Resistance, and Alternative Path Activity in Plant Mitochondria. *Plant Physiol* 70:959-964.
- Wolfsie JH. 1951. Treatment of cyanide poisoning in industry. *AMA Arch Ind Hyg Occup Med* 4:417-425.
- Wood GC. 1982. Acute cyanide intoxication: Diagnosis and management. *Clin Toxicol Consultant* 4:140-149.
- Wood JL, Cooley SL. 1956. Detoxication of cyanide by cystine. *J Biol Chem* 218:449-457.
- Woutersen RA. 1998. Toxicologic profile of acrylonitrile. *Scand J Work Environ Health* 24:5-9.
- Wright IH, Vesey CJ. 1986. Acute poisoning with gold cyanide. *Anaesthesia* 41:936-939.
- Wu ML, Tsai WJ, Ger J, Deng JF, Tsay SH, Yang MH. 2001. Cholestatic hepatitis caused by acute gold potassium cyanide poisoning. *J Toxicol Clin Toxicol* 39:739-743.
- Würzburg H. 1996. Treatment of cyanide poisoning in an industrial setting. *Vet Hum Toxicol* 38:44-47. www.emea.europa.eu/humandocs/PDF's/EPAR/Cyanokit (visited 14/07/2008)
- Borron M, Faure J, Morena H, Vincent M, Faure H. 1974. L'intoxication cyanhydrique aiguë: Données actuelles sur le métabolisme du cyanure et le traitement par hydroxocobalamine. *J Eur Toxicol* 7:22-29.
- Yamamoto HA. 1989. Hyperammonemia, increased brain neutral and aromatic amino acid levels, and encephalopathy induced by cyanide in man. *Toxicol Appl Pharmacol* 99:415.
- Yamamoto HA. 1990. Protection against cyanide-induced convulsions with α -ketoglutarate. *Toxicology* 61:221-228.
- Yamamoto HA. 1992. Nitroprusside intoxication: Protection of α -ketoglutarate and thiosulphate. *Food Chem Toxicol* 30:887-890.
- Yamamoto K. 1976. Acute combined effects of HCN and CO, with the use of the combustion products from PAN (polyacrylonitrile) - gauze mixtures. *Z Rechtsmed* 78:303-311.

Yamamoto K. 1977. Acute combined effects of HCN and CO, with special reference to a theoretical consideration of acute combined effects on the basis of the blood cyanide and COHb analyses. *J Combust Toxicol* 4:69-78.

Yamamoto K, Kuwahara C. 1981. A study on the combined action of CO and HCN in terms of concentration-time products. *Z Rechtsmed* 86:287-294.

Yatziv A, Simcha S. 1969. [Cyanide poisoning during childhood due to consumption of apricot pits]. *Harefuah* 76:535-536 [Hebrew; English translation].

Yen D, Tsai J, Wang LM, Kao WF, Hu SC, Lee CH, Deng JF. 1995. The clinical experience of acute cyanide poisoning. *Am J Emerg Med* 13:524-528.

Youngman RJ, Eyer P, Elstner EF. 1980. Reactions of 4-dimethylaminophenol with illuminated chloroplast lamellae. *Biochem Physiol Pflanzen* 175:789-796.

Yunker, M. H. and Higuchi, T. (1958), Kinetics of degradation of amyl nitrite ampuls II. Stabilization. *J. Pharm. Sci.*, 47: 621–625.

Zaknun JJ, Stieglbauer K, Trenkler J, Aichner F. 2005. Cyanide-induced akinetic rigid syndrome: Clinical, MRI, FDG-PET, beta-CIT and HMPAO SPECT findings. *Parkinsonism Relat Disord* 11:125-129.

Zavotsky KE, Mentler P, Gronczewski C, Torres P. 2004. A 52-year-old acyanotic man with severe hypoxia and profound metabolic acidosis following an industry chemical exposure. *J Emerg Nurs* 30:113-116.

Zerbe NF, Wagner BKJ. 1993. Use of vitamin B12 in the treatment and prevention of nitroprusside-induced cyanide toxicity. *Critical Care Med* 21:465-467.

Zhou L, Jin S, Chen C, Chen Y. 1999. The diagnosis and treatment of acute acrylonitrile poisoning: a clinical study of 144 cases. *J Occup Health* 41:172-176.

Zilker T. 1999. Personal communication, Cyanide Congress (Orphan Europe), Fulda.

Zilker T. 2007. ECETOC questionnaire on cyanide and nitrile poisoning [response for 8 cases]. Personal communication. Center Klinikum rechts der Isar, Toxikologische Abteilung. Technical University, Munich, Germany.

Zilker T. 2010. ECETOC questionnaire on cyanide and nitrile poisoning [response for 1 cyanide case]. Personal communication. Klinikum rechts der Isar, Toxikologische Abteilung. Technical University, Munich, Germany.

Zilker T and Eyer P. 2005. 4-Dimethylaminphenol (4-DMAP) as antidote for poisoning by cyanide. In Brent J, Wallace KL, Burkhart KK, Phillips SD, Donovan JW, eds, Critical care toxicology; Diagnosis and management of the critically poisoned patient. Elsevier Mosby, Saunders, Philadelphia, Pennsylvania, USA, pp 1609-1615.

Zilker T and Felgenhauer N. 2000. 4-DMAP as cyanide antidote: Its efficacy and side effects in human poisoning. Clin Toxicol 38:217.

Zilker T and Schweizer W. 1987. Zyankali-Intoxikation. Der Notarzt 3:59-60.

Zilker T and Solingen, 2010. Personnal communication.

Zoltani CK, Platoff GE, Baskin SI. 2009. Patent title: Methods and compositions for treatment of poison-caused pathology. US Army Research Laboratory; ATTN AMSRL CS CC IP. Adelphi, MD US IPC8 Class: AA61K31185FI USPC Class: 424699.

Zündorf I, Dingermann T. 2008. Neue Cyanid-Antidots. Pharm Unserer Zeit 2:116.

*****References not quoted***

The following references were consulted by the Task Force, but not quoted for various reasons given [in square brackets]. (The list includes references cited by others as indicated, except for some also used directly elsewhere in the report).

*Acrylonitrile Group. 2005a. Chemical emergency medical guidelines acrylonitrile (CH₂=CH CN) A. Information and recommendations for first responders. Draft final. Acrylonitrile Group, Washington DC, USA [Review].

*Acrylonitrile Group. 2005b. Chemical emergency medical guidelines acrylonitrile (CH₂=CH CN) B. Information and recommendations for medical treatment at the site. Draft final. Acrylonitrile Group, Washington DC, USA [Review].

*Acrylonitrile Group. 2005c. Chemical emergency medical guidelines acrylonitrile (CH₂=CH CN) C. Information and recommendations for hospitals / emergency departments. Draft final. Acrylonitrile Group, Washington DC, USA [Review].

*Agarwal PK, Kumari R. 2003. Nitroprusside in critically ill patients with aortic stenosis, author reply. *Engl J Med* 349:811-813 [Comment on Khot et al, 2003].

*AHFS. 1988. American Hospital Formulary Service. In McEnvoy KG, Litvak K, Welsh OH, Campbell JF, Ziegler K, Douglas PM, Shannon-Lass EP, Thomson K, McQuerrie GM, Schmadel LK, Fredrickson MK, Mendham NA, Morisseau AL, eds, Drug information. American Society of Hospital Pharmacists, Bethesda, Maryland, USA.

*Akinrele IA. 1986. Hydrocyanic acid hazard during large scale cassava processing. *Trop Sci* 26:59-65.

*Alarie Y. 1985. The toxicity of smoke from polymeric materials during thermal decomposition. *Annu Rev Pharmacol Toxicol* 25:325-347.

*Aldridge WN. 1951. The conversion of cyanogen chloride to cyanide in the presence of blood proteins and sulphhydryl compounds. *Biochem J* 48:271-276.

*Aleksandrov NP, Korobov RN, Filippov VG. 1979. [Poison content in the blood and tissues of mammals after the inhalational uptake of carbon monoxide and benzene and cyanide vapors]. *Farmakol Toksikol* 42:670-674 [Russian].

*Anderson RA, Harland WA. 1982. Fire deaths in the Glasgow area III: The role of hydrogen cyanide. *Med Sci Law* 22:35-40.

- *Anderson RA, Watson AA, Harland WA. 1981. Fire deaths in the Glasgow area II: The role of carbon monoxide. *Med Sci Law* 21:288-294.
- *Anderson RA, tenHowe PW, Watson AA. 1982. Onderzoek naar de doodsoorzaken bij brand. *Ned T Geneesk* 126:182-188.
- *Anderson RA. 1986. Fire gases. In Curry AS, ed, *Analytical methods in human toxicology*, Part 2. MacMillan, Basingstoke, UK, pp 289-317.
- *Anonymous. 1953. Antidotes for cyanide. *Br Med J* 1:36-37.
- *Anonymous. 1972. Food and cyanide. *Br Med J* 2:481-482.
- *Anonymous. 1973. Chronic cassava toxicity. *Lancet* 2:245-246.
- *Anonymous. 1973. Stability of amyl nitrite in cyanide antidote kits. *Clin Toxicol* 6:133-134.
- *Anonymous. 1975. Use and toxicity of nitroprusside. *N Engl J Med* 292:1081-1082.
- *Anonymous. 1976. Cyanide intoxication from sodium nitroprusside in anesthesia. *Med Lett Drugs Ther* 18:68.
- *Anonymous. 1978. Controlled intravascular sodium nitroprusside treatment. *Br Med J* 6140:784-785.
- *Anonymous. 2002. Cyanogen agents. *J R Army Med Corps* 148:383-386.
- *Antal J. 1894. Experimentelle Studien zur Behandlung der Vergiftung durch Blausäure-Einatmung. *Ungar Arch Med* 3:117-128 [Experimental].
- *Aquanno JJ, Chan KM, Dietzler DN. 1981. Accidental poisoning of two laboratory technologists with sodium nitrite. *Clin Chem* 27:1145-1146.
- *Archer AW, Geyer R. 1982. Production of cyanide in a sodium nitrite poisoning case using an acid distillation screening test. *J Forensic Sci Soc* 22:333-334.
- *Asahina S, Friedman MA, Arnold E, Millar GN, Mishkin M, Bishop Y, Epstein SS. 1971. Acute synergistic toxicity and hepatic necrosis following oral administration of sodium nitrite and secondary amines to mice. *Cancer Res* 31:1201-1205 [Experimental].
- *Asmus E, Garschagen H. 1953. Über die Verwendung der Barbitursäure für die photometrische Bestimmung von Cyanid und Rhodanid. *Z Anal Chem* 138:414-422 [Analytical method].

*Ast DB, Bushel A, Chase HC. 1950. A clinical study of caries prophylaxis with zinc chloride and potassium ferrocyanide. *J Am Dent Assoc* 41:437-442.

*Atkins D. 1977. Cyanide toxicity following nitroprusside-induced hypotension. *Can Anaesth Soc J* 24:651-660.

*ATSDR. 1998. Toxicological profile for cyanide. Harper C, Agency for Toxic Substances and Disease Registry, Division of Toxicology, Atlanta, GA, USA and Goldhaber S, Research Triangle Institute, Research Triangle Park, NC, USA. US Department of Health and Human Services, Public Health Service, Atlanta, Georgia, USA [Review].

*Au W, Hsu TC. 1979. Studies on the clastogenic effects of biologic stains and dyes. *Environ Mutagen* 1:27-35 [Experimental].

*Bailey B. 2003. Are there teratogenic risks associated with antidotes used in the acute management of poisoned pregnant women? *Birth Defects Res A Clin Mol Teratol* 67:133-140.

*Ballantyne B. 1987b. Hydrogen cyanide as a product of combustion and a factor of morbidity and mortality from fires. Chapter 10. In Ballantyne B, Marrs TC, eds, *Clinical and experimental toxicology of cyanides*. John Wright, Bristol, UK, pp 248-291.

*Ballantyne B. 1984. Relative toxicity of carbon monoxide and hydrogen cyanide in combined atmospheres. *Toxicologist* 4:69.

*Banea-Mayambu JP, Tylleskär T, Gitebo N, Matadi N, Gebre-Medhin M, Rosling H. 1997. Geographical and seasonal association between linamarin and cyanide exposure from cassava and the upper motor neurone disease konzo in former Zaire. *Trop Med Intern Health* 2:1143-1151 [Chronic toxicity].

*Banea-Mayambu JP, Tylleskär T, Tylleskär K, Gebre-Medhin M, Rosling H. 2000. Dietary cyanide from insufficiently processed cassava and growth retardation in children in the Democratic Republic of Congo (formerly Zaire). *Ann Trop Paediatr* 20:34-40 [Chronic toxicity].

*Barillo DJ, Goode R, Rush BF, Lin RL, Freda A, Anderson EJ. 1986a. Lack of correlation between carboxyhemoglobin and cyanide in smoke inhalation injury. *Curr Surg* 43:421-423.

*Barillo DJ, Rush BF, Goode R, Lin RL, Freda A, Anderson EJ. 1986. Is ethanol the unknown toxin in smoke inhalation injury? *Am Surg* 52:641-645.

*Barriot P, Baud FJ, Toffis V, Riou B, Astier A. 1988. Cyanide poisoning in fire-related smoke injury. *Chest* 94:87.

- *Bartelheimer E. 1962. Unterschiede der toxischen und cyanidantagonistischen Wirksamkeit von Kobaltchelate-Verbindungen (Co-Histidin und Co²-EDTA). Naunyn-Schmiedeberg's Arch Exp Pathol Pharmacol 243:254-268 [Experimental study on toxicity of cobalt edetate in rats].
- *Bartelheimer EW. 1962. Analyse der akuten Kobaltvergiftung im Tierversuch. Naunyn-Schmiedeberg's Arch Exp Pathol Pharmacol 243:237-253 [Experimental study].
- *Baselt RC, Cravey RH. 1990. Disposition of toxic drugs and chemicals in man, 3rd edition. Year Book Medical Publishers, Chicago, IL, USA, pp 224-229.
- *Baselt RC. 1982. Cyanide and nitrite. In Disposition of toxic drugs and chemicals in man, 2nd ed. Biomedical Publications, Davis, CA, USA, pp 209-214, 555-557 [Toxicity of sodium nitrite].
- *BASF. 1998. BASF chemical emergency medical guidelines. Cyanides (CN), information and recommendations A to D. Unpublished report E008-004. BASF, Ludwigshafen, Germany; BASF, Mount Olive, New Jersey, USA [Review].
- *Bastian G, Mercker H. 1959. Zur Frage der Zweckmäßigkeit der Inhalation von Amylnitrit in der Behandlung der Cyanidvergiftung. Naunyn-Schmiedeberg's Arch exp Path Pharmacol 237:285-295 [Experimental studies of amyl nitrite in animals and human volunteers].
- *Baud F. 2008. Intoxications par les fumées d'incendie. EMC (Elsevier Masson SAS, Paris), Toxicologie – Pathologie professionnelle 16-539-G-10.
- *Baud FJ, Barriot P, Toffis V, Riou B, Astier A. 1988. Cyanide poisoning in fire victims. Intensive Care Med 14:289.
- *Baud FJ, Barrot P, Riou B, Vicaut E, Astier A. 1992. Blood cyanide concentrations after smoke-inhalation – Reply. N Engl J Med 326:1362.
- *Baud FJ. 2009. Akute Vergiftungen mit Kohlenmonoxid und Zyaniden. Therapeutische Umschau 11:1-12 [Review].
- *Baumeister RG, Schievelbein H, Zickgraf-Rüdel G. 1975. Toxicological and clinical aspects of cyanide metabolism. Arzneimittelforschung 25:1056-1064.
- *Beaven GH, Holiday ER, Johnson EA, Ellis B, Petrow V. 1950. The chemistry of anti-pernicious anaemia factors. Part VI: The mode of combination of component alpha in vitamin B12. J Pharm Pharmacol 2:944-55.
- *Beck M, Monroe S. 1982. The Tylenol scare: The death of seven people who took the drug triggers a nationwide alert – and a hunt for a madman. Newsweek 100:32-36.

- *Benes V, Cerna V. 1959. Akrylonitril: Akute Toxizität und Wirkungs – Mechanismus. *J Hyg Epidemiol Microbiol Immunol* 3:106-116.
- *Benignus VA, Kafer ER, Muller KE, Case MW. 1987. Absence of symptoms with carboxyhemoglobin levels of 16-23% *Neurotoxicol Teratol* 9:345-348.
- *Benignus VA, Muller KE, Malott CM. 1990. Dose-effects functions for carboxyhemoglobin and behavior. *Neurotoxicol Teratol* 12:111-118 [Review].
- *Benko S, Petri G, Eisner Am, Kardos G, Szabo T, Bentzik M, Hetenyi G. 1951. A kaserletes helyi csontveloanoxia hatása a vörös verséjt számára [Effect of experimental local medullary hypoxia on the number of red blood corpuscles.]. *Magy Belorv Arch* 4:15-20.
- *Benner JP, Lawrence D, Brady W. 2009. Smoke signals. Recognition and treatment of combustion-induced cyanide toxicity. *JEMS* 34:56-63.
- *Berl WG, Halpin BM. 1979. Human fatalities from unwanted fires. National Bureau of Standards, Washington, DC, Center for Fire Research, NBS-GCR-79-168.
- *Berlin G, Brodin B, Hilden JO, Mårtensson J. 1984. Acute dapsone intoxication: A case treated with continuous infusion of methylene blue, forced diuresis and plasma exchange. *Clin Toxicol* 22:537-548 [Other poisoning than CN].
- *Bertol E, Mari F, Orzalesi G, Volpato I. 1983. Combustion products from various kinds of fibers: Toxicological hazards from smoke exposure. *Forensic Sci Int* 22:111-116.
- *Betz H, Fruhling L. 1950. Action du cyanure de potassium sur la léthalité de souris irradiées a doses fractionnées. *C R Seances Soc Biol Fil* 144:595-597.
- *Betz H, Fruhling L. 1950. Étude de la régénération des organes hematopoietiques chez les souris irradiées a fortes doses et protégées par injection de KCN. *C R Seances Soc Biol Fil* 144:1013-10135.
- *Betz H, Hervé A. 1950. L'effet protecteur de KCN sur la léthalité de souris irradiées a diverses doses de rayons X. *C R Seances Soc Biol Fil* 144:1015-1017.
- *Betz H. 1950. Sur le mécanisme d'action protectrice du cyanure de potassium sur la léthalité des souris irradiées. *C R Seances Soc Biol Fil* 144:593-595.
- *Binet L, Wellers G, Dubrisay J. 1951. La valeur et le mécanisme de l'action antidote du tétrathionate de sodium dans l'intoxication cyanhydrique. *Presse Med* 59:641-643.

- *Birky M, Malek D, Paabo M. 1983. Study of biological samples obtained from victims of MGM Grand Hotel fire. *J Anal Toxicol* 7:265-271.
- *Birky MM, Clarke FB. 1981. Inhalation of toxic products from fires. *Bull NY Acad Med* 57:997-1013.
- *Birky MM, Halpin BM, Caplan YH, Fisher RS, McAllister JM, Dixon AM. 1979. Fire fatality study. *Fire Mater* 3:211-217 [Forensic medicine].
- *Birky MM, Paabo M, Brown J.E. 1979/80. Correlation of autopsy data and materials involved in the Tennessee Jail fire. *Fire Safety J* 2:17-22 [Forensic medicine].
- *Blake G. 1840. Observations and experiments in the mode in which various poisonous agents act on the animal body. *Edinburgh Med Surg J* 53:35-49 [Experimental].
- *Blanc P, Hogan M, Mallin K, Hryhorczuk D, Hessel S, Bernard B. 1985. Cyanide intoxication among silver-reclaiming workers. *JAMA* 253:367-371 [Chronic neurotoxicity].
- *Blaschke TE, Melmon KL. 1980. Antihypertensive agents and the drug therapy of hypertension. In Gilman AG, Goodman LS, Gilman A, eds. *The pharmacological basis of therapeutics*, 6th ed. MacMillan Publishing, New York, USA, 807-808 or 805-806?.
- *Blass N, Fung D. 1976. Dyed but not dead - Methylene blue overdose. *Anesthesiology* 45:458-459 [Mechanism of action against methemoglobinaemia and side effects].
- *Bodansky O, Gutman H. 1947. Treatment of methemoglobinemia. *J Pharmacol Exp Ther* 89:46-56 [Mechanism of action against methemoglobinaemia and side effects].
- *Boivin MJ. 1997. An ecological paradigm for a health behavior analysis of "konzo", a paralytic disease of Zaire from toxic cassava. *Soc Sci Med* 45:1853-1862 [Chronic toxicity].
- *Borron S. 2006. Blood cyanide and plasma cyanocobalamin concentrations after treatment of acute cyanide poisoning with hydroxocobalamin or vehicle in a canine model. *Clin Toxicol* 44:630.
- *Borron SW, Uhl W, Nolting A, Hostalek U. 2007. Interference of the cyanide antidote hydroxocobalamin with carboxyhemoglobin measurements should not limit clinical use in suspected or confirmed cyanide poisoning. *Ann Emerg Med* 50:624-625 [Comment on Lee et al, 2007]; author reply 626.
- *Borron SW, Favier C, Benaissa L, Baud FJ. 1999. Are soot deposits predictive of carbon monoxide poisoning in fire victims? *Clin Toxicol* 37:607.

- *Bose A. 1959. Modification of ^{59}Fe uptake in the circulating blood of lethally irradiated rats protected with cyanide. *Int J Radiat Biol* 1:383-386.
- *Boulin R, Uhry P, Nepveux, Eliachar. 1950. La reaction au bleu de Prusse chez les malades atteints de cirrhose bronzée [Prussian blue in hemochromatosis]. *Sem Hop* 26:4245-4246.
- *Boxer GE, Rickards JC. 1951. Chemical determination of vitamin B12. II. The quantitative isolation and colorimetric determination of millimicrogram quantities of cyanide. *Arch Biochem* 30:372-381.
- *Brandt-Rauf PW, Fallon LF, Tarantini T, Idema C. 1988. Health hazards of fire fighters: exposure assessment. *Br J Ind Med* 45 606-612.
- *Brewer GJ, Tarlov AR. 1961. Studies on the mechanism of primaquine-type hemolysis: The effect of methylene blue. *Clin Res* 9:65 [Mechanism of action against methemoglobinaemia and side effects].
- *British Pharmacopoeia. 1980. Hydroxocobalamin, Dicobat edetate. Her Majesty's Stationery Office, London, UK 229-230, 579.
- *Brown KL, Wilson RF, White MT. 2007. Carbon monoxide-induced status epilepticus in an adult. *J Burn Care Res* 28:533-536.
- *Brunnemann KD, Yu L, Hoffmann D. 1977a. Chemical studies on tobacco smoke. XLVII: Gas chromatographic determination of hydrogen cyanide and cyanogens in tobacco smoke. *J Anal Toxicol* 1:38-42 [Analytical study cited by IARC, 1986].
- *Burgess WA, Treitman RD, Gold A. 1979. Air contaminants in structural firefighting. Final Report to National Fire Prevention and Control Administration and the Society of Plastics Industry, Inc. Harvard School of Public Health, Boston.
- *Burrows GE, Liu DH, Isom GE, Way JL. 1982. Effect of antagonists on the physiologic disposition of sodium cyanide. *J Toxicol Environ Health* 10:181-189 [Study in mice].
- *Burrows GE, Liu DHW, Way JL. 1973. Effect of oxygen on cyanide intoxication: V. Physiologic effects. *J Pharmacol Exp Ther* 184:739-748 [Experimental study in dogs related to preference of oxygen via air ventilation with and without several antidotes].
- *Butenuth J, Schneider H, Schneider V. 1970. Spinale Mechanismen in einem Fall von Hirntod nach Cyanidintoxikation. *Dtsch Z Nervenheilkd* 197:255-284.
- *Büttner H, Fortwich F, Hansen HW. 1967. Behandlung einer schweren Nitrobenzolvergiftung mit Toluidinblau. *Verh Dtsch Ges Inn Med* 72:665-666 [Treatment of methemoglobinaemia].

- * CAC 1999. 1991. Codex standard for edible cassava flour (African regional standard). In Codex Alimentarius, Vol XII, Suppl 4 (Codex STAN 176) Food and Agriculture Organization of the United Nations, Rome, Italy [Cited by WHO, 1993].
- *Calabrese EJ. 1983. Principles of animal extrapolation. John Wiley, New York, USA [Experimental study].
- *Calafat AM, Stanfill SB. 2002. Rapid quantitation of cyanide in whole blood by automated headspace gas chromatography. J Chromatogr B Analyt Technol Biomed Life Sci 772:131-137.
- *California Medical Association. 1953. TREATMENT of cancer with laetriles: A report by the Cancer Commission of the California Medical Association. Calif Med 78:320-326.
- *Campian EC, Benz FW. 2008. The acute lethality of acrylonitrile is not due to brain metabolic arrest. Toxicology 253: 104-109[Abstract].
- *Caplan YH. 1977. Relationship of cyanide to deaths caused by fire. FPP TR31, The John Hopkins University, Applied Physics Laboratory, Laurel, Maryland, USA, 20810.
- *Carden E. 1970. Hyperbaric oxygen in cyanide poisoning. Anaesthesia 25:442-443.
- *Casadei E, Cliff J, Neves J. 1990. Surveillance of urinary thiocyanate concentration after epidemic spastic paraparesis in Mozambique. J Trop Med Hyg 93:257-261.
- *Castello C, Giussani A, Berti A. 1969. Avvelenamento da acido cianidrico: Considerazioni clinico-terapeutiche e presentazione di un caso [Poisoning with cyanhydric acid: Clinico-therapeutic considerations and presentation of a case]. Minerva Anestesiol 35:437-447.
- *Cha SI, Kim CH, Lee JH, Park JY, Jung TH, Choi WI, Han SB, Jeon YJ, Shin KC, Chung JH, Lee KH, Kim YJ, Lee BK. 2007. Isolated smoke inhalation injuries: Acute respiratory dysfunction, clinical outcomes, and short-term evolution of pulmonary functions with the effects of steroids. Burns 33:200-208.
- *Chan GM. 2006. Carbon monoxide poisoning, myocardial injury, and mortality. JAMA 295:2601 [Comment on Henry et al, 2006].
- *Chanas B, Wang H, Ghanayem BI. 2003. Differential metabolism of acrylonitrile to cyanide is responsible for the greater sensitivity of male vs female mice: Role of CYP2E1 and epoxide hydrolases. Toxicol Appl Pharmacol 193:293-302 [Abstract].
- *Chang SS, Chan YL, Wu ML, Deng JF, Chiu T, Chen JC, Wang FL, Tseng CP. 2004. Acute Cycas seed poisoning in Taiwan. Toxicol Clin Toxicol 42:49-54.

- *Chaturvedi AK, Smith DR, Canfield DV. 2001. Blood carbon monoxide and hydrogen cyanide concentrations in the fatalities of fire and non-fire associated civil aviation accidents, 1991-1998. *Forensic Sci Int* 121:183-188.
- *Chen KK, Rose CL, Clowes GHA. 1933. Methylene blue, nitrites and sodium thiosulfate against cyanide poisoning. *Proc Soc Exp Biol Med*, 31:250-252 [Experimental study in dogs].
- *Chiwona-Karlton L, Tylleskär T, Mkumbira J, Gebre-Medhin M, Rosling H. 2000. Low dietary cyanogen exposure from frequent consumption of potentially toxic cassava in Malawi. *Int J Food Sci Nutr* 51:33-43.
- *Chobanian SJ, Cattau EL, Winters C, Johnson DA, Van Ness MM, Miramadi A, Morwitz SL, Colcher H. 1987. In vivo staining with toluidine blue as an adjunct to the endoscopic detection of Barrett's esophagus. *Gastrointest Endosc* 33:99-101 [Not related to CN poisoning].
- *Chou KJ, Fisher JL, Silver EJ. 2000. Characteristics and outcome of children with carbon monoxide poisoning with and without smoke exposure referred for hyperbaric oxygen therapy. *Pediatr Emerg Care* 16:151-155.
- *Christel D, Eyer P, Hegemann M, Kiese M, Lorcher W, Weger N. 1977. Pharmacokinetics of cyanide in poisoning of dogs, and the effect of 4-dimethylaminophenol or thiosulfate. *Arch Toxicol* 38:177-190.
- *Christian GD. 1969. Medicine, trace elements, and atomic absorption spectroscopy. *Anal Chem* 41:24A-40A [Analytical study of cobalt in body fluids].
- *Clark CJ, Reid WH, Gilmour WH, Campbell D. 1986. Mortality probability in victims of fire trauma: revised equation to include inhalation injury. *Br Med J (Clin Res Ed)* 292:1303-1305.
- *Clark County Fire Department (CCFD). 1981. Clark County Fire Department report on MGM Grand Hotel fire. Clark County Fire Department, Las Vegas, Nevada.
- *Cliff J, Essers S, Rosling H. 1986. Ankle clonus correlating with cyanide intake from cassava in rural children from Mozambique. *J Trop Pediatr* 32:186-189 [Chronic poisoning].
- *Cliff J, Lundquist P, Rosling H, Sörbo B, Wide L. 1986. Thyroid function in a cassava-eating population affected by epidemic spastic paraparesis. *Acta endocrinologica* 113:523-528.
- *Cliff J, Lundqvist P, Mårtensson J, Rosling H, Sörbo B. 1985. Association of high cyanide and low sulphur intake in cassava-induced spastic paraparesis. *Lancet* 2:1211-1213.
- *Cliff J, Nicala D, Saute F, Givragy R, Azambuja G, Taela A, Chavane L, Howarth J. 1997. Konzo associated with war in Mozambique. *Trop Med Int Health* 2:1068-1074.

- *Cliff J, Nicala D, Saute F, Givragy R, Azambuja G, Taela A, Chavane L, Gani A. 1999. Ankle clonus and thiocyanate, linamarin and inorganic sulphate excretion in school children in communities with Konzo, Mozambique. *J Trop Pedriatics* 45:139-142.
- *Cliff J, Nicala D. 1997. Long term follow-up of konzo patients. *Trans Royal Soc Trop Med Hyg* 91:447-449 [Chronic poisoning].
- *Coentrão L, Neves A, Moura D. 2010. Hydroxocobalamin treatment of acute cyanide poisoning with a jewellery-cleaning solution. *BMJ Case Rep*: Published online. doi: 10.1136/bcr.01.2010.2603 or www.ncbi.nlm.nih.gov/pmc/articles/PMC3028453/ [Review of 9 KCN poisonings, published after the deadline].
- *Cohen MA, Guzzardi LJ. 1988. Toxic smoke inhalation and cyanide poisoning. *Am J Emerg Med* 6:203-205.
- *Cohen MA, Guzzardi LJ. 1988. Toxic smoke inhalation and cyanide poisoning. *Am J Emerg Med* 6:203-205.
- *Cole P. 1978. The safe use of sodium nitroprusside. *Anaesthesia* 33:473-477.
- *Condorelli L. 1959. [Peripheral vascular receptors with respiratory effects]. *Boll Soc Ital Biol Sper* 35:1973-1975 [Italian].
- *Conn EE. 1973. Cyanogenic glucosides (Committee on Food Protection, Food and Nutrition Board, National Research Council). In: *Toxicants occurring naturally in foods*, 2nd ed. Washington, DC, National Academy of Sciences 299-308 [Review, covered by individual papers].
- *Conn EE. 1979. Cyanogenic glycosides. *Int Rev Biochem Biochem Nutr* 1 A27:21-43 [Cited by Seeger and Neumann, 1988].
- *Conn EE. 1981. Biosynthesis of cyanogenic glycosides. In Vennesland B, Conn EE, Knowles CJ, Westley J, Wissing F, eds. *Cyanide in biology*, chapter 11. Academic Press, London, UK 183-196 [Review, no mention of treatment].
- *Conn JB, Norman SL, Wartman TG. 1951. The equilibrium between vitamin B12 (cyanocobalamin) and cyanide ion. *Science* 113:658-659.
- *Conway EJ. 1950. Microdiffusion analysis and volumetric error. Van Nostrand Reinhold, New York, USA [Analytical study].

- *Cooke RD, Coursey DG. 1981. Cassava, a major cyanide-containing food crop. In Vennesland B, Conn EE, Knowles CJ, Westley J, Wissing F, eds. Cyanide in biology, chapter 7. Academic Press, London, UK 11-28 [No mention of treatment].
- *Copeland AR. 1985. Accidental fire deaths. The 5-year Metropolitan Dade County experience from 1979 until 1983. *Z Rechtsmed* 94:71-79.
- *Cotton FA, Wilkinson G. 1966. Advanced inorganic chemistry, 2nd ed. Interscience, New York, USA 869 [Basic science].
- *Cousineau A Legg FG. 1935. Hydrocyanic acid gas and other toxic gases in commercial fumigation. *Am J Public Health* 25: 277-294 [General toxicity].
- *CPTC. 2009. Smoke, cyanide and carbon monoxide: The toxic twins of smoke inhalation. Smoke suppl 1-26. Cyanide Poisoning Treatment Coalition, Indianapolis, Indiana, USA.
- *Crampton RF, Gaunt IF, Harris R, Knowles JF, Langman MJ, Linnell JC, Matthews DM, Mollin DL, Pettigrew AR, Smith WT, Waters AH, Wilson J, Wise IJ. 1979. Effects of low cobalamin diet and chronic cyanide toxicity in baboons. *Toxicology* 12:221-234 [Experimental study].
- *Curry SC, Patrick HC. 1991. Lack of evidence for a percent saturation gap in cyanide poisoning. *Ann Emerg Med* 20:523-528.
- *Curry SC, Carlton MW, Raschke RA. 1997. Prevention of fetal and maternal cyanide toxicity from nitroprusside with coinfusion of sodium thiosulfate in gravid ewes. *Anesth Analg* 84:1121-1126.
- *Cutting WA. 1978. Cassava - a valuable food but a possible poison. *Trop Doct* 3:102-103.
- *Danes B, Leinfelder PJ. 1951. Cytological and respiratory effects of cyanide on tissue cultures. *J Cell Physiol* 37:427-446.
- *Danes B, Leinfelder PJ. 1951. Cytological and respiratory effects of cyanide on tissue cultures. *J Cell Physiol* 37:427-446.
- *Daunderer M. 1979. Todliche Rauchvergiftung mit Blausäure durch Schwelbrände. *Fortschr Med* 97:1401-1405.
- *Daunderer M. 1980. Antidot-Therapie: Toluidinblau bei Methämoglobinämie. *Fortschr Med* 98:462-464 [Treatment of methemoglobinaemia].

*Deichmann WB, Gerarde HW. 1969. Toxicology of drugs and chemicals. Academic Press, New York, USA, London, UK 597 [General toxicology, toluidine blue suggested as an antagonist to heparin].

*Demling RH. 1993. Smoke inhalation injury. *New Horizons*, 1:422-434.

*DFG. 1958. Cyanwasserstoff. In Greim H, ed. 2000. Cyanwasserstoff, Gesundheitsschädliche Arbeitsstoffe, Toxikologisch-arbeitsmedizinische Begründung von MAK-Werten, 31st ed. Verlag Chemie, Weinheim, Germany [Review].

*DFG. 2001. Cyanwasserstoff. Kalium- und Natriumcyanid. In Greim H, ed, Gesundheitsschädliche Arbeitsstoffe, Toxikologisch-Arbeitsmedizinische Begründung von MAK-Werten (Maximale Arbeitsplatzkonzentrationen), 32nd ed. Wiley-VCH, Weinheim, Germany [Review].

*Dixon K. 1962. Spectrophotometric determination of sodiumthiosulfate in body fluids by use of an iodine-amylose complex. *Clin Chim Acta* 7:453-458 [Analytical study].

*Djerad A, Monier C, Houzé P, Borron SW, Lefauconnier JM, Baud FJ. 2001. Effects of respiratory acidosis and alkalosis on the distribution of cyanide into the rat brain. *Toxicol Sci* 61:273-282.

*Downs JCU. 2005. Incidence and findings at postmortem. *Encycl Am VOL?*:305-319.

*Drasch G, von Meyer L, Kauert G, Weger N. 1982. Ein Fall einer vaginalen Cyanid-Beibringung. *Arch Kriminol* 170:21-28.

*Du Pont, 1981. Inhalation toxicity of common combustion gases. Haskell Laboratory, Unpublished report 238-81. Haskell Laboratory, E.I. du Pont de Nemours and Company. Newark, DE., USA [Cited by US-NAC, 2000a].

*Duque E, Duque J, Henao J, Mejia G, Arango J, Arroyave I, Pena L, Tobón R, Carvajal J, Zuluaga G, Garcia A, Sanín E, Gutiérrez J, Velásquez A, Arbeláez M. 2004. Organs transplanted from intoxicated donors. *Transplant Proc* 36:1632-1633.

*Eben A., Lewalter J. 1988. Cyanide determination in blood. In Angerer J, Schaller KH, eds. Analyses of hazardous substances in biological materials, vol 2: Methods for biological monitoring. VCH Verlagsgesellschaft, Weinheim, Germany 133-144 [Analytical study].

*Eckert KG, Eyer P. 1983. Differences in the reactions of isomeric ortho- and para-aminophenols with haemoglobin. *Biochem Pharmacol* 32:1019-1027 [Analytical study].

- *El Ghawabi SH, Gaafar MA, El-Saharti AA, Ahmed SH, Malash KK, Fares R. 1975. Chronic cyanide exposure: A clinical, radioisotope, and laboratory study. *Br J Ind Med* 32:215-219 [Chronic cyanide poisoning].
- *Erdman AR. 2007. Is hydroxocobalamin safe and effective for smoke inhalation? Searching for guidance in the haze [Comment on Borron et al, 2007b]. *Ann Emerg Med* 49:814-816; Erratum in *Ann Emerg Med* 50:187.
- *Erdman AR. 2008. In reply [to Borron and Baud, 2007b]. *Ann Emerg Med* 50:110-111.
- *Ermans AM, Delange F, Van der Velden M, Kinthaert J. 1972. Possible role of cyanide and thiocyanate in the etiology of endemic cretinism. *Adv Exp Med Biol* 30:455-486 [Review].
- *Esposito FM, Alarie Y. 1988. Inhalation toxicity of carbon monoxide and hydrogen cyanide gases released during the thermal decomposition of polymers. *J Fire Sci* 6:195-242. doi: 10.1177/073490418800600303
- *Etteldorf JN. 1951. Methylene blue in treatment of methemoglobinemia in premature infants caused by marking ink: A report of eight cases. *J Pediatr* 38:24-27 [Mechanism of action against methemoglobonaemia and side effects].
- *Evelyn RA, Malloy HT. 1938. Microdetermination of oxyhemoglobin, methemoglobin and sulfhemoglobin in a single sample of blood. *J Biol Chem* 126:655-662 [Analytical method].
- *Eyer P, Kiese M. 1976. Biotransformation of 4-dimethylaminophenol: Reaction with glutathion and some properties of the reaction products. *Chem-Biol Interact* 14:165-178 [Experimental study].
- *Fairley A, Linton EC, Wild FE. 1934. The absorption of hydrocyanic acid vapour through the skin with notes on other matters relating to acute cyanide poisoning. *J Hyg* 34:283-294 [Early data; cited by ATSDR, 1998].
- *Famuyiwa OO, Akanji AO, Osuntokun BO. 1995. Carbohydrate tolerance in patients with tropical ataxic neuropathy--A human model of chronic cyanide intoxication. *Afr J Med Med Sci* 24:151-157 [Review].
- *Feigl F, Anger V, Oesper RE. 1966. Spot tests in organic analysis, 7th ed. Elsevier Science, Amsterdam 547-548 [Analytical study].
- *Feigl F, Feigl HE. 1949. On the reactivity of inner-complex-bonded palladium - A new specific test for cyanides in alkaline solutions. *Anal Chim Acta* 3:300-309 [Analytical study].

- *Feldstein M, Klendshoj NC. 1954. The determination of cyanide in biological fluids by microdiffusion analysis. *J Lab Chem Med* 44:166-170 [Analytical study].
- *Feldstein M, Klendshoj NC. 1954. The determination of volatile substances by microdiffusion analysis. *J Forensic Sci* 2:39-58 [Covered by Feldsten and Klendshoj, 1954aAnalytical study].
- *Fernando GC, Busuttil A. 1991. Cyanide ingestion: Case studies of four suicides. *Am J Forensic Med Pathol* 12:241-246 [Erratum by Fernando and Busuttil, 1993].
- *Fernando GC, Busuttil A. 1993. Cyanide ingestion: Case studies of four suicides. *Am J Forensic Med Pathol* 14:353 [Erratum to Fernando and Busuttil, 1991].
- *Fischer H, Masel H. 1969. Zur protrahierten Cyankalium-Vergiftung. *Munch Med Wochenschr* 111:941-943.
- *Flanagan RJ, Watson ID, Whelpton R. 2009. Fundamentals of analytical toxicology. John Wiley and Sons, Hoboken, New Jersey, USA [Review of basic principles and practical information on the analysis of drugs, poisons and other relevant analytes in biological specimens].
- *Fontana F. 1795. Treatise on the venom of viper; on the American poisons; and on the Cherry Laurel and some other vegetable poisons. J Skinner (trans), London [General review].
- *Forst AW, Felix W. 1953. Vergiftung mit Alkalizyanid bei anacidem Magen. *Z Biol* 106:77 80.
- *Fortin JL, Judic-Peureux V, Manzon C, Grimon D, Desmettre T, Capellier G. 2009. Intoxication cyanhydrique en milieu carcéral. *Urg Prat* 92:39-41.
- *Fortin JL, Ruttiman M, Domanski L, Kowalski JJ. 2004. Hydroxocobalamin: Treatment for smoke inhalation-associated cyanide poisoning. Meeting the needs of fire victims. *JEMS* 29:18 21
- *Fortin JL. 2007. Acide cyanhydrique: Un risque méconnu pour le sapeur-pompier en intervention. *Urg Prat* 82:45-47.
- *Fortin JL, Ruttimann M, Capellier G, Bigorie A, Ferlicot S, Thervet E. 2007. Successful organ transplantation after treatment of fatal cyanide poisoning with hydroxocobalamin. *Clin Toxicol (Phila)* 45:468-471
- *Frankenberg L, Sörbo B. 1973. Formation of cyanide from o-chlorobenzylidene malononitrile and its toxicological significance. *Arch Toxikol* 31:99-108.

*Frankenberg L. 1980. Enzyme therapy in cyanide poisoning: effect of rhodanese and sulfur compounds. *Arch Toxicol* 45:315-323.

*Freeman AG. 1972. Food and cyanide. *Br Med J* 3:49.

*Freeman AG. 1986. Optic neuropathy and chronic cyanide toxicity. *Lancet* 1:441-442.

*Friedman A, Tomankiewicz Z. 1983. Dynamika obrazu klinicznego i zapisu EEG w zatruciu cyjanowodorem [Clinical manifestations and EEG findings in hydrogen cyanide poisoning]. *Neurol Neurochir Pol* 17:659-663 [Polish].

*Friehoff FJ, Lobermann KH. 1952. Toluidinblau bei der Behandlung von toxischen Hämoglobinämien. *Ther Ggw* 91:446-450 [Treatment of methemoglobinaemia].

*Friehoff FJ, Lobermann KH. 1953. Zur Farbstofftherapie toxischer Hämoglobinämien unter Berücksichtigung der Pathophysiologie dieser Vergiftungen [Dye therapy of toxic haemoglobinaemias, bearing in mind the pathophysiology of such poisonings]. *Arzneimittelforschung* 3:616-619 [Treatment of methemoglobinaemia].

*Fukumoto Y, Nakajimo H, Uetake M, Matusyama A, Yoshida T. 1957. A study on the sense of smell with respect to potassium cyanide solution and its hereditary transmission. *Jpn J Hum Genet* 2:7-16 [Detection of cyanide by humans].

*Gajatto S. 1951. Tossicità del cianuro di potassio somministrato per via sottocutanea nel coniglio e limiti entro i quali è realizzabile un utile antidotismo locale tra cianuro di potassio e 2,3-dimercaptopropanolo [Toxicity of subcutaneous potassium cyanide in the rabbit and limits of a useful local antidotism between potassium cyanide and 2,3-dimercaptopropanol]. *Boll Soc Ital Biol Sper* 27:40-42.

*Garcia-Moreno C. 2009. Gender inequality and fire-related deaths in India. *Lancet* 373:1230-1231 [Comment on Sanghavi et al, 2009].

*Garnier R, Bismuth C, Riboulet-Delmas G, Efthymiou ML. 1981. Poisoning from fumes from polystyrene fire. *Br Med J (Clin Res Ed)* 283:1610-1611.

*Gast JH, Arai K, Aldrich FL. 1952. Quantitative studies on urinarythiosulfate excretion by healthy human subjects. *J Biol Chem* 196:875-884 [Excretion of thiosulfate in non-poisoned persons].

*Geller RJ, Barthold C, Saiers JA, Hall AH. 2006. Pediatric cyanide poisoning: Causes, manifestations, management, and unmet needs. *Pediatrics* 118:2146-2158.

- *Geller RJ, Barthold C, Sakers JA, Hall AH. 2006. Pediatric cyanide poisoning: Causes, manifestations, management, and unmet needs. *Pediatrics* 118; 2146-2158 [doi: 10.1542/peds.2006-1251; www.pediatrics.org/cgi/content/full/118/5/2146].
- *George G, Ramsay K, Rochester M, Seah R, Spencer H, Vijayasankar D, Vasicuro L. 2002. Facilities for chemical decontamination in accident and emergency departments in the United Kingdom. *Emerg Med J* 19:453-457.
- *Gerlach E, Luebben K. 1959. [Phosphate metabolism in pigeon and human erythrocytes under the influence of 2,4-dinitrophenol, sodium cyanide, monidoacetate and of thyroxin and triethylens melamine. Comparative studies using P32-marked orthophosphate]. *Pflugers Arch Gesamte Physiol Menschen Tiere* 269:520-554 [German].
- *German A, Carrere C, Hubert-Brigau J. 1959. [Influence of cyanidation and of heating on the characterization of cyanocobalamine added to liver extracts]. *Ann Pharm Fr* 17:408-19 [French].
- *Ghanayem BI, Farooqui MY, Elshabrawy O, Mumtaz MM, Ahmed AE. 1991. Assessment of the acute acrylonitrile-induced neurotoxicity in rats." *Neurotoxicol Teratol* 13:499-502 [Abstract].
- *Gifford RW. 1959. Current practices in general medicine. 7. Treatment of hypertensive emergencies including use of sodium nitroprusside. *Proc Staff Meet Mayo Clin* 34:387-94.
- *Gill JR, Goldfeder LB, Stajic M. 2003. The Happy Land homicides: 87 deaths due to smoke inhalation. *J Forensic Sci* 48:161-163.
- *Gold A, Burgess WA, Clougherty EV. 1978. Exposure of firefighters to toxic air contaminants. *Am Ind Hyg Assoc J* 39:534-539.
- *Goldfrank L, Bresnitz EA, Weisman RS, Lewin NA. 1984. The inhaled agents and other disorders of oxygen transport. In: Hanson W, ed. *Toxic emergencies*. Churchill Livingstone, New York, USA 204-211.
- *Goldstein F, Rieders F. 1951. Formation of cyanide in dog and man following administration of thiocyanate. *Am J Physiol* 167:47-51.
- *Goldstein F, Rieders F. 1953. Conversion of thiocyanate to cyanide by an erythrocytic enzyme. *Am J Physiol* 173:287-290 [Analytical study].
- *Goldstein GM, Doull J. 1971. Treatment of nitrite-induced methemoglobinemia with hyperbaric oxygen. *Proc Soc Exp Biol Med* 138:137-139 [Adverse effect of antidote].

- *Goodhart GL. 1994. Patient treated with antidote kit and hyperbaric oxygen survives cyanide poisoning. *South Med J* 87:814-816.
- *Goodman LS, Gilman A, eds. 1975. Hydroxocobalamin. In *The pharmacological basis of therapeutics*, 5th ed. MacMillan, New York, USA 1334.
- *Gosselin RE, Smith HP, Hodge HC, eds. 1984. Cyanide. In *Clinical toxicology of commercial products*, 5th ed. Williams & Wilkins, Baltimore, Maryland, USA 123-130.
- *Gosselin RE, Smith HP, Hodge HC, eds. 1984. Nitrite. In *Clinical toxicology of commercial products*, 5th ed. Williams & Wilkins, Baltimore, Maryland, USA III:314-319 [Review; infant case of sodium nitrite poisoning].
- *Grabois B. 1955. Fatal exposure to methyl cyanide. *NY State Dep Labor Div Ind Hyg Mon Rev* 34:7-8. [Covered by Amdur, 1959, who reports more details.]
- *Graeme KA, Curry SC, Bikin DS, Lo Vecchio FA, Brandon TA. 1999. The lack of transplacental movement of the cyanide antidote thiosulfate in gravid ewes. *Anesth Analg* 89:1448-1452.
- *Gramling P. 1984. Diagnosing acute cyanide poisoning. *Ann Emerg Med* 13:147.
- *Grand AF, Kaplan HR, Lee GH. 1981. Investigation of combustion atmospheres in real building fires. Southwest Research Institute, Final Report, SWRI Project 01-6067, USFA Grant 80027, Submitted to the Society of the Plastics Industry, Inc. And U.S. Fire Administration.
- *Grant WM. 1980. The peripheral visual system as a target. In: Spencer PS, Schaumburg HH, ed. *Experimental and clinical neurotoxicology*. Williams Wilkins, Baltimore, Maryland, USA 77-91 [Not targeted organ by acute cyanide].
- *Grant WM. 1986. Sodium nitrite. In *Toxicology of the eye*, 3rd ed. Charles C Thomas Publisher, Springfield, Illinois, USA, 840 [Sodium nitrite poisoning].
- *Green CR, Ham K, Tange JD. 1969. Kidney lesions induced in rats by p-aminophenol. *Brit Med J* 1:162-164 [Experimental study].
- *Greenberg DM. 1980. The case against laetrile: The fraudulent cancer remedy. *Cancer* 45:799-807 [Review].
- *Griffon H, Le Breton R. 1951. Sur un cas d'intoxication aiguë mortelle par ingestion massive de cyanure de potassium pur. *Ann Med Leg Criminol Police Sci Toxicol* 31:266-269.

- *Gruener N, Shuval HI, Behroozi K, Cohen S, Shechter H. 1973. Methemoglobinemia induced by transplacental passage of nitrites in rats. *Bull Environ Contam Toxicol* 9:44-48 [Experimental study].
- *Grzywa M, Brajerski W. 1971. Przypadkowe, ostre zatrucie cyjankiem sodu zakończone wyleczeniem [Cyanide poisoning connected with galvanization]. *Przegl Lek* 27:278-280.
- *Grzywa M, Cwiklińska T, Trojanowska M. 1969. Przypadkowe, ostre zatrucie cyjankiem sodu zakończone wyleczeniem [Successful treatment of acute accidental sodium cyanide poisoning]. *Pol Tyg Lek* 24:283-284.
- *Guidotti T. 2006. Acute cyanide poisoning in prehospital care: new challenges, new tools for intervention. *Prehosp Disaster Med* 21:s40-48.
- *Habal R, Roldan CJ. 2008. Toxicity, cyanide. Hnatiuk OW, Talavera F, Crausman RS, Rice TD, Pinsky MR, eds. In Griffing GT, Pinsky MR. Critical care articles. eMedicine, Omaha Nebraska, USA. <http://emedicine.medscape.com/article/165886-overview> [Overview]..
- *Haber F, Weiss JJ. 1934. The catalytic decomposition of hydrogen peroxide by iron salts. *Proc R Soc Lond A* 147:332-351 [Experimental study].
- *Hagler L, Coppes RI. 1982. Inhibition of methemoglobin and metmyoglobin reduction by cobalt. *Biochem Pharmacol* 31:1779-1782 [Experimental study].
- *Haguenoer JM, Dequidt J, Jacquemont MC. 1975. Intoxications expérimentales par l'acétonitrile, 4ème note: Influence de l'hydroxocobalamine sur l'intoxication à moyen terme. *J Eur Toxicol* 2:113-121 [Does NOT deal with acute toxicity].
- *Haley TJ, Stolarsky F. 1951. A study of the acute and chronic toxicity of toluidine blue and related phenazine and thiazine dyes. *Stanford Med Bull* 9:96-100.
- *Hall AH, Kulig KW, Rumack BH. 1986. Drug- and chemical-induced methaemoglobinaemia: Clinical features and management. *Med Toxicol* 1:253-260.
- *Halpin BM, Berl WG. 1978. Human fatalities from unwanted fires. Report NBS-GCR-79-168, National Bureau of Standards, Washington, DC, USA.
- *Hantson P, Butera R, Clemessy JL, Michel A, Baud FJ. 1997. Early complications and value of initial clinical and paraclinical observations in victims of smoke inhalation without burns. *Chest* 111:671-675.

*Haouach H, Fortin JL, Chaybany B, Essouri S, Berton L, Catineau J, Galinski M, Lapostolle F. 2005. Prehospital use of hydroxocobalamin in children exposed to fire smoke. *Ann Emerg Med* 46:S30.

*Harrison MR. 1977. Toxic methemoglobinemia: A case of acute nitrobenzene and aniline poisoning. *Anaesthesia* 32:270-272. [Other poisoning, methylene blue].

*Hart GB, Strauss MB, Lennon PA, Whitcraft DD. 1985. Treatment of smoke inhalation by hyperbaric oxygen. *J Emerg Med* 3:211-215.

*Hartelius H. 1950. Cyannatriumforgiftning och dess framgångsrika behandling [Successful treatment in a case of sodium cyanide poisoning]. *Nord Med* 44:1747-1749 [Swedish].

*Hartzell GE, Packham SC, Switzer WF. 1983. Toxic products from fires. *Am Ind Hyg Assoc J* 44:248-255.

*Hartzell GE, Switzer WG, Priest DN. 1985. Modeling of toxicological effects of fire gases: V. Mathematical modelling of intoxication of rats by combined carbon monoxide and hydrogen cyanide atmospheres. *J Fire Sci* 3:330-342 [Experimental study].

*Harvey SC. 1980. Sodium nitrite. In Osol A, ed, *Remington's pharmaceutical science*, 16th ed. Mack Publishing, Easton, Pennsylvania, USA, p 782 [Synthesis of sodium nitrite].

*Hauschild F. 1953. Tödliche Kalkstickstoffvergiftung und die Frage des gestörten Alkoholabbaues. *S Afr J Clin Sci* 14:311-320.

*Henning G, Mittag G, Schmidt HH, Klost W. 1973. Erfahrungen mit Sonderlehrgängen für Erste Hilfe bei Zyan-Vergiftungen im Betrieb. *Zentralbl Arbeitsmed* 23:1-4.

*Henry CR, Satran D, Lindgren B, Adkinson C, Nicholson CI, Henry TD. 2006. Myocardial injury and long-term mortality following moderate to severe carbon monoxide poisoning. *JAMA* 295:398-402.

*Henry CR, Satran D, Lindgren B, Adkinson C, Nicholson CI, Henry TD. 2006b. Myocardial injury and long-term mortality following moderate to severe carbon monoxide poisoning. In reply [to Chan, 2006; Perez et al, 2006]. *JAMA* 295:2602.

*Hicks SP. 1950. Brain metabolism in vivo; the distribution of lesions caused by cyanide poisoning, insulin hypoglycemia, asphyxia in nitrogen and fluoroacetate poisoning in rats. *AMA Arch Pathol* 49:111-137.

*Higgins EA, Fiorca V, Thomas AA, Davis HV. 1972. Acute toxicity of brief exposures to HF, HCl, NO₂, and HCN with and without CO. *Fire Technol* 8:120-130 [Cited by US-NAC, 2000a].

- *Hill IR. 1989. Incapacitation and fires. *Am J Forens Med Pathol* 10:49-53.

- *Höbel M, Kreye VA, Nemes Z, Pill J. 1978. Organotoxic effects of excessive doses of sodium nitroprusside in the rabbit. *Klin Wochenschr* 56 Suppl 1:153-159 [Experimental study].

- *Hoffman RS, Sauter D. 1989. Methemoglobinemia resulting from smoke inhalation. *Vet Hum Toxicol* 31:168-170.

- *Hogod C. 1979. Effect of exposure to 0.5 ppm hydrogen cyanide singly or combined with 200 ppm carbon monoxide and/or 5 ppm nitric oxide on coronary arteries, aorta, pulmonary artery, and lungs in the rabbit. *Int Arch Occup Environ Health* 44:13-23.

- *Holmes RK, Way JL. 1982. Mechanism of cyanide antagonism by sodium nitrite. *Pharmacologist* 24:182 [Abstract, experimental study].

- *Hoppe-Seyler F. 1876. Über die Prozesse der Gährungen und ihre Beziehung zum Leben der Organismen. *Pflügers Arch gesamte Physiol Menschen Tiere* 12:1-17 [Early report that cyanide inhibits tissue oxidation reactions].

- *Houeto P, Borron SW, Baud FJ. DATE. Early death due to smoke inhalation: Role of volatile organic compounds. ?? thesis. Hôpital Lariboisière, Université Paris VII, Paris, France.

- *Houeto P, Buneaux F, Galliot-Guilley M, Baud FJ, Levillain P. 1994. Determination of hydroxocobalamin and cyanocobalamin by derivative spectrophotometry in cyanide poisoning. *J Analyt Toxicol* 18:154-158. WS thesis

- *Houeto P, Borron SW, Sandouk P, Imbert M, Levillain P, Baud FJ. 1996. Pharmacokinetics of hydroxocobalamin in smoke inhalation victims. *Clin Toxicol*, 34:397-404.

- *Houeto P, Borron SW, Baud FJ, Muszynski J, Buisine A, Gourelain H, Cheftel E, Marlière F. 1999. Assessment of erythrocyte cholinesterase activity in victims of smoke inhalation. *Clin Toxicol* 37:321-326.

- *Houeto P, Borron SW, Marlière F, Baud FJ, Levillain P. 2001. Development of a method for measuring volatile organic compounds in the blood of fire victims using 'purge and trap' gas chromatography. *Indoor Built Environ* 10:62-69.

- *Hrushesky WJ, Olshefski R, Wood P, Meshnick S, Eaton JW. 1985. Modifying intracellular redox balance: An approach to improving therapeutic index. *Lancet* 1:565-567 [Experimental study, methylene blue].

- *Hug E, Marzeni AD. 1933a. La fixation de l'acide cyanhydrique par les hématies qui contiennent de la méthémoglobine. *C R Séances Soc Biol Fil* 114:84-86 [Experimental study].

*Hug E, Marzeni AD. 1933b. Mécanisme de l'action antidote du sodium vis-à-vis de l'intoxication par l'acide cyanhydrique. C R Séances Soc Biol Fil 114:86-87 [Experimental study].

*Hug E. 1932. L'intoxication par l'acide cyanhydrique. Action antidote du bleu de méthylène, du nitrite de sodium et du sulfure de sodium. C R Séances Soc Biol Fil 111:88-90 [Experimental study].

*Hug E. 1933. Action combinée du nitrite de sodium dans le traitement de l'intoxication cyanhydrique chez le lapin. C R Séances Soc Biol Fil 114:87-89 [Experimental study].

*Iakovlev BV. 1959. Opyt tsianizatsii passazhirskikh vagonov uskorennym sposobom [Experience with a rapid method of cyanide disinfection of passenger railroad cars]. Gig Sanit 24:85-86 [Russian].

*Ihedioha JI. 2002. The clinicopathologic significance of enriching grated cassava mash with red palm oil in the production of gari. Plant Foods Hum Nutr 57:295-305.

*Iida T, Tanaka M. 1953. Studies on the components of the fushun shale oil III: Nitriles from the crude light oil. Pharm Bull 1:211-212.

*Ikegaya H, Iwase H, Hatanaka K, Sakurada K, Yoshida K, Takatori T. 2001. Diagnosis of cyanide intoxication by measurement of cytochrome c oxidase activity. Toxicol Lett 119:117-123.

*IPCS. 2003. Hydrogen cyanide, liquefied. International Chemical Safety Card 0492. International Programme on Chemical Safety, World Health Organization, Geneva, Switzerland [Review].

*Isom GE, Burrows GE, Way JL. 1982. Effect of oxygen on the antagonism of cyanide intoxication-cytochrome oxidase, *in vivo*. Toxicol Appl Pharmacol 65:250-256.

*Isom GE, Way JL. 1974. Effect of oxygen on cyanide intoxication VI: Reactivation of cyanide-inhibited glucose metabolism. J Pharmacol Exp Ther 189:235-243.

*Israeli A. 1988. Management of cyanide poisoning. Ann Emerg Med 17:108-109.

*Italian Pharmacopoeia. 1985. Hydroxocobalamin. In Farmacopea Ufficiale della Repubblica Italiana, 9th ed. Permanent Commission of the Italian Pharmacopoeia, Istituto Superiore di Sanità, Roma, Italy. Istituto Poligrafico e Zecca dello Stato, Rome, Italy.

*ITI. 1985. Sodium nitrite. In Toxic and hazardous industrial chemicals safety manual. International Technical Information Institute, Tokyo, Japan, p. 485.

- *Ivankovich AD, Braverman B. 1988. Sodium thiosulphate decreases blood cyanide concentration following sodium nitroprusside. *Br J Anaesth* 60:744-746.
- *Ives JM, Hughes EE, Taylor HK. 1971. Toxic atmospheres associated with real fire situations. Report NBS 10807, National Bureau of Standards, Washington, DC, USA.
- *Izmerov NF, ed. 1982. IRPTC scientific reviews of soviet literature on toxicity and hazards of chemicals 22: Nitrites. Centre of International Projects, GKNT, Moscow, Russia.
- *Jackson LC. 1988. Possible adaptation to serum thiocyanate overload associated with chronic sublethal dietary cyanide ingestion. *Hum Biol* 60:615-622.
- *Jagnoor J, Ivers R, Kumar R, Jha P. 2009. Fire-related deaths in India: How accurate are the estimates? *Lancet* 2009 374:117; author reply 118 [Comment on Sanghavi et al, 2009].
- *Janiszewska M, Kostanecki W. 1970. Zmiany skorne w przypadku ostrego smiertelnego zatrucia cjankiem sodu [Skin lesions in a case of acute fatal sodium cyanide poisoning]. *Przegl Dermatol* 57:755-757 [Polish].
- *Jankovic J, Jones W, Burkhart J, Noonan G. 1991. Environmental study of firefighters. *Ann Occup Hyg* 35:581-202.
- *Järvalho B, Lavenius B, Sällsten G. 1986. Cancer morbidity in workers exposed to cutting fluids containing nitrites and amines. *Br J Ind Med* 43:563-565 [Not related to CN poisoning] .
- *Jensen PH, Nielsen SV. 1981. Cyankaliumforgiftning [Potassium cyanide poisoning]. *Ugeskr Laeger* 143:2978 [Danish].
- *Johanning RJ, Zaske DE, Tschida SJ, Johnson SV, Hoey LL, Vance-Bryan K. 1995. A retrospective study of sodium nitroprusside use and assessment of the potential risk of cyanide poisoning. *Pharmacotherapy* 15:773-777.
- *Johnstone RT. 1948. Aliphatic nitrile compounds. In *Occupational medicine and industrial hygiene*, Mosby, St Louis, Missouri, USA, pp 130-145.
- *Jones GR, Israel MS. 1970. Mechanism of toxicity of injected CS gas. *Nature* 228:1315-1317.
- *Jones GRN. 1988. Cyanide and fire victims. *Lancet* 1988:457.
- *Jones J, McMullen MJ, Dougherty J. 1987. Toxic smoke inhalation: Cyanide poisoning in fire victims. *Am J Emerg Med* 5:317-321.

- *Jouglard J, Fagot G, Deguigne B, Arlaud JA. 1971. L'intoxication cyanhydrique aiguë et son traitement d'urgence. *Mars Med* 108:571-575.
- *Judkins K. 1989. Algorithm for managing injury from smoke inhalation. *BMJ* 299:1216 [Comment on Langford and Armstrogn, 1989].
- *Juskowa J. 1972. Electrocardiogram in poisoning with mercuric oxycyanide. *Pol Med J* 11:297-303.
- *Kales SN, Christiani DC. 2004a. Acute chemical emergencies. *N Engl J Med* 350:800-808.
- *Kales SN, Christiani DC. 2004b. Acute chemical emergencies. *N Engl J Med* 350:2103-2104 [Reply to Nogué-Xarau et al, 2004].
- *Kalmus H, Hubbard DJ. 1960. The chemical senses in health and disease. Charles C. Thomas, Springfield, Illinois, USA [Many people unable to perceive odour of HCN].
- *Kapamadzija B. 1970. Pokušaj samoubistva kalijum cijanidom (Suicidolosko-psihopatološki prikaz) [Suicide attempt with potassium cyanide. (Suicidologic-psychopathologic presentation)]. *Neuropsihijatrija* 18:347-352 [Croatian].
- *Karalliedde L, Wheeler H, Maclehose R, Murray V. 2000. Possible immediate and long-term health effects following exposure to chemical warfare agents. *Public Health* 114:238-248.
- *Katsumata Y, Sato K, Yada S, Suzuki O, Yoshino M. 1983. Kinetic analysis of anaerobic metabolism in rats during acute cyanide poisoning. *Life Sci* 33:151-155.
- Kedderis GL, Strother DE. 2007. The bioactivation and acute toxicity of acetonitrile. Unpublished manuscript, draft. BP, Nitriles Toxicology and Health Risk Assessment, Arlington, Virginia, USA [Draft review].
- *Keilin J. 1950. Reactions of haems with cyanides, isocyanides and amino acids. *Nature* 165:151.
- *Kempeneers J, Lefebvre L. 1953. Un cas d'intoxication mercurielle. *Rev Med Liege* 8:770-772.
- *Kerns W, Beuhler M, Tomaszewski C. 2008. Hydroxocobalamin versus thiosulfate for cyanide poisoning. *Ann Emerg Med* 51:338-339 [Comment on Hall et al, 2007].
- *Khot UN, Novaro GM, Popović ZB, Mills RM, Thomas JD, Tuzcu EM, Hammer D, Nissen SE, Francis GS. 2003. Nitroprusside in critically ill patients with left ventricular dysfunction and aortic stenosis. *N Engl J Med* 348:1756-1763..

- *Kiese M, Lörcher W, Weger N, Zierer A. 1972. Comparative studies on the effects of toluidine blue and methylene blue on the reduction of ferrihaemoglobin in man and dog. *Europ J Clin Pharmacol* 1972; 4, 115-118 [Mechanism of action against methemoglobonaemia and side effects].
- *Kiese M, Waller HD. 1951. [Reduction of haemoglobin and oxygen for reversibly reducible dyes in red cells.] *Naunyn-Schmiedebergs Arch Pharmakol* 213:44 [German; treatment of methemoglobinaemia].
- *Kiese M. 1945. [Reduction of haemoglobin. IV.] *Naunyn-Schmiedebergs Arch Exp Pathol Pharmakol* 204:288-312. [German; treatment of methemoglobinaemia].
- *Kiese M. 1974. Methaemoglobinaemia, a comprehensive treatise. CRC Press, Cleveland, Ohio, USA [Other poisoning].
- *Kimmerle, G. 1974. Aspects and methodology for the evaluation of toxicological parameters during fire exposure. *Combust Toxicol* 1:4-51 [Cited by US-NAC, 2000a].
- *Kirk RL, Stenhouse NS. 1953. Ability to smell solutions of potassium cyanide. *Nature* 171:698-699 [Detection study].
- *Kizer KW. 1984. Hyperbaric oxygen and cyanide poisoning. *Am J Emerg Med* 2:113.
- *Klimmek R, Roddewig C, Fladerer H, Weger N. 1982. Cerebral blood flow, circulation, and blood homoeostasis of dogs during slow cyanide poisoning and after treatment with 4-dimethylaminophenol. *Arch Toxicol* 50:65-76.
- *Klír P, Strejc P, Neuwirthová H, Nazundarová K. 1976 . Nálezy u protrahované otravy kyanidy [Findings in protracted cyanide poisoning]. *Soud Lek* 21:25-28 [Czech].
- *Korhonen A, Hemminki K, Vainio H. 1983. Toxicity of rubber chemicals towards three-day chicken embryos. *Scand J Work Environ Health* 9:115-119 [Experimental study].
- *Kulling P. 1992. Hospital treatment of victims exposed to combustion products. *Toxicol Lett* 64-65:283-289.
- *Kulling P. 1992. Hospital treatment of victims exposed to combustion products. *Toxicol Lett* 64/65:283-289.
- *Kun E, Fanshier DW. 1959. Enzymic transfer of sulfur from beta-mercaptopyruvate to cyanide. *Biochim Biophys Acta* 33:26-28.

- *Kuroki T, Abbondandolo A, Drevon C, Huberman E, Laval F. 1980. Mutagenesis assays with mammalian cells. IARC Monogr Eval Carcinog Risk Chem Hum Suppl 2 Suppl:107-133 [Experimental study on sodium nitrite].
- *Lafin SM, Lucchesi M, Fedak M, Barone JA. 1992. A need to revise the cyanide antidote package instructions. J Emerg Med 10:623-624.
- *Laforge M, Gourlain H, Fompeydie D, Buneaux F, Borron SW, Galliot-Guilley M. 1999. Ferrocyanide ingestion may cause false positives in cyanide determination. J Toxicol Clin Toxicol 37:337-340.
- *Landa Z, Klouda P, Pleskotova D. 1965. In Veleminsky J, Gichner T, Ridesova A, eds, Induction of mutations and the mutation process. Proceeding of a Symposium, Prague, 26-28 September 1963. Czechoslovak Academy of Sciences Publishing House, Prague, Czechoslovakia, pp 115-122 [Experimental study on toluidine blue].
- *Langford RM, Armstrong RF. 1989. Algorithm for managing injury from smoke inhalation. BMJ 299:902-905.
- *Lavon O, Bentur Y. 2010. Does amyl nitrite have a role in the management of pre-hospital mass casualty cyanide poisoning? Clin Toxicol (Phila) 48:477-484 [Review].
- *Le Lay J. 1950. Possibilités d'intoxication par les cyanures alcalins et l'acide cyanhydrique dans l'industrie metallurgique. Méd Usine Rev Hyg Ind Mal Prof 12:22-25.
- *Lee CW, Tam JC, Kung LK, Yim LK. 2003. Validity of CO-oximetric determination of carboxyhaemoglobin in putrefying blood and body cavity fluid. Forensic Sci Int 132:153-156.
- *Lee HM, Reed J, Greeley GH, Englander EW. 2009. Impaired mitochondrial respiration and protein nitration in the rat hippocampus after acute inhalation of combustion smoke. Toxicol Appl Pharmacol 235:208-215 [Experimental study].
- *Lehmann P. 1977. Laetrile: The fatal 'cure'. FDA Consum 11:10-15.
- *Leumann EP, Brandenberger H. 1977. Hemodialysis in a patient with acute mercuric cyanide intoxication. Concentrations of mercury in blood, dialysate, urine, vomitus, and feces. Clin Toxicol 11:301-308.
- *Leung P, Sylvester DM, Chiou F, Way LI, Way EL, Way JL (1986) Stereospecific effect of naloxone hydrochloride on cyanide intoxication. Toxicol Appl Pharmacol, 83:525-530 [Experimental study].

*Levin BC, Paabo M, Gurman JL, Harris SE. 1987. Effects of exposure to single or multiple combinations of the predominant toxic gases and low oxygen atmospheres in fires. *Fundam Appl Toxicol* 9:236-250.

*Levin BC, Rechani PR, Gurman JL, Landron F, Clark HM, Yoklavich MF, Rodriguez JR, Droz L, Mattos de Cabrera F, Kaye S. 1990. Analysis of carboxy-hemoglobin and cyanide in blood from victims of the Dupont Plaza Hotel fire in Puerto Rico. *J Forensic Sci* 35:151-168.

*Levin BC. 1996. New research avenues in toxicology: 7-gas N-Gas Model, toxicant suppressants, and genetic toxicology. *Toxicology* 115:89-106.

*Levine B, D'Nicuola J, Kunsman G, Smith M, Stahl C. 1996. Methodologic considerations in the interpretation of postmortem carboxyhemoglobin concentrations. *Toxicology* 115:129-134.

*Levine B, Moore KA, Fowler D. 2001. Interaction between carbon monoxide and ethanol in fire fatalities. *Forensic Sci Int* 124:115-116.

*Levine S, Klein M. 1959. Effect of chlorpromazine on cyanide intoxication. *Proc Soc Exp Biol Med* 102:192-194.

*Levine S, Wenk EJ. 1959. Cyanide encephalopathy produced by intravenous route. *J Nerv Ment Dis* 129:302-305.

*Levine,MS, Radford EP. 1978. Occupational exposures to cyanide in Baltimore fire fighters, *J Occup Med*, 20:53-56 [Chronic toxicity].

*Lewis SA, O'Haver TC, Harnly JM. 1985. Determination of metals at the microgram-per-liter level in blood serum by simultaneous multielement atomic absorption spectrometry with graphite furnace atomization. *Anal Chem* 57:2-5 [Analysis of cobalt].

*Leybell I, Hoffman RS, Baud FJ, Borron SW. 2006. Toxicity, cyanide. In Lee DC, van de Voort JT, Benitez JG, Halamka J, Tarabar A, eds. *eMedicine specialities*, Omaha Nebraska, USA. [Overview; superseded by Leybell et al, 2008]..

*Leybell I, Hoffman RS, Borron SW. 2008. Toxicity, cyanide. Lee DC, van de Voort JT, Benitez JG, Halamka J, Tarabar A, eds. In Adler J, ed. *Emergency medicine articles, toxicology*. eMedicine, Omaha Nebraska, USA. <http://emedicine.medscape.com/article/814287-overview> [Overview]..

*Li CG, Rand MJ. 1993. Effects of hydroxocobalamin and haemoglobin on no-mediated relaxations in the rat anococcygeus muscle. *Clin Exp Pharmacol Physiol* 20:633-40

*Li FP. 1969. Suicide among chemists. *Arch Environ Health* 19:518-520.

- *Linnell JC, Wilson J, Crampton RF, Smith WT, Knowles JF, Gaunt IF, Wise IJ, Matthews DM. 1979. Effects of low cobalamin diet and chronic cyanide toxicity on cobalamin distribution in baboons. *Toxicology* 14:81-90 [Experimental study].
- *Lochner W, Mercker H, Nasser M. 1959. Über den anaeroben. Energiegewinn des Warmblüterherzens in situ unter Cyanidvergiftung. *Naunyn Schmiedebergs Arch Exp Pathol Pharmacol* 236:365-381.
- *Ludwin SK, Northway WH, Bensch KG. 1974. Oxygen toxicity in the newborn. Necrotizing bronchiolitis in mice exposed to 100% oxygen. *Lab Invest* 31:421-435 [Other poisoning].
- *Lumsden CE. 1950. Cyanide leukoencephalopathy in rats and observations on the vascular and ferment hypotheses of demyelinating diseases. *J Neurol Neurosurg Psychiatry* 13:1-15.
- *Lundquist P, Rammer L, Sörbo B. 1989. The role of hydrogen cyanide and carbon monoxide in fire casualties: A prospective study. *Forensic Sci Int* 43:9-14.
- *Lundquist P, Rosling H, Sörbo B. 1985. Determination of cyanide in whole blood, erythrocytes and plasma. *Clin Ther* 31:591-595 [Analytical study].
- *Lynch EL, Thomas TL. 2004. Pediatric considerations in chemical exposures: are we prepared? *Pediatr Emerg Care* 20:198-208.
- *Lynch RD. 1975. On the non-existence of synergism between inhaled hydrogen cyanide and carbon monoxide. Fire Research Note No.1035, Department of the Environment and Fire Offices' Committee, Joint Fire Research Organization.
- *MacDonald IA, Rao BG. 1982. Nitrite-induced volatile fecal mutagens: Some preliminary studies. *Cancer* 49:1405-1408 [Experimental study].
- *Mack RB. 1983. Aha!!, The smell of bitter almonds ... cyanide poisoning. *N C Med J* 44:27-28.
- *MacQuiston TAC. 1936. Fatal poisoning by sodium nitrite. *Lancet* 2:1153-1154 [Other poisoning].
- *Maeda H, Fukita K, Oritani S, Nagai K, Zhu BL. 1996. Evaluation of post-mortem oxymetry in fire victims. *Forensic Sci Int* 81:201-209.
- *Magendie F. 1817. Sur l'emploi de l'acide prussique dans le traitement de plusieurs maladies de poitrine, et particulièrement dans la phtisie pulmonaire. *Ann Chim Phys* 6:347-360 [Therapeutic use of cyanide].

- *Majdevac M, Sovljanski R, Kapamadzija B. 1975. Trovanje kalijevim cijanidom [Poisoning by potassium cyanide]. *Arh Hig Rada Toksikol* 26:151-155 [Croatian].
- *Makene WJ, Wilson J. 1972. Biochemical studies in Tanzanian patients with ataxic tropical neuropathy. *J Neurol Neurosurg Psychiatry* 35:31-33.
- *Mansouri A. 1985. Methemoglobinemia. *Am J Med Sci* 289:200-209 [Other poisoning].
- *Manz A. 1968. Über einen Überlebensfall nach akuter Vergiftung durch ein Schwefelwasserstoff-Blausäuregemisch. *Zentralbl Arbeitsmed* 18:167-171.
- *Marigo M, Ferrara SD, Cima L, Giusti P, Floreani M. 1979. Some cases of acute intoxications from compulsive use of isoaminile. *Arch Toxicol* 42:295-301.
- *Marrs TC, Ballantyne B. 1987. Clinical and experimental toxicology of cyanides: An overview. Chapter 23 in Ballantyne B, Marrs TC, eds, *Clinical and experimental toxicology of cyanides*. John Wright, Bristol, UK, pp 473-495 [Overview].
- *Marrs TC, Bright JE, Inns RH. 1989. Methaemoglobin production and reduction by methylene blue and the interaction of methylene blue with sodium nitrite in vivo. *Human Toxicol* 8:359-364 [Mechanism of action against methemoglobonaemia and side effects].
- *Marrs TC, Bright JE, Swanston DW. 1982. The effect of prior treatment with 4-dimethylaminophenol (DMAP) on animals experimentally poisoned with hydrogen cyanide. *Arch Toxicol* 51:247-253 [Experimental study].
- *Marrs TC, Swanston DW, Bright JE. 1985. 4-Dimethylaminophenol and dicobalt edetate (Kelocyanor) in the treatment of experimental cyanide poisoning. *Hum Toxicol* 4:591-600 [Experimental study].
- *Martindale. 1989. Methylene blue, hydroxocobalamin, amyl nitrite. In Reynolds JEF, ed, *The extra pharmacopoeia*, 29th ed. Pharmaceutical Press, London, UK, pp 843-844, 1260-1261, 1492-1493 [More recent edition: Martindale, 2008].
- *Mathes K, Gross F. 1939. Über den Nachweis von Methämoglobin und Cyanmethämoglobin im strömenden Blut. *Naunyn-Schmiedebergs Arch Exp Pathol Pharmacol* 191:706-714 [Amyl nitrite did not yield methaemoglobin in human volunteers].
- *Max B. 1988. This and that: An odorless crime and the mystery of a poet's heart. *Trends Pharmacol Sci* 9:276-278.

- *Mayes RW. 1991. The toxicological examination of the victims of the British Air Tours Boeing 737 accident at Manchester in 1985. *J Forensic Sci* 36:179-184.
- *McAllister JL, Roby RJ, Levine B, Purser D. 2008. Stability of cyanide in cadavers and in postmortem stored tissue specimens: A review. *J Anal Toxicol* 32:612-620.
- *McAuley F, Reive DS. 1983. Rapid quantitation of cyanide in blood by gas chromatography. *J Anal Toxicol* 7:213-215 [Analytical study].
- *McGaughey CA. 1951. Hydrocyanic acid poisoning in nutrias caused by cassava (*Manihot utilissima* Pohl). *Br Vet J* 107:279-280.
- *McNamara BP. 1976. Estimates of the toxicity of hydrocyanid acid vapors in man. Technical report No. EB-TR-76023. Edgewood Arsenal Aberdeen Proving Ground. Department of the Army, Maryland, USA [Not therapeutic study].
- *Meister A. 1973. On the enzymology of amino acid transport. *Science* 180:33-39 [Experimental study].
- *Metz E, Balcerzak S, Sagone AL. 1976. Mechanism of methylene blue stimulation of the hexose-monophosphate shunt in erythrocytes. *J Clin Invest* 58:797-802 [Experimental study in human erythrocytes in vitro].
- *Ministry of Health, Mozambique. 1984. Mantakassa: An epidemic of spastic paraparesis associated with chronic cyanide intoxication in a cassava staple area of Mozambique 1. Epidemiology and clinical and laboratory findings in patients; 2. Nutritional factors and hydrocyanic acid content of casava products. *Bull World Health Organ* 62:477-484, 485-492 [Chronic cyanide toxicity].
- *Mirvish SS, Salmasi S, Cohen SM, Patil K, Mahboubi E. 1983. Liver and forestomach tumors and other forestomach lesions in rats treated with morpholine and sodium nitrite, with and without sodium ascorbate. *J Natl Cancer Inst* 71:81-85 [Experimental study].
- *Mladoveanu C, Gheorghiu P. 1929. Le nitrite de soude comme antidote de l'empoisonnement experimental par le cyanure de potassium. *C R Soc Biol (Paris)*, 102:164-166 [Experimental study].
- *Mokhlesi B, Leikin JB, Murray P, Corbridge TC. 2003. Adult toxicology in critical care, part II: Specific poisonings. *Chest* 123:897-922.
- *Montgomery R, Reinhart CF, Terril JB. 1975. Comments on fire toxicity. *Comb Toxicol* 2:179-212 [Experimental study on HCN generated by pyrolysis of different materials].

- *Morikawa T, Yanai E, Nishina T. 1987. Toxicity evaluation of fire effluent gases from experimental fires in a building. *J Fire Sciences* 5:248-271.
- *Moriya F, Hashimoto Y. 2001. Potential for error when assessing blood cyanide concentrations in fire victims. *J Forensic Sci*, 46:1421-1425.
- *Morocco AP. 2005. Cyanides. *Crit Care Clin* 21::691-705, vi.
- *Morris B. 1987. Postmortem features and criteria for the diagnosis of acute lethal cyanide poisoning. In Ballantyne B, Marrs TC eds. *Clinical and experimental toxicology of cyanides*. John Wright, Bristol, UK, 217-247 [Forensic study].
- *Morse DL, Harrington JM, Heath CW. 1976. Laetrile, apricot pits, and cyanide poisoning. *N Engl J Med* 295:1264.
- *Moseley RH, Jarose S, Permod P. 1992. Hepatic Na(+)-dicarboxylate cotransport: Identification, characterization, and acinar localization. *Am J Physiol* 263:G871-879 [Detailed study on perfusion of α -ketoglutarate in isolated rat liver].
- *Mouzas GL, Smith RL, Syed A. 1983. A scare of cyanide poisoning. *Br J Clin Pract* 37:245-248.
- *Mulford CE. 1966. Solvent extraction techniques for atomic absorption spectroscopy. *At Absorpt Newsl* 5:88-90 [Analytical study].
- *Myers RA. 1980. Treatment protocol for smoke inhalation victims. *Occup Health Saf* 49:56-57, 59-60.
- *Nagaratnam N, Alagaratnam K, Thambapillai AJ, Wijemanne HS. 1974. Acute renal failure following potassium ferrocyanide poisoning treated with peritoneal dialysis. *Forensic Sci* 14:87-89 [Other poisoning].
- *Nederlandse Pharmacopee. 1958. 6th ed. Staatsdrukkerij, The Hague, Netherlands.
- *Newton GL, Dorian R, Fahey RC. 1981. Analysis of biological thiols: Derivatization with monobromobimane and separation by reverse-phase high-performance liquid chromatography. *Anal Biochem* 114:383-387 [Analytical study of thiols at the picomole level].
- *NFPA. 1978. Fire protection guide on hazardous materials, 7th ed. National Fire Protection Association, Boston, Massachusetts, USA.
- *Ng LL, Naik RB, Polak A. 1982. Paraquat ingestion with methemoglobinemia treated with methylene blue. *Br J Med* 284:1445-1446 [Other poisoning].

- *Noeller TP. 2001. Biological and chemical terrorism: recognition and management. *Cleve Clin J Med* 68:1001-10016.
- *Nogué-Xarau S, Dueñas A, Burillo G. 2004. Acute chemical emergencies. *N Engl J Med* 350:2102-2104 [Comment on Kales and Christiani, 2004a; author reply 2103-2104].
- *Olsen NS, Klein RJ. 1947. Effect of cyanide on the concentration of lactate and phosphates in brain. *J Biol Chem* 167:739-746 [Experimental study].
- *Oluwole OS, Onabolu AO, Cotgreave IA, Rosling H, Persson A, Link H. 2003. Incidence of endemic ataxic polyneuropathy and its relation to exposure to cyanide in a Nigerian community. *J Neurol Neurosurg Psychiatry* 74:1417-1422.
- *Oluwole OS, Onabolu AO, Cotgreave IA, Rosling H, Persson A, Link H. 2002. Low prevalence of ataxic polyneuropathy in a community with high exposure to cyanide from cassava foods. *J Neurol*. 249:1034-1040.
- *Oluwole OS, Onabolu AO, Cotgreave IA, Rosling H, Persson A, Link H. 2003. Incidence of endemic ataxic polyneuropathy and its relation to exposure to cyanide in a Nigerian community. *J Neurol Neurosurg Psychiatry* 74:1417-1422.
- *Oluwole OS, Onabolu AO. 2004. High mortality of subjects with endemic ataxic polyneuropathy in Nigeria. *Acta Neurol Scand* 110:94-99.
- *Orellano T, Dergal E, Alijani M, Briggs C, Vasquez J, Goldbaum LR, Absolon KB. 1976. Studies on the mechanism of carbon monoxide toxicity. *J Surg Res* 20:485-487.
- *Osuntokun B.O. 1976. Nutritional problems in the African region. *Bull Schweiz Akad Med Wiss* 31:252-376.
- *Osuntokun BO, Osuntokun O. 1971. Tropical amblyopia in Nigerians. *Am J Ophthalmol* 72:708-716.
- *Osuntokun BO. 1981. Cassava diet, chronic cyanide intoxication and neuropathy in the Nigerian Africans. *World Rev Nutr Diet* 36:141-73.
- *Owasoyo JO, Iramain CA. 1980. Acetylcholinesterase activity in rat brain: Effect of acute cyanide intoxication. *Toxicol Lett* 6:1-3.
- *Padberg LR, Martin T. 1939. Three fatal cases of sodium nitrite poisoning. *J Am Med Assoc* 113:1733 [Other poisoning].

- *Padwell A. 1997. Cyanide poisoning: Case studies of one homicide and two suicides. *Am J Forensic Med Pathol* 18:185-188.
- *Pahl MV Vaziri ND. 1982. In vivo and in vitro hemodialysis studies of thiocyanate. *J Toxicol Clin Toxicol* 19:965-974 [Excretion of thiocyanate].
- *Pane GA, Mohler SR, Hamilton GC. 1985. The Cincinnati DC-9 experience: Lessons in aircraft and airport safety. *Aviat Space Environ Med* 56:457-461.
- *Patz A. 1968. Comment on oxygen therapy. *Arch Ophthalmol* 79:507 [Other poisoning].
- *Patz A. 1968. New role of ophthalmologist in prevention of retrolental fibroplasia. *Trans Am Ophthalmol Soc* 66:940-985 [Other poisoning].
- *Patzsch H. 1953. Gesundheitsstörungen nach dem Genuß von weissen Bohnen. *Medizinische* 43:1390-1391.
- *Paulet G, Menoret P. 1954. Etude comparée de la sensibilité du sang de l'homme (frais ou conservé) et du sang de chien à divers méthémoglobinisants. *C R Séances Soc Biol Fil* 148:1014-1017 [Experimental study].
- *Paulet G, Chary R, Bocquet P, Fouilhoux M. 1960. Valeur comparée du nitrite de sodium et des chélates de cobalt dans le traitement de l'intoxication cyanhydrique chez l'animal (chien-lapin) non-anesthésié. *Arch Int Pharmacodyn* 127:104-117 [Experimental study].
- *Paulet G, Desnos J, Battig J. 1966. De la toxicité de l'acrylonitrile. *Arch Mal Prof* 27:849-856.
- *Pearce LL, Bominaar EL, Hill BC, Peterson J. 2003. Reversal of cyanide inhibition of cytochrome c oxidase by the auxiliary substrate nitric oxide: An endogenous antidote to cyanide poisoning? *Biol Chem* 278:52139-52145.
- *Peleg R, Goldberger Y, Bursztejn de Myttemaere S, Heifetz M. 1979. [Sodium nitroprusside poisoning]. *Harefuah* 97:122-123 [Hebrew].
- *Pelton WJ. 1950. The effect of zinc chloride and potassium ferrocyanide as a caries prophylaxis. *J Dent Res* 29:756-759.
- *Perez A, Johnson-Arbor K, McKay CA. 2006. Carbon monoxide poisoning, myocardial injury, and mortality. *JAMA* 295:2601 [Comment on Henry et al, 2006].
- *Persson B, Simonson M. 1998. Fire emissions into the atmosphere. *Fire Technology* 34:266-279.

- *Pettersen JC, Cohen SD. 1985. Antagonism of cyanide poisoning by chlorpromazine and sodium thiosulfate. *Toxicol Appl Pharmacol* 81:265-273.
- *Pettersen JC, Cohen SD. 1993. The effects of cyanide on brain mitochondrial cytochrome oxidase and respiratory activities. *J Appl Toxicol* 13(1):9-14.
- *Pharmacopée Française. 1988. Hydroxocobalamine. In 10th ed Commission nationale de la Pharmacopée française, Agence Française de Sécurité Sanitaire des Produits de Santé, Saint-Denis, France. Adrapharm, Paris, France.
- *Pharmacopoea Nordica. 1964. Editio Svecica. Apotekarsocietetens förlag, Stockholm, Sweden.
- *Philip JM. 1970. Toxicity and naturally occurring chemicals in food. *R Soc Health J* 90:237-242.
- *Piantadosi CA, Sylvia AL, Jöbsis FF. 1983. Cyanide-induced cytochrome a,a3 oxidation-reduction responses in rat brain *in vivo*. *J Clin Invest* 72:1224-1233 [Experimental study].
- *Pirie NW. 1986. Cassava toxicity. *Lancet* 1:561.
- *Pitchumoni CS, Jain NK, Lowenfels AB, DiMagno EP. 1988. Chronic cyanide poisoning: unifying concept for alcoholic and tropical pancreatitis. *Pancreas* 3:220-222.
- *PoisIndex. 1987. Microfiche data base, 53rd ed. Micromedex, Denver, Colorado, USA.
- *Polson CJ Tattersall RN. 1969. Clinical toxicology. London, Pitman, pp 128-159 [HCN used as fumigant]..
- *Pritchard JD. 2007. HPA compendium of chemical hazards: Hydrogen cyanide. Version 2. Health Protection Agency, London, UK [Review]. www.hpa.org.uk
- *Prockop LD, Chichkova RI. 2007. Carbon monoxide intoxication: An updated review. *J Neurol Sci* 262:122-130.
- *Product Information. 1984. Product Information for the cyanide antidote package. Eli Lilly, Indianapolis, Indiana, USA.
- *Pryor AJ, Johnson DE, Jackson NN. 1975. Hazards of smoke and toxic gases produced in urban fires. *Combust Toxicol* 2:64-112 [Cited by US-NAC, 2000a].
- *Puls R, Newschwander FP, Greenway JA. 1978. Case report: Cyanide poisoning from *Glyceria grandis* S. Wats. ex Gray (tall mannagrass) in British Columbia beef herd. *Can Vet J* 19:264-265.

- *Purser D. 1996. Behavioural impairment in smoke environments. *Toxicology* 115:25-40.
- *Purser DA, Grimshaw P, Berrill KR. 1984. Intoxication by cyanide in fires: A study in monkeys using polyacrylonitrile. *Arch Environ Health* 39:394-400.
- *Purser DA, Woolley WD. 1983. Biological studies of combustion atmospheres. *J Fire Sci* 1:118-144.
- *Purser DA. 1984. A bioassay model for testing incapacitating effects of exposure to combustion product atmospheres using *Cynomolgus* monkeys. *J Fire Sci* 2:20-36.
- *Radonjić V, Bosković B, Pribić A. 1983. Hitna pomoć kod trovanja cijanovodonikom prilikom fumigacije [Emergency treatment of hydrogen cyanide poisoning during fumigation]. *Arh Hig Rada Toksikol* 34:239-249 [Croatian].
- *Rammer L, Rosling H, Sörbo B, Tibbling L, Tydén H. 1988. Cyanidförgiftning — en försummad risk [Cyanide poisoning — an overlooked risk]. *Läkartidningen* 85:853-854, 856 [Swedish].
- *Ravishankar DK, Kashi SH, Lam FT. 1998. Organ transplantation from donor who died of cyanide poisoning: A case report. *Clin Transplant* 12:142-143.
- *Roberts RM. 1981. Carbon monoxide poisoning and fire-associated deaths. *Lancet* 1981:1816.
- *Román GC, Spencer PS, Schoenberg BS. 1985. Tropical myeloneuropathies: The hidden endemias. *Neurology* 35:1158-1170.
- *Rose CL, Worth RM, Chen KK. 1965. Hydroxo-cobalamin and acute cyanide poisoning in dogs. *Life Sci* 4:1785-1789 [Experimental study].
- *Roseman E, Goodman CW, McCulloch WS. 1946. Rapid changes in cerebral oxygen tension induced by altering the oxygenation and circulation of the blood. *J Neurophysiol* 9:33-40 [Experimental study].
- *Rosling H. 1989. Cassava associated neurotoxicity in Africa. In Volans GN, Sims J, Sullivan FM, Turner P ed, *Proceedings of the 5th International Congress of Toxicology*. Taylor Francis, Brighton, UK, pp 605-614 [Chronic toxicity].
- *Rudomin P, Rubio R. 1959. Actividad subliminal de los quimiorreflejos originados por el cianuro de potasio [Subliminal activity of the chemoreflexes originated by potassium cyanide]. *Acta Physiol Lat Am* 9:194-208.

*Rumack BH, ed. 1987. Cyanide poisoning management. In Poisindex: Poison information system. Micromedex, Denver, Colorado, USA, p 52:1-F4.

*Russel WO, Stahl WC. 1942. Fatal poisoning from potassium thiocyanate treatment of hypertension. J Am Med Assoc 119:1177-1181 [Other poisoning].

*Salomone J, Sohn AP, Ritzlin R, Gauthier JH, McCarty V. 1987. Correlations of injury, toxicology, and cause of death to Galaxy flight 203 crash site. J Forensic Sci 32:1403-1415.

*Samele Acquaviva E. 1968. Intossicazione acuta da caduta accidentale in un bagno per cadmiatura [Acute poisoning caused by an accidental fall into a cadmium-plating bath]. Minerva Anestesiol 34:1171 [Italian].

*Sandberg CG. 1967. A case of chronic poisoning with potassium cyanide? Acta Med Scand 181:233-236.

*Sanghavi P, Bhalla K, Das V. 2009. Fire-related deaths in India in 2001: a retrospective analysis of data. Lancet 373:1282-1288.

*Sass MD, Caruso CJ, Axelrod DR. 1969. Mechanism of the TPNH-linked reduction of methemoglobin by methylene blue. Clin Chem Acta 24:77 [Mechanism of action against methemoglobinaemia and side effects].

*Savateev NV, Tonkopii VD, Frolov SF. 1969. Vliianie kisloroda pri normal'nom i povyshennom dovlenii na techenie otravlenii tsianidami [Effect of oxygen at normal and increased pressure on the course of cyanide poisoning]. Farmakol Toksikol 32:328-330 [Russian].

*Savolainen H, Kirchner N. 1998. Toxicological mechanisms of fire smoke. Internet Journal of Rescue and Disaster Medicine 1:1-7. www.ispub.com

*Schardein JL. 1985. Chemically induced birth defects. Marcel Dekker, New York, USA, pp 74, 759, 810-824 [Other poisoning].

*Scheele CW. 1786. Chemical essays. Beddves T (trans) Reprint 1966, London, Dawson [Early report on isolation of HCN from Prussian blue].

*Schoepfle GM, Bloom FE. 1959. Effects of cyanide and dinitrophenol on membrane properties of single nerve fibers. Am J Physiol 197:1131-1135.

*Schrader JCC. 1802. Des expériences de M. Schrader sur l'acide prussique continu dans les plantes. J Phys Chim Hist Nat Arts 56:224-226 [Early report on cyanide obtained from bitter almonds].

- *Schuckmann F. 1994. Successful treatment of life-threatening propionitrile exposure with sodium nitrite/sodium thiosulfate followed by hyperbaric oxygen, by Scolnick et al. *J Occup Med* 36:1283.
- *Schumacher G. 1959. [Contribution to the treatment of cough]. *Ther Ggw* 98:581-582 [German].
- *Scott A. 1950. A cytological analysis of the effects of cyanide and 4,6-dinitro-orthocresol on the mitotic phases in *Arbacia punctulata*. *Biol Bull* 99:362-363.
- *Serra C, Lucarelli S. 1951. Un anno di terapia dinitrilica [One year of the treatment with dinitrile]. *Gazz Med Ital* 110:324-326 [Italian].
- *Sharpe AT. 1976. The chemistry of cyano complexes of the transition metals. Academic Press, New York, USA [Analytical study].
- *Sheehy MH, Way JL. 1974. Nitrite intoxication: Protection with methylene blue oxygen. *Toxicol Appl Pharmacol* 30:221-226 [Other poisoning].
- *Shimanko II, Babichev MN. 1968. [A case of severe mercury cyanide poisoning]. *Ter Arkh* 40:117-119 [Russian].
- *Shiono H, Maseda C, Akane A, Matsubara K. 1991. Rapid and sensitive quantitation of cyanide in blood and its application to fire victims. *Am J Forens Med Pathol*, 12:50-53.
- *Shusterman D, Alexeeff G, Hargis C, Kaplan J, Sato R, Gelb A, Becker C, Benowitz N, Gillen M, Thollaug S, Balmes J. 1996. Predictors of carbon monoxide and hydrogen cyanide exposure in smoke inhalation patients. *J Toxicol Clin Toxicol* 34:61-71.
- *Sigrist T, Dirnhofer R. 1979. Zur Entstehung der kombinierten inhalatorischen Blausaure-Kohlenmonoxid-Vergiftung. *Arch Kriminol* 163:145-159.
- *Silcock A, Robinson D, Savage NP. 1978. Fire in dwellings - an investigation of actual fires. Current Paper 80/78, Building Research Establishment, Fire Research Station, Borehamwood, U.K.
- *Silverman SH, Purdue GF, Hunt JL, Bost RO. 1988. Cyanide toxicity in burned patients. *J Trauma* 28:171-176 [with commentary *J Trauma* 6:555].
- *Silverman SH, Purdue GF, Hunt JL, Bost RO. 1988. Cyanide toxicity in burned patients. *Burns* 14:338-340 [Abstract].

- *Singer MM, Wright F, Stanley LK, Roc BB, Hamilton WK. 1970. Oxygen toxicity in man. A prospective study in patients after open heart surgery. *New Engl J Med* 283:1473 [Other poisoning].
- *Smith RP, Kruszyna H. 1976. Letter: Nitroprusside and cyanide. *Br J Anaesth* 48:396.
- *Smith RP, Kruszyna H. 1976. Toxicology of some inorganic antihypertensive anions. *Fed Proc* 35:69-72.
- *Smith RP, Olson MV. 1973. Drug-induced methemoglobinemia. *Semin Hematol* 10:253-268 [Other poisoning].
- *Smith RP. 1971. Spectrophotometric determination of methemoglobin and oxyhemoglobin in the presence of methylene blue. *Clin Toxicol* 4:273 [Analytical study].
- *Soars MH, Hendlin D. 1951. The use of potassium cyanide in the *Lactobacillus leichmannii* assay for vitamin B12. *J Bacteriol* 62:15-17.
- *Sörbo B, Ohman S. 1978. Determination of thiosulphate in urine. *Scand J Clin Lab Invest* 38:521-527 [Analytical study].
- *Spahr RC, Salisburey DJ, Krissberg A, Prin W. 1980. Intra-amniotic injection of methylene blue leading to methemoglobinemia in one of twins. *Int J Gynaecol Obstet* 17:477-478 [Other poisoning].
- *Speitel LC. 1996. Fractional effective dose model for post-crash aircraft survivability. *Toxicology* 115:167-77.
- *Spillane JD. 1969. Tropical neurology. *Proc R Soc Med* 62:403-410.
- *Standefor JC, Jones AM, Street E, Inserra R. 1979. Death associated with nitrite ingestion: report of a case. *J Forensic Sci* 24:768-771 [Other poisoning].
- *Stemler FW, Groff WA, Kaminskis A, Johnson RP, Froehlich HL. 1995. Comparison of the treatment of cyanide poisoning in the *Cynomolgus* monkey with sodium nitrite or 4-dimethylaminophenol (4-DMAP), with and without sodium thiosulfate. *Govt Rep Announc Index, Issue 2*. [Experimental study].
- *Stemler FW, Kaminskis A, Tezak-Reid TM, Stotts RR, Moran TS, Hurt HH, Ahle NW. 1995. Correlation of atmospheric and inhaled blood cyanide levels in miniature pigs. In Nelson GL, ed. *Fire and Polymers II: Materials and tests for hazard prevention*, chapter 12. *ACS Symp Ser* 599:312-322.

- *Stempfel. 1951. Sur la toxicité des cyanures alcalins utilisés dans l'industrie, notamment dans les ateliers de traitements thermiques. *Med Usine Rev Hyg Ind Mal Prof* 13:220-223.
- *Stevenson RN, Kingswood C, Cohen SL, Vesey CJ. 1988. Cyanide and fire victims. *Lancet* 2:1145.
- *Stimpfl T, Grassberger M, Andresen H. 2007. Zyanidvergiftung. *Rechtsmedizin* 17:343-350.
- *Stojicic N, Baumstark-Khan C, Hellweg CE, Grotheer HH, Reitz G, Kolanus W, Hemmersbach R. 2010. Toxicity of ethylene combustion condensates is directly proportional to their carbon content. *Toxicology* 269:35-40.
- *Stolarsky F, Haley TJ. 1951. Acute and chronic toxicity of toluidine blue and neutral red. *Fed Proc* 10:337 [Other poisoning].
- *Streicher E. 1951. Toxicity of colchicine, di-isopropyl fluorophosphate, intocostin, and potassium cyanide in mice at 4 degrees C. *Proc Soc Exp Biol Med* 76:536-538.
- *Strother DE. 2004. Overexposure to acrylonitrile: Review of mechanism of acute toxicity and antidotes. Unpublished report, draft. The AN Group Health Sciences Committee, Washington, DC, USA [Review].
- *Suarez KA, Bhonsle P. 1978. Enhanced hepatotoxicity of carbon tetrachloride following sodium nitrite pretreatment. *Arch Int Pharmacodyn Ther* 234:329-334 [Other poisoning].
- *Suzuki T, Saito K, Kobayasi Y. 1951. Adrenaline secretion of the suprarenal glands in cyanide anoxia. *Tohoku J Exp Med* 54:343-350.
- *Sykes AH. 1981. Early studies on the toxicology of cyanide. In: Vennesland B, Conn EE, Knowles CJ, Westley J, Wissing F ed. *Cyanide in biology*. New York, London, Academic Press 1-9 [Other poisoning].
- *Szirmai E. 1959. [Pharmacodynamic studies on a new nitrile]. *Z Gesamte Inn Med* 14:1131-1133 [German].
- *Takahama U, Oniki T, Hirota S. 2002. Oxidation of quercetin by salivary components. Quercetin-dependant reduction of salivary nitrite under acidic conditions producing nitric oxide. *J Agric Food Chem* 50, 4317 - 4322 [Cited by OECD, 2005].
- *Takahashi H, Matsuzaki H, Kumei K, Takahashi H. 1959. EEG-arousal by gamma-aminobutyric acid (GABA) in the rabbit under anesthesia, cyanpoisoning or asphyxia. *Jpn J Physiol* 9:207-217.

- *Tauberger G, Karzel K, Roezel V. 1974. Versuche zur Cyanidentgiftung mit Blut-Austauschtransfusionen. *Arch Toxicol* 32:189-197.
- *Tauberger G, Klimmer OR. 1963. Tierexperimentelle Untersuchungen einiger Kobaltverbindungen nach intravenöser Injektion. *Arch Int Pharmacodyn Ther* 143:219-239 [Experimental study].
- *Ten Brink WA, Wiezer JH, Luijpen AF, van Heijst AN, Pikaar SA, Seldenrijk R. 1982. Nitrite poisoning caused by food contaminated with cooling fluid. *J Toxicol Clin Toxicol* 19:139-147 [Other poisoning].
- *Terrill JB, Montgomery RR, Reinhardt CF. 1978. Toxic gases from fires. *Science* 200:1343-1347 [Screening test for toxic fire gases].
- *Tewe OO, Maner JH. 1980. Cyanide, protein and iodine interactions in the performance, metabolism and pathology of pigs. *Res Vet Sci* 29:271-276.
- *Tewe OO, Maner JH. 1981. Long-term and carry-over effect of dietary inorganic cyanide (potassium cyanide) in the life cycle performance and metabolism of rats. *Toxicol Appl Pharmacol* 58:1-7.
- *Tewe OO, Maner JH. 1981. Performance and pathophysiological changes in pregnant pigs fed cassava diets containing different levels of cyanide. *Res Vet Sci* 30:147-151.
- *Tewe OO, Maner JH. 1982. Cyanide, protein and iodine interactions in the performance and metabolism of rats. *J Environ Pathol Toxicol Oncol* 6:69-77.
- *Tewe OO, Pessu E. 1982. Performance and nutrient utilization in growing pigs fed cassava peel rations containing different cyanide levels. *Nutr Rep Int* 26:51-58.
- *Thiers RE, Williams JF, Yoe JH. 1955. Separation and determination of millimicrogram amounts of cobalt. *Anal Chem* 27:1725-1731 [Analytical study].
- *Tolbert BM, Hughes AM. 1959. C¹⁴-Labeled cyanide: Radioactivity excretion in mice and estimation of radiation dose to humans. *Metabolism* 8:73-78.
- *Tölle R, von Kracht E. 1969. Cerebrale Folgeerscheinungen einer Bromcyan-Vergiftung. *Nervenarzt* 40:594-595.
- *Tomoda A, Yoneyama Y. 1979. On the relationship of hemoglobin oxidation with the conformation of hemoglobin. *Experientia* 35:15-16 [Analytical study].

- *Tompsett SL. 1959. A note on the determination and identification of cyanide in biologic materials. *Clin Chem* 5:587-591.
- *Treitman RD, Burgess WA, Gold A. 1980. Air contaminants encountered by firefighters. *Am Ind Hyg Assoc J* 41:796-802.
- *Trimble EJ. 1996. The management of aircraft passenger survival in fire. *Toxicology* 115:41-61.
- *Tsou CL. 1951. On the cyanide inactivation of succinic dehydrogenase and the relation of succinic dehydrogenase to cytochrome b. *Biochem J* 48:lii-liii.
- *Tung A, Lynch J, McDade WA, Moss J. 1997. A new biological assay for measuring cyanide in blood. *Anesth Analg* 85:1045-1051.
- *Turrina S, Neri C, De Leo D. 2004. Case review: Effect of combined exposure to carbon monoxide and cyanides in selected forensic cases. *J Clin Forensic Med* 11:264-267.
- *Tylleskär T, Banea M, Bikangi N, Fresco L, Persson LA, Rosling H. 1991. Epidemiological evidence from Zaire for a dietary etiology of konzo, an upper motor neuron disease. *Bulletin WHO* 69:581-589.
- *Tylleskär T, Banea M, Bikangi N, Nahimana G, Persson LA, Rosling H. 1995. Dietary determinants of a non-progressive spastic paraparesis (Konzo): a case-referent study in a high incidence area of Zaire. *Int J Epidemiol* 24:949-956.
- *Uhl W, Nolting A, Gallemann D, Hecht S, Kovar A. 2008. Changes in blood pressure after administration of hydroxocobalamin: relationship to changes in plasma cobalamins-(III) concentrations in healthy volunteers. *Clin Toxicol (Phila)* 46:551-559; discussion 576-577.
- *United States Dispensatory. 1973. Dispensatory of the United States of America, 27th ed. JB Lippincott, Philadelphia, Pennsylvania, USA.
- *United States Environmental Protection Agency. 1994. Acrylonitrile fact sheet: support document (CAS No. 107-13-1). OPPT Chemical Fact Sheets, EPA 749-F-95-011a.
- *US-NAC. 2000a. Hydrogen cyanide, proposed acute exposure guideline levels (AEGLs). Unpublished report proposed 1:1/2000. National Advisory Committee to develop AEGLs. US Environmental Protection Agency, Washington DC, USA [Review].
- *US-NAC. 2000b. Acute exposure guideline levels (AEGLs) for acetone cyanohydrin, draft June 2000. Unpublished report. FoBiG, Forschungs- und Beratungsinstitut Gefahrstoffe GmbH.

Griem P. National Advisory Committee to develop AEGLs. US Environmental Protection Agency, Washington DC, USA [Review].

*Uwakwe AA, Monanu MO, Anosike EO. 1991. Whole blood cyanide levels of mainly dietary origin in a human population sample in Port Harcourt, Nigeria. *Plant Foods Hum Nutr* 41:117-124.

*Vale A. 2007. Cyanide. *Medicine* 35:11.

*van Heijst AN, Maes RA, Mtanda AT, Chuwa LM, Rwiza HT, Moshi NH. 1994. Chronic cyanide poisoning in relation to blindness and tropical neuropathy. *J Toxicol Clin Toxicol* 32:549-556.

*Vesey CJ, Cole PV. 1978. Cyanide toxicity following nitroprusside. *Can Anaesth Soc J* 25:437-438 [Letter, with letter by Smith F, West D, 1978].

*Vieira Filho JPB, Oliveira ASB, da Silva MRD, Amaral AL, Schultz RR. 1997. Polineuropatia nutricional entre índios Xavantes [Polyneuropathy deficiency among Xavante indians]. *Rev Assoc Med Bras* 43:82-88 [Portuguese].

*Vincer MJ, Allen AC, Evans JR, Nwaesei C, Stinson DA. 1987. Methylene-blue-induced hemolytic anemia in a neonate. *Can Med Assoc J* 136:503-504 [Other poisoning].

*von Lüpke H, Schmidt K. 1977. Kombinierte Kohlen-Monoxid/Cyanwassertoff Vergiftung nach Verschmoren von Kunststoff. *Z Rechtsmed* 79:69-71.

*Way JL, Gibbon SL, Sheehy M. 1966. Cyanide intoxication: protection with oxygen. *Science* 210:211 [Experimental study]

*Way JL, Sylvester D, Morgan RL, Isom GE, Burrows CB, Tamulinas CB, Way JL.. 1984. Recent perspectives on the toxicodynamic basis of cyanide antagonism. *Fundam Appl Toxicol* 4S:231-239 [Experimental study].

*Way JL, Tamulinas CB, Leung P, Ray L, Nizamani S, Sylvester D, Way JL, Chiou F. 1984. Pharmacologic and toxicologic basis of cyanide`antagonism. *Proc West Pharmacol Soc* 27:149-153 [Experimental study].

*Weisburger JH, Horn CL, Barnes WS. 1983. Possible genotoxic carcinogens in foods in relation to cancer causation. *Semin Oncol* 10:330-341 [Other poisoning].

*Wetherell HR. 1966. The occurrence of cyanide in the blood of fire victims. *J Forensic Sci* 11:167-173.

- *Whitman JG, Taylor AR, White JM. 1979. Potential hazard of methylene blue. *Anaesthesia* 34:181-182 [Other poisoning].
- *WHO. 1996. Guidelines for drinking water quality. 2nd ed. Vol 2: Health criteria and other supporting information. Geneva, Switzerland, 226-230.
- *WHO. 2003. Cyanide in drinking-water, background document for development of WHO guidelines for drinking-water quality. Report WHO/SDE/WSH/03.04/05. World Health Organization, Geneva, Switzerland.
- *WHO. 2004. Cyanide, chemical fact sheet 12.33. In Guidelines for drinking-water quality, 3rd ed. Volume 1, Recommendations. World Health Organization, Geneva, Switzerland, 339-340. ISBN 92 4 154638 7
- *Whysner J, Ross PM, Conaway CC, Verna LK, Williams GM. 1998. Evaluation of possible genotoxic mechanisms for acrylonitrile tumorigenicity. *Regul Toxicol Pharmacol* 27:217-239 [Review].
- *Wikipedia. 2009. Amygdalin. In Wikipedia, the free encyclopedia. [Review]. <http://en.wikipedia.org/wiki/Amygdalin>.
- *Wilkins RW, Haynes FW, Weiss S. 1937. The role of the venous system in circulatory collapse induced by sodium nitrite. *J Clin Invest* 16:85-91 [Other poisoning].
- *Wirth W. 1968. Schädigungsmöglichkeiten durch Antidote. *Arch Toxicol* 24:71-82
- *Wokes F, Willimott SG. 1951. The determination of cyanide in seeds. *J Pharm Pharmacol* 3:905-917.
- *Yeoh M.J, Braitberg G. 2004. Carbon monoxide and cyanide poisoning in fire related deaths in Victoria, Australia. *J Toxicol Clin Toxicol* Vol. 42, No. 6, pp. 855-863.
- *Yoshida M, Adachi J, Watabiki T, Tatsuno Y, Ishida N. 1991. A study on house fire victims: Age, carboxyhemoglobin, hydrogen cyanide and hemolysis. *Forensic Sci Int* 52:13-20.
- *Zamecnik J, Tam J. 1987. Cyanide in blood by gas chromatography with NP detector and acetonitrile as internal standard: Application on air accident fire victims. *J Anal Toxicol* 11:47-48.
- *Zikria BA, Weston GC, Chodoff M, Ferrer JM. 1972. Smoke and carbon monoxide poisoning in fire victims. *J Trauma* 12:641-645.

APPENDIX A: QUESTIONNAIRE

A.1 Personal Cases

In order to have information on the occurrence of cyanide and nitrile poisoning a structured questionnaire was mailed to the EAPCCT members and to the industry.

The aim of this survey was to assess the number of poisonings reported to Poison Centres / hospital (intensive care unit) / Industry and to summarise their experience (Part A) and when available to obtain more detailed information on exposure, substances involved, antidotal treatment used and outcome (Part B).

The questionnaire was distributed and responses were received through personal contacts of the TF members and the EAPCCT secretariat, with information from the Cyanide Forum and WHO/IPCS.

The return on questionnaire Part A was limited to 20 respondents from 14 European countries; 2 respondents (from Estonia and the Netherlands) stated that no cyanide poisoning was reported and their reply is not included in the overview below.

Overview of responses to questionnaire

Country	Centre	Cases of poisoning	Number	Period (years)	Antidote(s) used	Reference
Belgium	PCC	Cyanide	15	2004 - 2008	Hydroxocobalamine	Smet, 2009
Belgium	PCC	Nitrile	2	2004 - 2008	-	Smet, 2009
Czech Republic	Hospital	Cyanide	3	1990 - 1996	AN, SN STS	Pelclová, 2009
Denmark	PCC	Cyanide	2	Since 2006	Hydroxocobalamine	Ebbenhøj, 2009
Denmark	PCC	Nitrile	0	Since 2006	-	Ebbenhøj, 2009
Finland	PCC	Cyanide	5 deaths	1998 - 2001	-	Hoppu, 2009
France	Hospital	Cyanide	12		STS HOCO, Co-EDTA	Baud, 2009
France	Hospital	Nitrile	2		Hydroxocobalamine	Baud, 2009
Germany	Hospital	Cyanide	9	Since 1985	4-DMAP, STS	Zilker, 2007
Germany	Industry	Cyanide	58	Since 1977	4-DMAP, STS, NAC, HOCO AN, SN Oxygen	Steffens, 2008 Rödelsperger, 2009 Küpper, 2009 Busch, 2010

Country	Centre	Cases of poisoning	Number	Period (years)	Antidote(s) used	Reference
Germany	Industry	Nitrile	116	Since 1977	4-DMAP, STS ,NAC Oxygen	Steffens, 2008 Rödelsperger, 2009 Küpper, 2009 Busch, 2010
Germany	PCC (Mainz)	Cyanide	88	1995 - 2009	DMAP, STS HOCO	Sauer, 2009
Germany	PCC (Mainz)	Nitrile	20	1995 - 2009	DMAP,STS, HOCO	Sauer, 2009
Germany	PCC (Mainz)	Cyanogenic glycoside	98	1995 - 2009	DMAP, STS HOCO	Sauer, 2009
Ireland	Hospital	Cyanide	4	Since 2004	-	Casey, 2009
Ireland	Hospital	Nitrile	3	Since 2004	-	Casey, 2009
Italy	PCC (Firenze)	Cyanide	4	Since 2000	SN, STS, HOCO, dimercaprol	Botti, 2009
Norway	Hospital	Cyanide	5 - 10 cases/y	Since 2004	Hydroxocobalamine	Jacobson, 2009
Norway	Hospital	Nitrile	0	Since 2004	-	Jacobson, 2009
Slovak Republic	PCC	Cyanide	1	9 y	DMAP, STS	Ficekova, 2009
Spain	PCC	Cyanide or nitrile, incl 5 cyanogenic glycoside and 4 fire smoke	27	2006 - 2008		Martinez Arrieta, 2009
UK	PCC (NPIS Edinburgh)	Cyanide	3	Since 1987	None	Whyte, 2009
UK	PCC (NPIS Edinburgh)	Nitrile	0	Since 1987		Whyte, 2009

In the global overview 482 cases of cyanide / nitrile / cyanogenic compound poisonings were reported of which 217 cases came with further details and are listed in an extended table in annex.

A wide range of antidotes was used. In spite of the availability of antidotes cyanide poisoning is a severe and uncommon disease and the problem remains in a selection of appropriate antidote based on efficacy and safety depending on the condition of poisoning.

A.2 The questionnaire

EUROPEAN CENTRE FOR ECOTOXICOLOGY AND TOXICOLOGY OF CHEMICALS

To: EAPCCT members

C054

Brussels, 13th December 2009

Dear Colleague,

Questionnaire on cyanide and nitrile poisoning – Further responses?

Following our questionnaire on cyanide and nitrile poisoning, announced on the EAPCCT website last April / May, we have received several replies from national Poison Centres, and from the cyanide and nitrile industries (list of responses enclosed [Note: By August 2010, there were 20 respondents in all, see list attached]). The statistics and information given by respondents so far are very useful for our Task Force in compiling a critical review of the efficacy and safety of cyanides antidotes (terms of reference enclosed [Section 2.1 of report]). A report is expected to be published early in 2011. It will also include a systematic evaluation, using the Poison Severity Score, of more than 300 cases to establish a cyanides toxidrome and assess the efficacy of various antidotes.

For these reasons, it seems prudent to repeat the questionnaire and we are soliciting PCCs and ICUs that have not been able to reply so far. As before, respondents are kindly invited to answer question A below to summarise their experience, and complete the attached form B for each poisoning case with the available information. All replies may be sent directly to the undersigned by e-mail: henk.vrijhof@ecetoc.org. All contributions will be acknowledged in the final publication.

A. Cases of poisoning	Number	Period (years)	Antidote(s) used
Cyanide?			
Nitrile?			

We look forward to receiving a positive reply.

Yours sincerely,

Ir. H. Vrijhof

Chemicals Programme Manager

encl. Form **B**, ToR, TF

B. ECETOC questionnaire on poisoning cases with cyanides

1	Contact details				
1.1	Centre:				
1.2	Contact person:				
1.3	Phone:				
1.4	E-mail:				
2	Patient details				
2.1	Internal ID number:				
2.2	Date of poisoning:				
2.3	Sex:				
2.4	Age:				
3	Exposure details:	Yes / No	Remark		
3.1	Suspected?				
3.2	Confirmed?				
	How confirmed:				
3.3	Cyanide in blood?				
3.4	Thiocyanate in urine?				
3.5	Cyanide container (e.g. bottle) presented?				
3.6	Smell of bitter almonds?				
3.7	Lactic acidosis?				
3.8	Cyanide toxidrome (abnormal neurological status, respiratory pattern and heart rate)?				
3.9	Other?				
4	Substance details	Yes / No	Oral	Inhalation	Skin
4.1	HCN?				
4.2	Cyanide salt?				
4.3	Nitrile?				
4.4	Natural e.g. cassava, almond kernels, etc.?				
4.5	Sodium nitroprusside?				
4.6	Smoke?				
4.7	Intoxication mixed with ...?				
5	Poison Severity Score	Yes / No			
5.1	Severe?				
5.2	Moderate?				
5.3	Minor?				
6	Antidote treatment used	Yes / No	Dose	Time	Remark
6.1	Oxygen 100%?				
6.2	AN?				
6.3	SN?				
6.4	4-DMAP?				
6.5	STS?				
6.6	Hydroxocobalamine?				
6.7	Dicobalt edetate?				
6.8	Other (specify)?				
7	Outcome	Yes / No	Remark		
7.1	Death?				
7.2	Recovery?				
7.3	Other (sequelae)?				

20 Responses received (10-Aug-10)

- Anonymous. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 0 cyanide / nitrile cases]. Personal communication. Laboratory for Clinical and Forensic Toxicology and Drug Analyses, University Hospital, Groningen, Netherlands.
- Baud F. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 14 cyanide cases, June 2009]. Personal communication. Lariboisière Hospital, Paris, France.
- Botti P. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 4 cyanide cases]. Personal communication. Centro Antiveleni Firenze, Azienda Ospedaliero Universitaria Careggi, Firenze, Italy.
- Busch H. 2010. ECETOC questionnaire on cyanide and nitrile poisoning [response for 144 cases]. BASF, Ludwigshafen, Germany.
- Casey P. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 4 cyanide and 3 nitrile cases]. Personal communication. National Poisons Information Centre, Beaumont Hospital, Dublin, Ireland.
- Ebbehøj N. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 2 cyanide cases 2006-2009]. Personal communication. Arbejds- og Miljømedicinsk Klinik, Bispebjerg Hospital, København, Denmark.
- Hoppu K. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 0 cases]. Personal communication. Poison Information Centre, Helsinki University Central Hospital, Helsinki, Finland.
- Jacobsen D. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 5-10 cyanide cases per year]. Personal communication. Department of Acute Medicine. Ullevål University Hospital, Oslo, Norway.
- Küpfer U. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 9 cyanide cases 1977-2008]. Personal communication. Evonik Degussa, Wesseling, Germany.
- Martínez Arríeta R, Tena Quintero F. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 27 cyanide or nitrile cases 2006-2008]. Personal communication. Instituto Nacional de Toxicología y Ciencias Forenses, Madrid, Spain.
- Pelclová D. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 3 cyanide cases 1990-1996]. Personal communication. Charles University in Prague, 1st Faculty of Medicine, Prague, Czech Republic.
- Plackova S. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 1 cyanide case]. Personal communication. National Toxicological Information Center, Bratislava, Slovak Republic.
- Pöld K. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 0 cases 2008-2009]. Personal communication. Estonian Poison Information Centre, Tallinn, Estonia.
- Rödelsperger R. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 17 cyanide and 2 nitrile cases]. Personal communication. Evonik Röhm, Darmstadt, Germany.
- Sauer O. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 88 cyanide and 20 nitrile cases]. Personal communication. Giftinformationszentrum der Länder Rheinland-Pfalz und Hessen, Klinische Toxikologie, Universitätsklinikum, Mainz, Germany.
- Smet H. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 17 cases]. Personal communication. Poisons Centre, Hospitaalcentrum van de Basis Koningin Astrid, Brussels, Belgium.
- Steffens W. 2008. ECETOC questionnaire on cyanide and nitrile poisoning [response for 7 cases]. Personal communication. Bayer CropScience, Monheim, Germany.
- Whyte I. 2009. ECETOC questionnaire on cyanide and nitrile poisoning [response for 3 cyanide cases]. Personal communication. Hunter Area Toxicology Service, Newcastle, UK.
- Zilker T. 2007. ECETOC questionnaire on cyanide and nitrile poisoning [response for 8 cases]. Personal communication. Center Klinikum rechts der Isar, Toxikologische Abteilung. Technical University, Munich, Germany.
- Zilker T. 2010. ECETOC questionnaire on cyanide and nitrile poisoning [response for 1 cyanide case]. Personal communication. Klinikum rechts der Isar, Toxikologische Abteilung. Technical University, Munich, Germany.

APPENDIX B: DATABASE OF > 400 CASES

Please see separate spreadsheet issued with this report.

APPENDIX C: CASUISTICS

New actions:

GCS Glasgow coma scale

CONTENTS:	147
Akintowa and Tunwashe, 1992*	152
Baader and Wrbitzby, 2006	152
Bain and Knowles, 1967	152
Ballesteros et al, 2003	152
Baud et al, 1986*; Bismuth et al, 1987*	153
Baud et al, 2002*	153
Baud and Borron, 2008*	153
Beamer et al, 1983	154
Benaissa et al, 1995	154
Berlin, 1970	155
Berumen, 1983	156
Bismuth et al, 1984	156
Bismuth et al, 1983	156
Boggild et al, 1990	157
Bonsall, 1984	157
Borgohain et al, 1995	158
Borron, 1996, 2007	158
Bouachour et al, 1990 (Harry et al, 1985)	158
Bourrelier and Paulet, 1971	159
Brian, 1990	159
Brivet, 1983	159
Brown et al, 1987	160
Brueske, 1997	160
Bryson, 1978	160
Buchanan, 1976	160
California Bureau of Communicable Disease Control, 1972	161
California Bureau of Communicable Disease Control, 1975	161
Campbell and Jones, 2001	161
Caravati and Litowitz, 1988	162
Carella, 1988	162
Cesion and Jurriling, 2009	162
Cetnarowski and Conti, 1986	163
Chen and Rose, 1952	163
Cheok, 1978	165
Cherian and Richmond, 2000	165
Chin and Calderon, 2000	165
Cliff and Coutinho, 1995	166
Coentrão et al., 2010a (Coentrão and Moura, 2010b)	166
Coentrão and Moura, 2011	166
Dall and Hannah, 1964	167
Daunderer et al, 1974	167
Davies, 1975	168
Davis and Ewer, 1988	168

Dawood, 1969	168
De Busk and Seidel, 1969	169
Dequidt et al, 1974	169
DiNapoli et al, 1989	170
Dodds and McKnight, 1985	171
Edwards and Thomas 1978	171
Eiró Gonsalves et al, 1956	171
Espinoza et al, 1992	172
Favarel-Garrigues, 1982 / Dreykluft et al, 2004 (case 1)	172
Feihl et al, 1982	173
Feldmann, 1990	173
Fernández et al, 2008	173
Finelli, 1981	174
Fortin et al, 2007	174
Froldi et al, 2001	174
Gaillard, 1991	174
Gambaro et al, 2007	175
Gaultier and Le Breton, 1952	175
Geller et al, 1991	175
Ghiringhelli, 1955	176
Glenn et al, 1994	176
Goodhart, 1994	176
Graham et al, 1977	177
Grunske, 1949	177
Gunder et al, 1969	177
Hall et al, 1986	178
Hall et al, 1987	178
Hall et al, 1989a	179
Hall et al, 2009	179
Hantson, 1996	179
Harry, 1985	180
Heintz et al, 1990	180
Herbert, 1979	180
Hill, 1942	181
Hillman et al, 1974	181
Hoang The Dan et al, 1981	182
Humbert et al, 1977; Braico et al, 1979	182
Humphrey, 1978	183
Hung et al, 2009	183
IPCS, 1993a	183
IPCS, 1993b	184
Jacobs, 1984	184
Jaeger et al, 1977	184
Jamali, 1993	185
Jeannin et al, 1961	185
Jerretin, 1963	185
Johnson and Mellors, 1988	186
Johnson et al, 1989	186
Jones, 1992	187
Jouglard et al, 1974	187
Jourdan et al, 1993	187
Kampe et al, 2000	187

Kasamo, 1993	188
Kim et al, 1982	188
Kirk et al, 1993	189
Kläui et al, 1984	190
Klöver and Wenderoth, 1965	191
Kreff, 1955	191
Krieg and Saxena, 1987	191
Küpper, 2009	192
Kurt et al, 1990	193
Lam and Lau, 2000	193
Lang and Stintzy, 1960	194
Lasch and El Shawa, 1981	194
Lawson-Smith et al, 2010	195
Lazarus-Barlow and Norman, 1941*	195
Lee et al, 1982	196
Lee-Jones et al, 1970	196
Leor et al, 1986	197
Liebowitz and Schwartz, 1948	197
Litovitz et al, 1983	198
Lorz, 1950	199
Losek et al, 1991	199
Lundquist, 1992	199
Lurie, 1953	200
Lutier et al, 1971	201
Mannaioni, 2002	202
Marbury, 1982	203
Martín-Bermúdez, 1997	203
Martinelli et al, 2008	204
Mascarenhas et al, 1969	204
Maxwell, 1978	204
McKiernan, 1980	205
MacRae, 1974	205
Mellino, 1980	205
Merryfield, 1974	205
Messing, 1991	206
Michaelis et al, 1991	206
Michenfelder and Tinker, 1977	206
Miller and Toops, 1951	206
Mlingi et al, 1992	207
Montoliu et al, 1979	207
Morse et al, 1979	207
Morse et al, 1981	208
Moss et al, 1981	208
Motin et al, 1970	208
Mueller and Borland, 1997	208
Muraki et al, 2001	209
Musshoff et al, 2002	209
Mutlu, 2002	210
Nagler et al, 1978	211
Nakatani et al, 1992	211
Naughton, 1974	212
Nicoletta et al, 2007	213

O'Brien et al, 2005	213
Ortega and Creek, 1978	213
Oruřev et al, 1972	214
Pardee, 1947	214
Patel, 1986	214
Paulet, 1965	215
Pentore et al, 1996	215
Perschau et al, 1977	216
Peters et al, 1982	216
Pijoan, 1942	217
Posner, 1977	217
Potter, 1950	217
Prieto, 2005	218
Queisser, 1966	218
Quinlan et al, 2008	219
Rachinger et al, 2002	219
Racle et al, 1976	219
Ram et al, 1989	219
Ruangkanchanasetr et al, 1999	220
Rauscher, 1978	220
Riudavets et al, 2005	220
Robinson, 1959	220
Roedelsperger, 2009	221
Rosenberg et al, 1989	221
Rubino and Davidoff, 1979	222
Sadoff et al, 1978	222
Saincher and Swirsky, 1994	222
Sarikaya, 2006	222
Sartorelli, 1966	223
Schulz et al, 1979	223
Schulz et al, 1982	223
Schulz and Roth, 1982	223
Scolnick et al, 1993	224
Selden, 1990	224
Shragg et al, 1982	225
Singh et al, 1989	225
Sipe et al, 2001	226
Smith, 1977, 1978	226
Steffens et al, 1998; Steffens, 2002	227
Steffens, 2003a,b	227
Steffens et al, 2003	228
Steffens, 2010	228
Stickel, 2008 (Zilker and Eyer, 2005; Zilker and Schweizer, 1987)	229
Suchard et al, 1998	232
Sutter et al, 2010	232
Tartière et al, 1992	232
Tassan et al, 1990	232
Thiess and Hey, 1969	233
Thomas and Brooks, 1970	233
Townsend and Boni, 1975	233
Trapp, 1970	233
Turchen and Manoguerra, 1989; Turchen et al, 1991	234

Uitti, 1985	234
Valenzuela et al, 1992	235
Zaknum (2005) + Rachinger (2002), 2005	235
van Dijk et al, 1987 / van Heijst et al, 1987	235
van Heijst et al, 1987 / Douze et al, 1987	236
Vogel et al, 1984	236
Vogel, 1991	237
Wananukul and Kaojarern, 1992	237
Weng et al, 2004	237
Werner, 1979	238
Wesseon, 1985	238
Wolfsie, 1951	239
Wright and Vesey, 1986	239
Wu et al, 2001	240
Würzburg, 1996	241
Yacoub et al, 1974	241
Yatziv and Simcha, 1969	242
Yeh et al, 1992	242
Yen et al (1995)	242
Zaknum, 1905	243
Zavotsky et al, 2004	243
Zilker, 2009	244
Zilker and Eyer (2002)	244
Zilker and Eyer, 2005	245
Zilker, 2010	245

Akintowa and Tunwashe, 1992*

Three persons ingested a cassava-based meal ('Gari') and died shortly after hospital admission. The patients vomited and complained of abdominal pain immediately after the meal. They were unconscious with renal failure and died of cardiopulmonary arrest. The cyanide levels in the blood were 1.15, 0.85 and 1.35 mg/l, respectively and urine averaged 0.54 mg/l. Cassava contains cyanogenic glycosides which slowly release cyanide and this may have been responsible for the death of these patients. There is an urgent need to establish maximum tolerable levels of cyanide in 'Gari' and other cassava food products. *Comment: No antidotes were applied, only symptomatic treatment for food poisoning was given with no effect.*

Baader and Wrbitzby, 2006

A cleaning worker in a railroad tank wagon has contained acrylonitrile. Neither acrylonitrile nor cyanide levels are reported as the paper deals with adduct monitoring in colleagues, rescue personnel and hospital staff, who had no symptoms of poisoning.

Bain and Knowles, 1967

A 61 year old patient employed in a factory dealing with cyanides was seen by the plant physician after a distinct collapse. He was semi-comatose, had seizures, a tachycardia of 120/min and was feeling nauseous and vomiting. Two doses of 150 mg of Co-EDTA (in total 300 mg of the antidote) were administered i.v. within a 10 minute period as his clinical state had aggravated. After the first dose the patient vomiting once more, became coherent and responded to questions, and the convulsions ceased. 2 hours afterwards, the patient had apparently completely recovered. A blood sample taken just before the administration of the antidote revealed a blood cyanide concentration of 5.1 mg/l. *Comment: A case of moderate poisoning, yet high cyanide level in blood.*

Ballesteros et al, 2003

A 1.5-month-old infant under SNP infusion (initial dose 1µg/kg/min, increased to 5.5 µg/kg/min after 12 hours with a total of 36 hours) developed somnolence under the infusion. Convulsions set in 6 hours after termination of the SNP treatment. Cardiorespiratory arrest required intubation and resuscitation. Catecholamines were needed to stabilise circulation. The infant developed signs of hypoxic encephalopathy. Under suspicion of cyanide intoxication, an unspecified dose of HOCO was given i.m. and prompted 'spectacular' improvement of the neurological signs. A mild residual paresis of the right arm and axial hypotension persisted. The cyanide level,

however, measured 8 hours after cessation of the SNP infusion, but before HOCO application, was not elevated. *Comment: This case does not enable any assessment due to insufficient reporting of data.*

Baud et al, 1986*; Bismuth et al, 1987*

A worker tried to repair a pump wearing neither breathing protection nor adequate protective clothing. He rapidly lost consciousness. After arrival in the intensive care unit, he was decontaminated and received i.v. 4 g of HOCO and 8 g of STS combined over 30 minutes. During the infusion, he regained consciousness and recovered. The blood cyanide level was 5.71 mg/l.

Baud et al, 2002*

A retrospective clinical study was made of 11 acute cyanide-poisoned patients. The aim was to test the hypothesis that plasma lactate concentrations could be of confirmatory value in patients with histories consistent with acute pure cyanide poisoning because immediate laboratory confirmation of suspected cyanide poisoning is rarely possible and because clinicians must rapidly decide whether to administer specific antidotes, which may have severe side effects.

All 11 patients were admitted to the intensive care unit from 1988 to 1999; fire victims were excluded. Before antidotal treatment, the median plasma lactate concentration was 168 mg/dl, the median blood cyanide concentration was 4.2 mg/l. Using Spearman's test, there was a significant correlation between plasma lactate and blood cyanide concentrations ($r = 0.74$, $p = 0.017$). Before antidotal treatment, plasma lactate concentration correlated positively with anion gap and inversely with systolic blood pressure, spontaneous respiratory rate, and arterial pH. During the course of cyanide poisonings, a plasma lactate concentration of ≥ 72 mg/dl (8 mmol/l) was sensitive (94%) and moderately specific (70%) for a toxic blood cyanide concentration (≥ 1.0 mg/l). The specificity was substantially improved in patients not receiving catecholamines (85%). The immediate and serial measurement of plasma lactate concentrations was considered useful in assessing the severity of cyanide poisoning.

Baud and Borron, 2008*

Case 1: A 42 year old male with smoke inhalation was found in cardiac arrest. He was resuscitated, intubated, mechanically ventilated with pure oxygen, and received colloidal fluids. Spontaneous circulation returned as well as consciousness. His blood pressure was 130/80 and plasma lactate was 12.8 mmol/l. He then received 5 g of HOCO prior to a session of hyperbaric

oxygen (HBTO) and survived. *Comment: In this case, improvement occurred before HOCO application.*

Case 2: A 37 year old male was also found in cardiac arrest after smoke inhalation. Resuscitation, intubation and mechanical ventilation with pure oxygen were installed, after which spontaneous circulation was achieved. Immediately after 5 g of HOCO spontaneous respirations returned but he remained comatose. He survived after administration of hyperbaric oxygen. *Comment: A case of sever poisoning in which symptomatic therapy improved the status, and further improvement was seen after HOCO. However, coma did not improve after HOCO, but did after hyperbaric oxygen.*

Beamer et al, 1983

This case of Laetrile poisoning is described as occurring as a result of a self administered medication error. A patient with heavy cancer had gone to Mexico to obtain Laetrile in the hope to treat his poignancy. As he had missed several days of his medication and took 12 to 18 tablets of Laetrile to catch up. He showed an acute onset of grand mal seizures. The time between intake and onset of symptoms is not stated. At admission, the blood pressure was 140/100 mm Hg and the heart rate 88 beats/min. He was tachypnoeic with a respiration rate of 32/min. The pupils were fixed and dilated, the skin was pink, and the patient was comatose. His arterial blood gas demonstrated a metabolic acidosis with ineffective respiratory compensation (pH: 7.09, PaO₂: 135 mm Hg, PaCO₂: 9 mm Hg, HCO₃: 3 mEq). The Lilly Cyanide Antidote Kit was used. AN was administered for 30 sec/min by inhalation over 3 min. The patient was intubated and ventilated with 100% oxygen. Then he received 0.3 g of SN i.v. followed by 12.5 g of STS. The blood gases did not improve within the next 5 minutes, bicarbonate was given and a second dose of 0.3 g of SN was administered when the MetHb level reached 6.6%. Within 5 minutes, the blood gas improved significantly (pH: 7.21, PaO₂: 562 mm Hg, PaCO₂: 26 mm Hg, HCO₃: 10 mEq). The MetHb level went up to 10.5%. Due to uncontrollable violent seizures, it was necessary to induce a neuromuscular blockade by pancuronium. The patient rapidly recovered full neurological function within 20 hours. *Comment: A case of severe poisoning in which treatment with SN and symptomatic measures were successful.*

Benaissa et al, 1995

Case 1: A female pharmacist (age 27) ingested mercury-oxy-cyanide. She developed chest and stomach pain and lost consciousness soon after. A quarter of an hour later the ambulance staff assessed the following parameters: blood pressure systolic 80 mm Hg, pulse 34/min, GCS 3 points and both pupils dilated and non-reactive to light. The patient was intubated and ventilated and because of cardio-circulatory failure cardiopulmonary resuscitation (CRP) was begun, which

was continued after admission to the ED. During the resuscitation, 4 g of HOCO and 8 g of STS were administered i.v. Half an hour later pulse and blood pressure had stabilised. 1 hour after the antidote application, she developed disseminated intravascular coagulation (DIC) with multiple internal bleedings followed by hypovolemic and haemorrhagic shock with anuria. The specific mercury antidotes dimercaptole and dimercaptosuccinate were given but without success. The woman died 5 hours later. The levels of cyanide and mercury had been determined once during the course with 6.99 mg/l (cyanide) and 0.44 mg/l (mercury). *Comment: A case of severe poisoning with initial cardiac arrest, HOCO prompted cardio-circulatory improvement, yet the patient died of DIC. The anuria probably was due to mercury poisoning.*

Case 2: A 14 year old boy had taken an unknown amount of mercury-cyanide. Approximately 50 minutes after the intoxication he was conscious with tachycardia (heart rate 120/min) and complained about diarrhoea, nausea, vomiting and stomach pain. During the next minutes he became agitated, hypotonic (decrease of the systolic blood pressure from 100 to 80 mm Hg) and more tachycardic (heart rate 140/min). The blood gases showed a pH of 7.33 and a lactate level of 19.6 mmol/l. The treatment consisted in emptying the stomach and administering i.v. fluids, dimercaprol (200 mg i.m. every 6 hours), dimercaptosuccinate (400 mg p.o. every 8 hours) and 5 g of HOCO. His condition improved temporarily. 7 hours later, he became anuric. He was given furosemide and dopamine, without any positive reaction. 5 days later peritoneal dialysis was initiated (creatinine of 1325 µmol/l). The cyanide level was determined 4 times, together with the lactate levels: at admission (0) and after 1, 5 and 11 hours: Cyanides (µmol/l): 217 , 35 , 4.6 and 13 (corresponds to 5.64, 0.95, 0.12 and 0.35 mg/l). He recovered completely and could be discharged after 26 days. *Comment: Apart from the anuria, this was only a case of mild cyanide poisoning, in which HOCO improved the status. The anuria was probably due to mercury poisoning.*

Berlin, 1970

A 17-month-old child drank about 7.5 ml of Drabkin's solution (contains sodium bicarbonate, potassium ferricyanide, and potassium cyanide), containing 50 mg/l of KCN which approximates 1/50th of a lethal dose. The child was rushed to the emergency department and given 0.3 g of SN followed by 12.5 g of STS (adult doses). Gasping resulted, interpreted as continuing cyanide poisoning. A second dose of 150 mg of SN was given with 12.5 g of STS. He vomited, had apnoeic spells and seizures. After treatment with bicarbonate, artificial ventilation, diazepam for seizures, he was noted to have brown blood. 15 mg of methylene blue was given i.v. then exchange transfusion begun. He suffered a cardiac arrest and died. Post-mortem cyanide concentration was 0.10 mg/l (10 µg/dl). *Comment: The MetHb concentration was not reported. However, this was a clear overdose of SN in a toddler that obviously had fatal outcome.*

Berumen, 1983

A 23 year old student saw his dog suddenly collapse and started external cardiac massage and mouth-to-nose ventilation effort. Moments later the dog died and the student felt nauseated, vomited, and lost consciousness. At the hospital an alert medical officer detected a bitter almonds odour on the patient's breath and administered the accepted treatment for cyanide poisoning, after which he recovered. It turned out that the dog had accidentally swallowed cyanide, and the volatile poison eliminated through the dog's lungs had been inhaled by his master during the mouth-to-nose artificial respiration. *Comment: Probably a case of mild poisoning but no information which antidote was administered.*

Bismuth et al, 1984

Case 1: This patient was found in cardiorespiratory arrest, his cyanide level was later found to be 1.8 mg/l. Resuscitation and application of Co-EDTA and HOCO doses were without success.

Case 2: A patient in coma with cardiorespiratory arrest, convulsions and circulatory failure. Resuscitation and application of Co-EDTA and HOCO could not amend the coma, which persisted until the patient died 6 months later. Cyanide level had been 6.8 mg/l initially.

Case 3: A patient in coma with respiratory arrest, circulatory failure. Resuscitation, Co-EDTA and HOCO were applied. The patient recovered completely, though the initial cyanide level had been 6 mg/l.

Comment: In all 3 cases, it is impossible to assess the effect of HOCO, as Co-EDTA was also administered. In 2 cases, the antidotes could neither reverse cardiac arrest nor improve coma. In the third case, the patient survived without sequelae indicating that the application of both antidotes was successful.

Bismuth et al, 1983

A retrospective study of 25 cases of cyanide poisoning brought to light the following points: cardiorespiratory arrests are frequent (7/25) and often inaugural; in severe intoxications (7/25), deep metabolic acidosis is the rule, and cyanide poisoning should always be suspected in cases of coma with severe acidosis; mild intoxications are frequently symptomless. Anxiety and agitation should not be considered as evidence of cyanide poisoning; they are merely due to fear in most cases. The present treatment of acute cyanide poisoning relies basically on symptomatic measures including sodium bicarbonate, cardiac massage and, above all, assisted ventilation with 100% oxygen. The author's experience did not support the concept of a lethal cyanide blood

level when patients can rapidly be transferred by a medical team to an intensive care unit. Survival depends more on prompt medical care than on the accessibility to sophisticated antidotes. *Comment:?????*

Boggild et al, 1990

A 22 year old female was found semiconscious at her place of work. On admission to hospital she was conscious. Her friends reported that she had probably taken an unknown poison approximately 8 hours before admission. She resisted gastric lavage. Clinical examination was unremarkable apart from transient sinus tachycardia. She remained drowsy but arousable over the next 18 hours, but then rapidly developed agitation followed by grand mal convulsions and asystolic cardiac arrest. Additionally a profound metabolic acidosis was found and the patient remained hypotensive. She died 30 hours after the supposed ingestion. Later investigation revealed that the patient had taken acetone and acetonitrile. Acetone is known to slow the metabolism of acetonitrile to cyanide. *Comment: A case of severe poisoning but no antidote was given.*

Bonsall, 1984

An accidental combined (skin and inhalation) cyanide intoxication occurred when a tank containing nitriles was washed with water. According to the authors HCN gas was liberated through this action. A male worker entered the tank to inspect it. He was wearing an overall and boots. Within 3 minutes he collapsed, lost consciousness and suffered from respiratory failure. His pupils were dilated. As no one was aware of the possibility of cyanide intoxication and decontamination was not performed, but he was ventilated mouth-to-mouth by a colleague. He arrived in the emergency department about 40 minutes after the incident. There he presented a coma with a positive Babinski's sign^a. Then he was decontaminated and an unspecified dose of STS injected twice. As the patient convulsed and vomited he was intubated and ventilated. His condition improved over the next hours so that he could be extubated 48 hours after admission. 72 hours after the intoxication he was conscious and orientated. He was discharged 14 days later in complete remission. *Comment: A case of severe poisoning in which STS and supportive treatment were effective.*

^a big toe goes up when the outside of the sole is stimulated

Borgohain et al, 1995

A 27 year old woman was admitted with stiffness and weakness of all 4 limbs making her more or less bedridden for the past 5 years. These symptoms started a month after she had attempted by ingesting approximately 300 mg of KCN procured from her father's place of work (goldsmith). She developed delayed onset of persistent generalised dystonia. Cranial CT scan showed bilateral putaminal hypodensities which were also seen on MRI scans to be hypotense on T1 and hypertense on T2 weighted images. Multimodality evoked potentials were normal. An improvement was noted with levodopa. No antidote was administered. *Comment: antidote efficacy cannot be judged as treatment was several years after intoxication and in any event no antidote was given.*

Borron, 1996, 2007

Borron reviewed a total of 18 clinical reports of the use of high dose hydroxycobalamin including case reports cited by Yacoub (1974), Jougla (1974), Bourneller (1971), Bismuth (1984), Racle (1976), Motin (1970), Lutler (1971), Danel (cited by Broward 1987), Dougharty (1990), Tassan (1990) and Tarti re (1992). HOCO was administered alone in 5 cases, and possible a further 2, or in combination with with EDTA (6), STS (2), EDTA/Nitrile (1) or EDTA/Nitrile/STS (1). Cyanide intoxication was confirmed in 7 cases by blood cyanide determinations. Recovery was complete in 15 cases, with sequelae in 1 (after 7gms of HOCO and STS) chronic coma in a second case and death in a third (both with CN poisoning confirmed and after 4gms of HOCO and EDTA).

Bouachour et al, 1990 (Harry et al, 1985)

A 54 year-old woman ingested 200 mg of NaCN and was found 30 minutes later in deep coma. The patient was intubated and ventilated with 100% oxygen. 1 hour after the incident and upon arrival in hospital, the patient presented a low score (3 points) on the GCS, hypotonia (blood pressure 70/40 mm Hg), hypothermia (33 C), tachycardia (heart rate 120/min), and high levels of lactic acid (29.5 mmol/l) and blood cyanide (10 mg/l). The patient was given 4 g of HOCO and 8 g of STS i.v. This resulted in a normal peripheral resistance, a normal aortic pressure and an improvement on the GCS (increase to 9 points), while lactate decreased to 7.5 mmol/l. Additional hyperbaric oxygen was initiated. Under this treatment, the patient's neurological status normalised completely (GCS 15). The cyanide level during further hospitalisation dropped to 1.2, 0.3 and 0.25 mg/l after 4, 6 and 20 hours, respectively. (The case was communicated earlier by Harry et al, 1985.) *Comment: A case of severe poisoning in which HOCO and STS applied together were partially successful for the neurological symptoms.*

Bourrelier and Paulet, 1971

During an accident implicating molten NaCN, both exposed persons were treated with two ampoules and more of Co-EDTA. One of the victims, a 50 year old male, was in a coma and received 2 ampoules of Co-EDTA but had regained consciousness by the end of the infusion. He received oxygen and glucose. The other, a 27 years old male, was in a profound coma, apnoeic and in a state of intense shock. He immediately received 100% oxygen after intubation and initially three ampoules of Co-EDTA and then two further ampoules. In addition, he received two doses of 100 mg of HOCO. The cardiovascular state normalised. 1 hour after administration of the five ampoules adverse effects attributed to the Co-EDTA were noted including a phase of tachycardia with hypertension and profuse sweating. 2½ hours later, the gasping had disappeared being replaced with irregular spontaneous respirations which subsequently regularised. He recovered but had some difficulties with speech. *Comment: A case of severe poisoning which indicated that 200 mg of HOCO in an adult represents an inefficacious dose. Co-EDTA was fully effective for the cardiovascular, but only partially for the neurological status.*

Brian, 1990

A case of a 7 year old girl year old that developed weakness, tremor and vomiting one hour after eating cassava was reported from Papua New Guinea. After 4 to 65 hours later she was conscious, but still had tremor and restlessness. She was given 2 ampoules of Co-EDTA without clinical effect/improvement, but consecutive facial oedema. *Comment: A case of mild poisoning in which no cyanide or thiocyanate measurements were performed and Co-EDTA was not effective.*

Brivet, 1983

A 43 year old male analytical chemist attempted suicide by ingesting KCN (unknown quantity). Ten minutes after ingestion he developed a headache and 10 minutes later his blood pressure dropped and he became comatose. The first treatment comprised the application of 100% oxygen and gastric lavage. The patient developed seizures followed by cardiac arrest. He was resuscitated by CPR and administered with catecholamines, sodium bicarbonate and i.v. fluids. This resulted in a blood pressure of 80/50 mm Hg and a heart rate of 130/min. The pH was 6.8 at admission and 7.34, 7.49 and 7.47 after 2, 12 and 18 hours respectively. The lactate level was measured at the beginning and after 12 hours with 20 mmol/l and 1.96 mmol/l respectively. The patient developed a mild rhabdomyolysis probably due to the reanimation procedure. He recovered completely during the next 20 hours. *Comment: No antidote was used. Comment: A case of moderate poisoning in which HOCO reversed the neurological symptoms.*

Brown et al, 1987

A 24 year old man was cleaning a vat containing silver cyanide when he was overcome by fumes and collapsed into several inches of slurry. Upon arrival in the casualty emergency department 15 minutes later he was unresponsive, apnoeic, and hypotensive. His skin was brick red, which is indicative of cyanide toxicity. After resuscitation, administrations of 300 mg of Co-EDTA were given at 1, 2, 3, and 7 minutes. Spontaneous respiration resumed, although he remained haemodynamically unstable. 24 hours after admission urine output and blood pressure were satisfactory. Electroencephalograms obtained at this time and 48 hours after admission showed no cerebral activities. As he remained haemodynamically stable and blood cyanide values were rapidly falling to non-toxic concentrations the possibility of organ donation was considered. Both kidneys were perfused immediately after removal and transplanted within 24 hours. *Comment: A case of sever poisoning in which respiration and cardiovascular status could be improved by Co-EDTA, yet the brain damage could not be reversed.*

Brueske, 1997

A 35 year old man was exposed to copper cyanide gas while attempting a valve repair. He developed nausea, dizziness and lethargy/disorientation. A nurse practitioner administered AN for 10 minutes. After about 45 minutes he was taken to hospital. In the ED, he was dizzy and short of breath, with hypertension, tachycardia and tachypnoea. He remained lethargic and disoriented. He was given 300 mg of SN followed by 12.5 g of STS. His level of consciousness rapidly improved, but he remained tachycardic. He was admitted to the ICU where he stayed for 24 hours before being discharged the following day without sequelae. *Comment: A case of moderate poisoning in which the "Lily-Kit" was effective.*

Bryson, 1978

This is a comment to a publication in Lancet (1977; **11**, p1167) in which casualty officers were instructed to offer Kelocyanor for all patients displaying symptoms of cyanide poisoning. Bryson responded saying that in the past 2 years several cases of unpleasant reactions to Kelocyanor had (600mg) been referred to him and cautioned the need for clinical evaluation and case history before its use (see also Edwards and Thomas 1978).

Buchanan, 1976

A 23 year old goldsmith attempted suicide by ingesting KCN. He was found collapsed only 1.5 hours later. In the ED he presented a coma, with shallow respiration and a pink coloured skin.

The heart rate was weak at 40-50/min and the systolic blood pressure was 110 mm Hg. Five minutes after the clinical examination respiration ceased making intubation and mechanical ventilation necessary. The bradycardia persisted and the blood pressure dropped to 0 mm Hg systolic. The young man received dextrose, sodium chloride, noradrenaline, cortisone, aramine and methoxamine but this did not improve his condition. With adrenaline and isoprenaline infusions, a blood pressure of 80 mm Hg could be established. After 2.5 hours a blood level of 5 mg cyanide/l was determined he was treated with 0.3 g of Co-EDTA and 4 g of glucose. 1 hour later the patient was also given 0.5 g of SN and 25 g of STS. The antidotal treatment was completed with 3 g of HOCO 30 minutes after the nitrite application. There was no change in the man's condition and 17.5 hours after the suicide attempt cardiac arrest occurred. CPR failed and the man died 37.5 hours after the incident because of irreversible asystolia. *Comment: A case of severe poisoning in which the delayed application of several antidotes, including HOCO, could not save the patient.*

California Bureau of Communicable Disease Control, 1972

In California a couple shared a drink of pureed apricot kernels (total of 30) and dried apricots. 1 hour later the woman complained of abdominal discomfort, tachycardia and a strange feeling, her husband also developed headache, light-headedness, tachycardia and a strange feeling. *Comment: This is a case of mild poisoning but no antidotes were given nor cyanide measurements reported.*

California Bureau of Communicable Disease Control, 1975

In California a 56 year old woman ate a handful of apricot kernels and developed a headache, light-headedness, tachycardia, flushing and weakness 1 hour later. Vomiting was induced and the vomit was found to contain 1.7 mg/l cyanide, while she had no cyanide in blood. *Comment: This is a case of mild poisoning but no antidotes were given.*

Campbell and Jones, 2001

A 30 year old male ingested KCN. 1 hour later he suffered cardiac arrest and was resuscitated. SN at an unspecified dose was not successful, so he then received 2.5 g of HOCO. While the cardiovascular situation improved, there was no CNS improvement, and he was declared brain dead after 5 days. This was attributed to cardiac arrest hypoxia. Cyanide in blood had been found to be above 4 mg/l. *Comment: In this case HOCO improved the cardiovascular, but not the neurological situation after cardiac arrest.*

Caravati and Litowitz, 1988

A case of dermal/inhalation uptake is reported in a 2 year old child, surviving a cyanide level of 6 mg/l.

2 cases of paediatric accidental ingestion of an acetonitrile-containing cosmetic are reported. 1 of the children, a 16-month-old boy, was found dead in bed the morning after ingesting the product. No therapy had been undertaken, as the product was mistakenly assumed to be an acetone-containing nail polish remover. The second child, a 2 year old boy, experienced signs of severe cyanide poisoning, but survived with vigorous supportive care. Both children had blood cyanide levels in the potentially lethal range. The observed delayed onset of severe toxic reactions supports the proposed mechanism of acetonitrile conversion to inorganic cyanide via hepatic microsomal enzymes. *Comment: These are cases of severe poisoning but no antidotes were given.*

Carella, 1988

A 46 year old woman and her husband were brought to hospital after drinking from a bottle of a soft drink which had been adulterated with cyanide. The man died before arriving at the hospital. On admission, his wife was found to be unconscious, readily reacting to pain and moving her limbs without obvious symmetry. Babinski's sign was present on the right. An analysis of gastric aspirates from the patient and her husband confirmed the presence of cyanide. CT scan and cerebrospinal fluid examination were normal. EEG showed occasional decerebrate posturing, but she slowly recovered after 2 weeks. She was transferred to a rehabilitation unit where she was found to have dystonia and dysarthria, slight plastic limb hypertonia, slight hemiparesis with right Babinski's sign and diffuse hyperactivity of tendon reflexes. 1 year later, the patient was admitted to another hospital because of increasing speech and swallowing difficulties, drooling and unsteadiness of gait. Subsequently, drooling and dysphagia became more marked and sustained abnormal postures of the mouth and tongue developed. 5 years after poisoning, the patient was readmitted to the rehabilitation unit. In conclusion this case confirms that Parkinsonism, dystonia and lid opening apraxia may evolve after cyanide poisoning.

Cesion and Jurriling, 2009

The authors report on a 54 year old man suffering from smoke inhalation. The COHb was 29% and although HCN poisoning was suspected, this was not confirmed by measurement. Lactate was high 16 mmol/l. The patient was treated with hyperbaric oxygen and 5 g of HOCO. The course is not reported. *Comment: As it is unclear, whether the patient really suffered cyanide poisoning, the efficacy of HOCO cannot be assessed in this case. No side effects were reported.*

Cetnarowski and Conti, 1986

A patient received a total dose of 215 mg (3.07 mg/kg) of SNP over 33 hours. The patient showed unexplained and worsening nausea, vomiting, mental alterations and metabolic acidosis with dramatic improvement after the admission of amyl nitrite. The authors commented that the administered dose was well below any recommended total maximum dose. *Comment: A case of moderate poisoning in which AN showed improvement.*

Chen and Rose, 1952

Chen and Rose reported 16 cases of cyanide poisoning treated with parts of their cyanide antidote kit, among which 10 received SN after exposures to HCN gas/fumes.

Case 1: A 24 year old man tried to start a syphon by mouth and thereby accidentally swallowed a solution containing 2.3% silver cyanide and 6.9% NaCN. He became weak, had a burning sensation in the throat and felt his heart pounding. 45 minutes later he showed fine tremors together with a very rapid pulse and received 10 ml of a 3% SN (300 mg NaNO₂) and 50 ml of 12.5% STS solution (6.25 g STS). He was discharged from the clinic after 24 hours. *Comment: A case of mild poisoning only and recovery is not necessarily attributable the use of antidotes.*

Case 2: A man was found unconscious in a HCN fumigated barracks and given artificial respiration and an unknown amount of methylene blue. Despite absence of respiration and being regarded as hopeless he was later given 0.3 g of SN and 1 g of STS i.v. Both antidotes were applied for a second time about 90 minutes later. He began to respond to stimuli and was fully conscious the next day. After 1 week he developed acute excitement, restlessness, and mania. After 2 months, he was discharged with persistent athetoid movements of the hands and personality changes. *Comment: A case of severe poisoning with delayed treatment. CNS symptoms not fully reversed.*

Case 3: A male patient was accidentally poisoned with cyanide. He received SN and STS and reportedly recovered. *Comment: The patient's recovery was attributed to the medication. This case is inconclusive as not enough data about the condition of the patient were available.*

Case 4: A customs inspector lost consciousness about 1 hour after he had stayed in a room with 460 sacks of mung beans that had been HCN fumigated the previous day. Immediately after he had been given AN pearls for inhalation he recovered but felt weak for one month. *Comment: A case of moderate poisoning with successful treatment with AN.*

Case 5: A 64 year old man was overcome by HCN fumes and was found unconscious, cyanotic and pulseless. He received artificial respiration and AN, resulting in

‘hypermotor activity’. One hour later he received 0.3 g of SN and 12.5 g of STS i.v. AN inhalation was continued together with oxygen. He became conscious about 2 hours after exposure. The patient was sent home 4 hours later. *Comment: A case of severe poisoning in which quick application of AN, but delayed application of SN/STS were effective.*

Case 6: A 22 year old female was exposed to cyanogen chloride by inhalation. She was found unconscious and flaccid, followed by restlessness and hypermotor activity. 15 to 20 pearls of AN and 300 mg of SN were simultaneously administered. STS was not given (technical difficulties). Epinephrine was injected subcutaneously due to low blood pressure. 3 hours later she regained her rationality and was discharged with complaints of persistent abdominal, chest and back muscle tenderness the next day. *Comment: A case of moderate poisoning in which AN and SN were effective after some delay.*

Case 7: A 61 year old man was overcome by HCN. He was found unconscious, cyanotic, and in cardiorespiratory arrest. 20 minutes later he was treated with AN by inhalation, artificial respiration, and additionally with nikethamide intramuscularly. When he regained consciousness approximately 45 minutes later 0.3 g of SN and 12.5 g of STS were administered to him, along with epinephrine, caffeine, and sodium benzoate. He was taken home 2 hours after exposure. *Comment: A case of severe poisoning. From the description it is not fully clear if the patient regained consciousness before the administration of SN. If so AN was effective in a severe poisoning case.*

Case 8: A 67 year old man was overcome by HCN. He was found unconscious, cyanotic, and in cardiorespiratory arrest. The man was unconscious several minutes after exposure to hydrocyanic acid. He was pale and his pulse was weak and irregular. Respiration was normal. He responded well to AN only. *Comment: A case of moderate poisoning in which AN was effective.*

Case 9: A man, aged 44, was accidentally exposed to HCN gas. He became unconscious for a few minutes. AN was administered from several pearls, one at a time. He was nauseated, vomited several times, and developed a severe headache. He was sent home in an ambulance. *Comment: This case is inconclusive as the moderately poisoned patient may have regained consciousness before he got treatment. The headache was probably due to the AN. Specific treatment was not really appropriate in this case. However, AN was tolerated well.*

Case 10: A man was exposed to HCN from a cake of precipitate. He became dizzy and vomited. This continued for approximately 2 hours, at which time he arrived at the first-aid station. He was immediately given 0.3 g of SN and 12.5 g of STS. After 45 minutes the patient had recovered and was discharged 32 hours later. *Comment: This is a case of mild poisoning. SN/STS would not be conclusive at all as there was no indication that nitrite treatment had been required.*

Summary of the cases presented by Chen and Rose: Assessment was difficult as symptoms were often insufficiently described: 10 mild, 2 moderate and 2 severe intoxications. Four out of 10 patients received AN alone; the remaining 6 received the nitrite antidote kit. The moderately and severely poisoned patients all received SN, 3 in combination with STS, and 1 with AN. They all recovered completely.

Cheok, 1978

A report from Sarawak describes two children having eaten a bitter cassava cake – so bitter that the parents did not eat more than a bite. The cake was later found to contain 3 ppm cyanide (as CN⁻). Both children (1.5 and 2.5 years of age) fell ill 7 and 7.5 hours after eating the cake with vomiting, drowsiness, and weakness, in one case also cyanosis. As no SN, which the authors recommend for treatment followed by STS, was available, the children were treated with STS alone (50 ml of 25% solution, 2.5-5 ml/min i.v.) and one with oxygen via nasal catheter. Both children recovered quickly and fully. *Comment: No measurements of cyanide in blood or for thiocyanate are available; in both cases gastric aspirate was negative for cyanide. The clinical improvement after antidote application is not conclusive in the absence of measurements cyanide detection.*

Cherian and Richmond, 2000

A fatal accident occurred aboard a stern freeze trawler 40 miles off the coast of West Africa. The warm temperature and failure to clean the storage tank immediately after its previous use had left a mixture of sea water and fish to rot. After opening the door of this salt water tank to flush it out, the crew member collapsed immediately, apparently having been overcome by toxic fumes. Unaware of the danger several crew members went to help him and were also overcome. 3 men died within 1 minute and 6 suffered other injuries, such as blackouts, seizures, chest pain, vomiting and difficulty in breathing. Several volatile agents were detected in the blood and whole lung. Besides cyanide (blood level <50 µg/l) and MetHb (14%) methane could also be identified. *Comment: A case of severe poisoning although involvement of cyanide is doubtful, the use of antidotes is not described.*

Chin and Calderon, 2000

A 19 year old woman drunk from an open soda bottle contaminated with cyanide (cyanide could be detected) by her boyfriend who had been working in a jewellery business for some time. He called EMS and on their arrival she was apnoeic and unresponsive. She was intubated and taken to hospital, unresponsive (GCS = 5); apnoea, tachycardia of 110/min, hypotonia of 90 mm Hg systolic blood pressure and a blood-pH of 7.01 and with 10 mmol/l lactic acid were observed.

She received naloxone, thiamine, glucose and i.v. fluids, together with activated charcoal and sodium bicarbonate. Thirty minutes after administration of 12.5 g of STS, the pH increased to 7.17 and the MetHb counted for 9.5% (no MetHb forming drug had been administered until that time; the TF believes that the blood samples had been mixed up). 300 mg of SN were administered and half the initial dose of both antidotes was repeated 60 minutes later “to avoid excessively high levels of methaemoglobinaemia yet continue to treat the patient’s acidosis”. The patient made a good recovery and could be extubated after 5 days. 8 days after the incident she was discharged. 5 months after the intoxication she complained about short-term memory problems as well as a weakness in the lower legs. *Comment: A case of moderate intoxication in which the victim could be rescued with the nitrite-kit but neurological and motor sequelae persisted. The elevated MetHb level before SN remains unexplained.*

Cliff and Coutinho, 1995

One 3 year old child died, probably from aspiration pneumonia. Cyanide poisoning was diagnosed by the history of eating cassava and the similarity of the clinical picture with acute cyanide intoxication. *Comment: No measurements at all were done.*

Coentrão et al., 2010a (Coentrão and Moura, 2010b)

A 50-year-old goldsmith who tried suicide by ingestion of a jewellery cleaner solution containing approximately 1.2 g of potassium cyanide presented unconsciousness, with severe lactic acidosis and arteriolisation of venous blood gases. Following hydroxocobalamin treatment, neurologic and metabolic disorders rapidly improved. He was discharged home 4 days later, without neurological sequelae.

Coentrão and Moura, 2011

A retrospective chart review of all admissions for acute cyanide poisoning by ingestion for the years 1988 to 2008 was conducted in a tertiary university hospital serving the largest population in the country working in jewelry and textile facilities. Of the 9 patients admitted to hospital during the study period, 8 (7 males, 1 female; age 36 ± 11 years, mean \pm SD) attempted suicide by ingestion of potassium cyanide used in their profession as goldsmiths or textile industry workers. 5 patients had severe neurologic impairment and severe metabolic acidosis (pH 7.02 ± 0.08 , mean \pm SD) with high anion gap (23 ± 4 mmol/L, mean \pm SD). Of the 5 severely intoxicated patients, 3 received antidote therapy (sodium Thiosulphate or hydroxocobalamin) and resumed full consciousness in less than 8 hours. All the patients survived without major sequelae. *Comment by the authors: Metabolic acidosis with high anion is a good surrogated marker of severe cyanide poisoning. STS and HOCO are both safe and effective antidotes.*

Dall and Hannah, 1964

As examples, Dall and Hannah (1964) reported on “Oxygen therapy in cyanide poisoning” in a 39-year-old male accidentally poisoned by oral ingestion of a solution of KCN. After having ‘called for medical attention’ he was found in deep coma, hypopnoea, seizures, respiratory and cardiac arrest. In addition to oxygen, reported treatment included the administration of a number of ampules of AN, external cardiac massage, artificial ventilation after endotracheal intubation, STS (25 g) and SN (300 mg). The patient fully recovered. However, the causal relationship between oxygen administration and survival is complicated by the associated treatment, the efficacy of which might have been underestimated. Additionally, Bismuth et al (1984) emphasised about the priority of oxygen in cyanide poisoning and reported details of treatment in 4 out of 25 cyanide poisoning. One of the four with minor poisoning recovered with oxygen alone. Among the three remainders two died while one survived without sequelae after having received in addition to oxygen and supportive treatment, including cardiopulmonary resuscitation for transient cardiac arrest, Co-EDTA, and HOCO. *Comment: It seems rather difficult to draw any definitive conclusion from these complex and severe cases receiving multiple combined treatments.*

Daunderer et al, 1974

A 26 year old female working in a chemical laboratory tried to commit suicide by drinking about 10 g of KCN solved in sour vine. She vomited and then collapsed directly after having drunk the solution. In the hospital she presented the following symptoms: good general condition, irregular breathing, mild peripheral cyanosis, red shining parts of the skin, odour of bitter almonds, deep coma, no reaction to painful stimulus, wide pupils, minimal reaction to light, blood pressure 140/90 mm Hg, HF 130/min and weak pulse. She was intubated, ventilated and gastric lavage was performed. 45 minutes after ingesting the poison she was administered 250 mg of DMAP together with 300 mg of Cobalt-EDTA i.v. No adverse effects were seen and a short time after this therapy the pupils became reactive to light again and deep tendon reflexes as well as reaction to pain could be triggered. About 45 minutes after that treatment 60 ml of a 10% STS-solution were applied (6 g). During this infusion the patient became awake and orientated. In the former course metabolic acidosis, hypotonia, tachycardia and a general oedema occurred. All these side effects could be managed within a few days. Methaemoglobin was not measured. *Comment: A case of severe poisoning in which the combination of DMAP, Cobalt-EDTA (both given in the recommended doses) and STS worked well. The generalised oedema can be correlated to the Cobalt-EDTA. Fortunately all adverse effects and poisoning consequences could be managed.*

Davies, 1975

A 40 year old male received SNP while anaesthetized during surgery. The blood pressure could not be lowered sufficiently, so the SNP dose was increased. Approximately 400 mg had been given in total. After cessation of the infusion the blood pressure dropped, and the skin was pink. Bradycardia necessitated administration of atropine but blood pressure dropped further. Refractory cardiac arrest followed. The level of cyanide in blood was 5 mg/l.

Davis and Ewer, 1988

A 22 year old electrician drank a solution containing cyanide to commit suicide. A short time later he told his father. On admission to the ED 45 minutes later he presented a GCS of 5 points, dilated and fixed pupils, a hypertonia (blood pressure 160/80 mm Hg), a hyperreflexia, tachypnoea, tachycardia, a flush and facial oedema (*note: this was prior to being given the cobalt-EDTA*). There was no reaction to painful stimuli. 2 ampoules of AN were administered by inhalation together with 100% oxygen. Then he received 300 mg of Co-EDTA together with 50 ml of a 50% glucose solution, 0.3 g of SN and 12.5 g of STS. Furthermore gastric lavage was performed and activated charcoal administered. As his condition did not change during the next hour, the cobalt-EDTA-treatment was repeated and hyperbaric oxygen initiated. At this time the patient was still unconscious with loss of reaction to pain, but the pupils were now reactive to light and he made spontaneous movements. After the second dose of Co-EDTA lactic acidosis he quickly recovered and was orientated within 20 minutes. He vomited and was given sedatives as he had become agitated. Furthermore, facial oedema was detected (*Note: this was already mentioned with the initial symptoms, one does not know, when it really occurred*). He finally recovered completely. *Comment: This is a case of severe poisoning. No blood-cyanide-levels were measured. The Co-EDTA was given in high dose, as the first antidote and the patient made a remarkable recovery only after the second cobalt-dose, but the authors attribute the good course to the hyperbaric oxygen (which was initiated at the same time as the second antidote application).*

Dawood, 1969

According to a report from Singapore a 3.5 year old girl had started to vomit ten hours after ingestion of cassava and arrived in hospital in a state of shock with acidosis, drowsiness, irritability, shortness of breath and paleness. The eyes were staring, the pupils were dilated and non-reactive to light, blood pressure was hypotonic, and reflexes were missing. Neither cyanide nor thiocyanate could not be detected in the gastric aspirate. The tapioca contained 94 mg/kg hydrogen cyanide. The child recovered without specific treatment, no cyanide or thiocyanate measurements were done. *Comment: Cyanide poisoning was only suspected, and no antidotes were given.*

De Busk and Seidel, 1969

Case 1: A 22 year old student in chemical engineering took 1.5 g of NaCN in a suicide attempt. He presented a flush, arrhythmia and was unresponsive to stimulation. As the vomit smelt like bitter almonds and a suicide note was found, cyanide intoxication was quickly suspected. The ECG was altered, consisting of a short QT time, an increased ST-line and a loss of the P-wave. The cyanide blood level or other blood-parameters were not determined. Primarily intubation and ventilation as well as gastric lavage were performed and AN pearls were crushed over the respirator. Afterwards he was administered 300 mg of SN (10 ml of a 3% solution). The estimated interval from cyanide ingestion to i.v. SN administration was 20 minutes. Cardiac arrest occurred but could be reversed with CPR. Then 12.5 g of STS (50 ml of a 25% solution) were injected resulting in spontaneous movements and vital parameters being re-established. As the mental state did not improve a course of repeat AN/STS treatment was administered resulting the cyanosis clearing completely and the patient fully recovering. *Comment: A case of severe poisoning in which the patient recovered from a high dose of cyanide even though he had to be resuscitated. Resuscitation was required after the first nitrite dose, and only STS and a repetition of AN/STS led to full recovery.*

Case 2: A laboratory technician presented respiratory distress, followed by collapse and a possible seizure during the half-hour preceding admission. On arrival in hospital she was comatose, the extremities were flaccid and deep tendon reflexes were absent. No blood pressure was obtainable and the femoral pulse was weak with a pulse rate of 110/min. Respiration was irregular at 6-10 breaths per minute. The pupils were dilated and unresponsive to light. The ECG showed atrial fibrillation, occasional ectopic ventricular beats with T waves originating high on the R wave. Treatment included intubation, ventilation with 100% oxygen, gastric lavage and metaraminol (probably a catecholamine) i.v. for circulatory support. Because of a smell of almonds acute cyanide poisoning was suspected and 0.3 g of SN was given over four minutes and 17.5 g of STS (both agents were taken from the Lilly cyanide Kit). The patient became moderately cyanotic (MetHb). After 15 minutes she had regained consciousness and could be asked about what had happened. She said she had taken 4-6 g of KCN in a suicidal attempt. The cyanide came from her laboratory. The approximate time between ingestion of cyanide and treatment with SN was 50 minutes. Recovery was uneventful with prompt reversal of the ECG changes. No cyanide or MetHb measurements were taken. *Comment: This case is conclusive as regards the prompt improvement after nitrite treatment that followed the respiratory and circulatory insufficiency caused by a high cyanide dose.*

Dequidt et al, 1974

Case 1: A 19 year old man was accidentally exposed to acetonitrile vapours. After 4 hours GI symptoms developed with convulsive coma accompanied respiratory pauses by 12 hours

followed by collapse and cardiac arrest by 24 and 26 hours respectively. From the time of collapse he had been intubated and had received gastric lavage. He survived the cardiac arrest with symptomatic treatment but with persistence of an unstable blood pressure and repeated convulsions. 41 hours after the exposure, cyanide poisoning was confirmed by a free blood cyanide concentration of 1.12 µg/ml. He received 600 mg of Co-EDTA. On the following day he remained in a profound coma. About 65 hours after the exposure, reactivity to painful stimuli reappeared and the patient then received 4 g of HOCO. The patient was subsequently diagnosed as brain dead. *Comment: Antidotal treatment of this severe poisoning was delayed for more than 24 hours so Co-EDTA or HOCO efficacy cannot be assessed.*

Case 2: There is a second case report of death due to inhalation in spite of administration, after more than 24 hours, of Co-EDTA (2x600 mg) and HOCO (4 g) in a worker after he had cleaned with acetonitrile and hot water. The first symptoms developed after about 12 hours. 24 hours after admission to hospital the blood cyanide level was 11.2 mg/l and the maximum level of acetonitrile in blood was 311 mg/l. *Comment: Again, antidotal treatment of this severe poisoning was delayed for more than 24 hours so Co-EDTA or HOCO efficacy cannot be assessed.*

DiNapoli et al, 1989

A 29 year old man was found unresponsive a few minutes after self-injecting undetermined amounts of KCN and NaAs₂O₃ (arsenite) i.v. in a suicide attempt. On arrival at ED he was unconscious and unresponsive to stimuli. He was tachycardic with deep gasping respiration. He was given naloxone, dextrose and thiamine but his condition did not change. AN pearls were administered followed by a return of the spontaneous movements. In the blood gases a pH of 7.12 and a pO₂ of 54 mm Hg were detected. After infusion of 0.3 g of SN and 12.5 g of STS the patient regained consciousness within seconds and the blood pH increased to 7.28 and the pO₂ to 85 mm Hg. The MetHb level was 7.5%. The anion gap counted 24 mmol/l and the cyanide level 4.4 mg/l before antidotal treatment. The cyanide level fell to 0.18 mg/l 12 hours after the nitrite had been administered. After recovering from coma the patient had visual problems and was treated empirically for methanol poisoning, which was subsequently not confirmed. Blood and urine arsenic concentrations were also quite elevated, so the patient received BAL and later D-penicillamine. He was ultimately released from the hospital about 3 weeks later. *Comment: Though the arsenite level was not increased the man received specific treatment for arsenite intoxication (chelators). None of the antidotes, neither for arsenite nor for cyanide intoxication, seems to have caused adverse effects. In the end he made an uneventful and complete recovery after effective treatment of the cyanide poisoning.*

Dodds and McKnight, 1985

A 43 years old worker was completely immersed in a bath of hot copper cyanide for 3 minutes by accident. Only 15 minutes later he lost consciousness, became cyanotic, his respiration was irregular and the blood pH counted only 6.9. The base excess was 20 mmol/l. He was ventilated, the clothes were removed and 300 mg of Co-EDTA administered over 3 minutes. As this did not show a prompt improvement, the antidote administration was repeated. He developed massive facial oedema which commenced in the peri-orbital area. As his condition still did not improve, the nitrite antidote kit was additionally administered (AN pearls, SN and STS). It took 16 hours before he regained consciousness and recovered completely. *Comment: The antidotal therapy consisting of Co-EDTA (twice) and nitrites did not result in fast recovery but in the end the treatment seemed to have been efficient (however, as 16 hours had passed the recovery could also be attributed to physiological detoxification because normally the antidotes should have worked more quickly).*

Edwards and Thomas 1978

This is primarily a comment to a publication from Graham et al, Arch Intern Med 1977, 137:1042 (see also **Bryson, 1978**), but they describe an additional case.

A 48 year old chemist was presented deeply unconscious with unrecordable blood pressure; he had a cardiorespiratory arrest on arrival. He was asystolic and was resuscitated by a combination of intubation and ventilation with 100% oxygen, cardiac massage, and intracardiac injection of 1 mg of adrenaline. Direct-current (DC) shock was used to revert ventricular tachycardia to atrial fibrillation and 10 mg of practolol i.v. to control rapid ventricular rate. He remained hypotensive (60 mm Hg systolic, central venous pressure zero) and his arterial pH was 6.92. Serum *Comment: There is little to mindicate that this was cyanide poisoning other than he was a chemist.*

Eiró Gonsalves et al, 1956

6 cases of cyanide intoxication as a result of ingestion of wild cassava were admitted to the emergency ward of the pediatric clinic during a 5 month period.

Case 1: A six year old boy became unconscious about 2½ hours after the meal. He survived after i.v. application of a total of 90 ml of STS 2% and 50 ml of SN (concentration not reported).

Case 2: Another six year old boy had abdominal pain and was vomiting 5 hours after a meal and became somnolent. He recovered after i.v. administration of 2 ampoules of SN 2% and 2 ampoules of STS 10% (amount not reported).

Case 3: A five year old girl became somnolent and started vomiting 8 hours after a meal. She, too, recovered after 2 ml of SN 2% and 20 ml of STS 10%.

Case 4: A nine year old boy presented with initial tetanic convulsions of the extremities, followed by flaccid paralysis and inability to speak. He recovered after 15 ml of SN and 15 ml of STS (concentration not reported). The time between ingestion and treatment was not reported.

Case 5: A six year old boy was vomiting and feeling weak four hours after a meal. He inhaled an ampoule of AN and then received 5 ml of SN 10% and 60 ml of STS 10%, after which he recovered.

Case 6: After the same meal a four year old girl also was feeling weak and vomited. She too recovered after 5 ml SN of 10% and 30 ml of STS 10%. *Comment: No cyanide or thiocyanate measurements were performed, cyanide intoxication can only be assumed due to the symptoms observed, which are more or less indicative only, the effect of the antidotal treatment is not clearly described – all children had recovered on the following day, which may or may not be due to the antidotes given.*

Espinoza et al, 1992

8 male children between 8 and 11 years of age developed vomiting and excessive weakness after ingesting bitter cassava. On arrival at ER they presented respiratory failure, bradycardia, hypotension and cardiovascular collapse. Two children developed generalised seizures with relaxation of sphincters. All received 100% oxygen. The more serious half of the children were treated with 3% SN (0.2ml/kg bw) and 25% STS (0.04ml/kg). The less serious 4 were treated with 500mg hydroxocobalamin in dextrose solution. All 8 survived without sequelae.

Favarel-Garrigues, 1982 / Dreykluft et al, 2004 (case 1)

Case 1: A 36 year old chemical-technical assistant was presented to the hospital in a coma with tachypnoea (32/min), tachycardia (100/min) and wide, non-reactive pupils. The P_aCO_2 was only 19 mm Hg and he showed a severe acidosis (pH 6.96, lactate 20 mmol/l, anion gap 25 mmol/l). The patient had a history of multiple suicide attempts. His treatment included gastric lavage application of activated charcoal and sodium bicarbonate, because of the unknown poison. 9 hours after the incident the man became responsive again and recovered completely. He then

confirmed the suicide attempt with KCN. *Comment: No specific antidote was given. Documentation a bit confusing because the publication was part of a quiz.*

Case 2: A 27 year old woman ingested 2 g of KCN in a suicide attempt. 30 minutes later the ambulance staff found her comatose, hypotonic with dilated pupils, failed respiration and convulsing. She was ventilated with 100% oxygen and given drugs for haemodynamic support (it has not been mentioned which drugs). In the ED an unspecified amount of STS was administered together with sedatives. This resulted in intermittent improvement of the haemodynamic situation but 5 days later she died due of cerebral damage. *Comment: Information is incomplete although it would appear that in this case of severe poisoning STS alone was eventually not effective.*

Feihl et al, 1982

After ingesting 600 mg of KCN a 60 year old chemist arrived in hospital comatose and bradypnoic with severe metabolic acidosis. In addition to supportive care, he received SN and STS and recovered without sequelae ‘despite excessive methaemoglobinaemia’ (value not provided). The authors note an extraordinary blood cyanide concentration of 40 mg/l including a fraction bound to inactive cyanMetHb. *Comment: The actual free cyanide level is unclear. However, in this case of moderate poisoning SN/STS was effective.*

Feldmann, 1990

A 28 year old man was presented at the emergency department after ingesting 800 mg of KCN in a suicide attempt. He survived only with intensive medical and psychiatric intervention, and went on to develop severe Parkinsonian symptoms, including profound micrographia and hypersalivation. Magnetic resonance imaging showed bilateral and symmetrical basal ganglial abnormalities. Survival following cyanide (*TF insert: severe*) poisoning is rare; the clinical, radiologic, and neuropathologic sequelae are reviewed in other documented cases.

Fernández et al, 2008

A 38 year old man was exposed by inhalation and dermal to pentenenitrile solvent solution at work. Several hours afterwards he developed progressive weakness prompting him to decontaminate in a shower for 20 minutes. He became syncopal. EMS found him in atrial fibrillation at a rate of 112 bpm that increased to 177 upon arrival at hospital where he presented headache, nausea, vomiting and metabolic acidosis. He was treated with 100% oxygen,

ondansetron and AN followed by SN and STS. He had a confirmed serum thiocyanate level of 7.5 µg/ml and recovered without sequelae.

Finelli, 1981

A 30 year old man was found unconscious, apnoeic, and cyanotic after he had ingested an unknown quantity of a cyanide-containing insecticide. After resuscitation and the application of STS, he regained consciousness. He remained mute for 12 days and developed choreiforme (involuntary spasmodic) movements. After 16 days he was normal apart from mild dysarthria and minor movement disorders. A computer tomography revealed a bilateral symmetrical infarction of the globus pallidus and an infarction of the left cerebellar hemisphere. *Comment: In this case of severe poisoning neurological sequelae persisted after effective treatment with STS, but these were probably due to cardiorespiratory arrest.*

Fortin et al, 2007

A 47 year old male with third-degree burns and (later) proven cyanide poisoning from smoke inhalation was found in cardiac arrest. He was resuscitated and received 10 g of HOCO in two stages. After the first 5 g of HOCO the cyanide level in blood was 3.4 mg/l and after the second 5 g it was below 1 mg/l. While cardiac activity resumed he had to be declared brain dead on day 3. *Comment: A severe poisoning case in which HOCO improved the cardiac, but not the neurological situation after cardiac arrest.*

Froldi et al, 2001

This was a rare case of suicide by oral ingestion of 6 g of SNP in a 41 year old woman found dead. At autopsy total cyanide in blood was 5 mg/l. *Comment: Fatality, no antidote use.*

Gaillard, 1991

After a domestic dispute with his wife a 33 year old man tried to commit suicide with a solution containing cyanide. The ambulance staff found a semi-comatose, agitated, tachycardiac (heart rate 135/min), tachypnoeic, dizzy patient with a bilateral miosis (not mydriasis). They administered i.v. fluids and initiated ventilation with 100% oxygen by face-mask. At admission at the ED the patient was fully conscious with spontaneous respiration but after developing seizures had to be treated with clonazepam and intubated. Ventilation and sedation were performed. Gastric emptying was undertaken and activated charcoal administered. Additionally,

sodium bicarbonate and 4 g of HOCO were given about 2 hours after poison intake. A cyanide level of 150 $\mu\text{mol/l}$ (4.05 mg/l) was measured at admission and decreased to 15 $\mu\text{mol/l}$ (0.41 mg/l) after treatment. The HOCO-treatment was repeated twice within the next three days. The man was extubated on the fourth day in hospital and discharged on the seventh day. He recovered completely. *Comment: The clinical course is not well described, but obviously HOCO significantly decreased the cyanide blood level.*

Gambaro et al, 2007

Case 1: A 26 year old woman was taken to hospital presenting slowed respiration, gasping breath and bradycardia. She died more than 1 hour later. Symptomatology suggested cyanide poisoning, which remained unconfirmed.

Case 2: A 30 year old man was found dead his car. Laboratory analysis of the liquid in a bottle found nearby revealed cyanide. Forensic evaluation could not establish the time interval between ingestion and death. The analysis of the stomach content and postmortem blood samples was positive for cyanide.

Comment: cases of severe poisoning with fatal outcome but no antiotes were used.

Gaultier and Le Breton, 1952

A male in custody at a police station developed hand ataxia, followed by trembling. 30 minutes later convulsions, somnolence/coma and irregular respiration were observed, followed by respiratory arrest and death 3 hours after the start of the ataxia. Autopsy revealed cyanide poisoning. A tube with cyanide was found in the duodenum. *Comment: Cyanide poisoning was not recognised and no antidote was administered.*

Geller et al, 1991

A patient received gastric lavage and activated charcoal administration 30 minutes after ingesting acetonitrile. Hours later he developed mental status abnormalities. He vomited prior to a generalised seizure. The blood cyanide level documented a significant exposure (intoxication). Following the administration of STS, the patient made an uneventful recovery. *Comment: A case of moderate poisoning with apparent delayed effects in which STS was effective.*

Ghiringhelli, 1955

One case from Italy has been reported – a young man drank by mistake a few mls of adipic nitrile. Symptoms were weakness, headache, vertigo, nausea/vomiting, cyanosis, tachypnea, hypotension, mydriasis, somnolence, tonic-clonic convulsion, inability to stand independently. All these quickly resolved after application of STS. *Comment: a case of nitrile poisoning in which STS was completely effective.*

Glenn et al, 1994

A 54 year old man deliberately drank a potassium-gold cyanide solution (300 ml) that contained about 1.65 g of KCN and survived after treatment with the Lilly antidote kit and hyperbaric oxygen. When he arrived in hospital 6½ hours later vomiting with diarrhoea and was gradually disorientated. He was given oxygen and received antidotal treatment from the Lilly-kit. There was no proper response and the SN and STS treatment was repeated. Despite these measures, he became progressively obtund, with hypotension refractory to pressors. 8 hours after ingestion, the blood cyanide level was 0.09 mg/l, i.e. within normal limits. *Comment: The case is not conclusive. It seems that cyanide poisoning was not the main problem, as the patient was alive 6½ hours after ingestion and the cyanide level in blood was very low. The whole case probably describes gold salt poisoning.*

Goodhart, 1994

A 54 year old man working in a watch factory intentionally drank 300 ml of a gold-solution containing 1650 mg of KCN. 6 hours and a half later he was presented in the emergency centre with vomiting, diarrhoea and an altered mental status. The blood cyanides were measured eight hours after poisoning with 0.09 mg/l. The vital signs included a systolic blood pressure of 50 mm Hg and a heart rate of 118/min. He was given 100% oxygen by face mask, activated charcoal and sympathomimetics. Under this treatment he became comatose, cyanotic, anuric and apnoeic. The nitrite-antidote-kit (AN pearls, 0.3 g of SN and 12.5 g of STS i.v.) was administered and since the patient did not respond to the first dose this was repeated. Additionally six cycles of hyperbaric oxygen were performed. The blood cyanide level remained below the measurable range and the lactate only rose to 3.4 mmol/l. He stayed with on respirator for 18 days, developed ARDS, pneumonia, jaundice and had to be temporarily dialysed. Despite all this he could be discharged 41 days later in complete remission. *Comment: The clinical presentation was serious, but the blood-cyanide levels were quite low. Probably it was a gold intoxication.*

Graham et al, 1977

A young man (aged 21) ingested KCN with suicidal intent. He became cyanotic and lost consciousness. The vital signs included a respiration frequency of 24/min, heart rate of 68/min (arrhythmic) and blood pressure of 168/112 mm Hg. Both pupils were reactive to light. In the blood samples an anion gap of 35 mmol/l together with a mild acidosis (pH 7.27 under 5 l oxygen per minute) could be detected. Furosemide was administered for the treatment of pulmonary oedema (noticed in the chest X-Ray). Further therapy consisted of sodium bicarbonate and sodium-chloride i.v. as well as intubation and ventilation. 10 hours later the suspicion of cyanide intoxication was substantiated (the authors do not report how, but after recovery the patient confirmed that he had taken 200 mg KCN). Since so much time had passed between the intake of the cyanide and the diagnosis of intoxication no specific antidote was applied. 12 hours after the suicide-attempt the blood-cyanide-level was measured with 2.0 mg/l and fell down to 1.6 mg/l after 22 hours and 1.2 mg/l after 84 hours respectively. 24 hours after the incident the young man could be extubated and was discharged on the ninth day post exposition. *Comment: A case of moderate poisoning with supportive treatment only. Even if the time span between intoxication and diagnosis was long, a specific antidote that supports the excretion of the poison (e.g. STS) might have accelerated the recovery.*

Grunske, 1949

In 1949, the death of a child was reported following pest control operations with CAN. Two more similar cases have not been published. The child had lacrimation, nausea, tonic seizure 1 hour after exposure in a room ventilated after pest control operator treatment. The next morning the child was in respiratory distress and coma with tachycardia and pallor. She died on transport to hospital. *Comment: A case of sever poisoning but no antidote had been given and cyanide measurement was not done.*

Gunder et al, 1969

A 3.5 year-old Israeli year old girl ingested an unknown amount of apricot pits and developed later pallor, fast pulse, rapid breathing, dilated pupils, vomiting and temporary unconsciousness. A bitter almond smell was noticed. Gastric lavage and paraffin application prompted full recovery. *Comment: Cyanide poisoning after amygdalin intake can be supposedly based on symptoms. No specific treatment was done.*

Hall et al, 1986

A four year old boy with Down syndrome and an eye seizure disorder ingested 12 Laetrile tablets (equivalent to 300 mg amygdalin) after a meal of fruits, vegetables and nuts. Within 1.5 hours he slowly became unresponsive and had multiple seizure episodes. The child was transported from the local hospital to a referral hospital but in both hospitals no complete antidote set (Lilly-Kit) was available. The child was presented in the second hospital with bradycardia (60/min) and thus measurable blood pressure. He was intubated and ventilated. Gastric lavage was performed and 15 g of activated charcoal was instilled. The blood gases showed severe metabolic acidosis with ineffective respiratory compensation, pH: 6.85, pCO₂: 15 mm Hg, pO₂: 169 mm Hg, anion gap: 16 E/l. Initially only AN pearls were administered by intermittent inhalation. This increased the blood pressure to 100/50 mm Hg. The blood pH could only be improved a little by sodium bicarbonate. The full antidote-kit (probably Lilly-Kit) was available 6 hours post ingestion. During the last 2 hours the child was maintained on AN (the dose was not specified). Bradycardia and hypotension improved during periods of inhalation and worsened during periods of noninhalation. The antidote-kit, 5 ml of SN 3% (150 mg) and 25 ml of 25% STS (6.25 g) were sequentially administered. Within 30 minutes the child recovered with normal blood pressure seizures. The child was extubated 15 hours postingestion and was alert and awake. The blood gases improved rapidly and were normalised 15 hours after ingestion. The highest whole blood cyanide level was 16.3 mg/l after the antidote administration and 8.2 mg/l at admission to the second hospital. *Comment: A case of severe cyanide poisoning after Laetrile ingestion. AN was partially effective, but allowed to bridge the time until availability of SN/STS which was fully effective.*

Hall et al, 1987

A man (34 years old) took 1 g of KCN in a suicide attempt. He reported the incident after about half an hour and became lethargic. A further 15 minutes later he lost consciousness and was ventilated with an anaesthesia bag by the ambulance staff. Upon arrival at the ED, 1 hour after the intoxication, he presented apnoeic so was intubated and ventilated with 100% oxygen. Sodium bicarbonate was administered to correct an acidosis (pH 7.11 to 7.33). His pupils were fixed and dilated. Quarter of an hour after his admission 300 mg of SN and 12.5 g of STS were administered over 45 minutes. As this was finished the patient showed spontaneous respiration and movements and the pupils were again reactive to light. 4 hours after the suicide attempt the man was alert and was extubated 6 hours later. The cyanide levels in the blood samples were determined at 1, 1.5, 2, 2.5, 3, 5, 11 and 17 hours after the exposure as 5.5, 8.72, 15.65 (extremely high value!), 9.93, 1.74, 9.82 (*no explanation can be given for this new increase, after the blood cyanide had already fallen*), 0.62 and 0.53 mg/l. A level of 2% MetHb was measured after the application of the antidotes (*this is below the level that is generally considered as therapeutically effective*). The patient fully recovered. *Comment: This is a case of severe*

poisoning as confirmed by high cyanide blood levels with respiratory arrest in which quite high doses of SN/STS were fully effective, albeit MetHb levels remained low.

Hall et al, 1989a

After smoke inhalation a 78 year old patient was found comatose, apnoeic and asystolic. He could be resuscitated on site. In hospital he was treated with 300 mg of SN i.v., which induced severe hypotension. MetHb elevated very little. 4½ hours after hospital admission the patient was treated with hyperbaric oxygenation (hyperbaric oxygen-chamber) without success. The patient died the next day in irreversible circulatory failure. The cyanide level in blood was 0.34 mg/l on admission. *Comment: This was no significant cyanide poisoning as shown by the blood level. In the absence of therapeutical benefit the side effect of SN (hypotension) prevailed.*

Hall et al, 2009

In a review article, Hall et al (2009b) quoted a case report from Person (1993). Two children were found comatose in a fire. They received oxygen by inhalation and 400 mg/kg bw of STS and were transferred for hyperbaric oxygen treatment. Both children recovered completely. The treatment was well tolerated. The cyanide levels were 1.15 mg/l in one child and 1.1 mg/l in the other. *Comment: Two cases of moderate poisoning that reacted well to STS alone, while CO poisoning was probably concurrently treated with hyperbaric oxygen.*

Hantson, 1996

A woman, aged 38, ingested one teaspoon of potassium ferrocyanide ($K_3Fe(CN)_6$) in a suicide attempt. Half an hour later she arrived in hospital fully conscious, tachycardiac (heart rate 110/min) and hypertonic (blood pressure 150/80 mm Hg). The physician on duty suspected cyanide poisoning and performed intubation, ventilation and application of 5 g of HOCO i.v. 1 hour after the incident the patient was still conscious, the vital signs were all within the normal ranges and a cyanosis could not be detected. She also had not developed adverse effects because of the antidote. The blood gases only showed an elevated anion gap of 13 mmol/l. 12 hours after the intoxication the patient was extubated and could be discharged soon after. During her time as an in-patient the cyanide level was determined after 1, 3, 4, 6 and 8 hours as 20 (0.54), 15.4 (0.42), 12.3 (0.33), 8.85 (0.24) and 3.6 (0.1) $\mu\text{mol/l}$ (mg/l). MetHb, pH or the lactate were altered at any time. *Comment: As potassium ferrocyanide does not liberate cyanide, this was no cyanide poisoning. The antidote was not justified, but did not show any side effects.*

Harry, 1985

A 54 year old female laboratory assistant ingested 200mg of sodium cyanide. After 5 minutes she was confused and uncomfortable. 30 minutes later she was found comatose and unresponsive and was ventilated with oxygen. At 60 minutes she was resuscitated and was in hypertonic coma (class 3/15 on Glasgow score), temperature 33°C, blood pressure 70/40 mmHg, rate 120/min with lactic acidosis (29.5 mmol/l). Blood analysis showed vascular shock and collapse. At 20 minutes HOCO (4gms) and STS (8gms, Cyanokit) normalised systemic arteriolar resistance and perfusion with 500ml plasma resumed moderate capillary pressure. The antidotes partially ameliorated the neurological state (Glasgow score 9/15) and decreased the lactic acidosis. Administration of hyperbaric oxygen at 2.5 ATA normalised venous oxygen levels and neurological status. Haemodynamic analysis revealed a 48% reduction during the first 5 hours that was not improved by HOCO and STS administration without clinical or radiological signs of pulmonary oedema. Total cyanide levels were measured as being 10mg/l at 1 hr, and 1.2, 0.3 and 0.25 mg/l at 4, 6 and 20 hours respectively, with an apparent distribution volume of 0.4L/kg.

Heintz et al, 1990

A chemistry student (aged 23) ingested 1500 mg of KCN in a suicide attempt (he admitted it after recovering). He developed a seizure and lost consciousness. His respiration was fast and the blood gases demonstrated a lactic acidosis (pH 6.96, BE -28 mmol/l and lactate 22 mmol/l). The blood cyanide was determined 6 hours after the incident as 6 mg/l. Until then he had received supportive treatment only, since the poison was not known. The therapy consisted of gastric lavage, administration of activated charcoal and ascarthatic sodium-sulphate, ventilation with 100% oxygen and sodium bicarbonate. After the result of the blood cyanide level was available, he was given 1 g/hour STS over 24 hours (24 g in a whole). At the end of the antidote administration he was conscious and could be extubated. The pH normalised within 9 hours of therapy, the BE within 12 hours, the respiration parameters within 24 hours and the blood lactate level was within the normal range after 2 days. So the STS showed a good efficacy, despite of having been administered six hours after the intoxication. The patient was discharged on the 9th day post exposure. *Comment: A case of mild poisoning based upon the clinical presentation although blood cyanide levels were high. STS alone was effective when accompanied by supportive treatment.*

Herbert, 1979

In 1940, a 14 year old Indian school girl was hospitalised with stupor, convulsions, fibrillary muscular twitching, pale cyanosis, laboured respiration, fever, hypotension and tachycardia. She died within 24 hours. Autopsy revealed large amounts of choke cherries and pits in the

gastro-intestinal tract. Crushed pits fed to guinea pigs caused convulsions and death. *Comment: Cyanide measurements were not done, however, in regard of the symptoms and the animal experiments cyanide poisoning is probable in this case. Some further cases have only been mentioned in a critical overview on Laetrile as cancer cure.*

Hill, 1942

A young female swallowed a quantity of SNP in water, possibly 5 grams. Death seems to have occurred after about 1 hour. The post mortem status resembled cyanide poisoning. Cyanide was found in the stomach contents (0.013 g). *Comment: A case of severe poisoning without use of antidote.*

Hillman et al, 1974

A 68 year old man accidentally drank from a bottle filled with a sodium-cyanide solution. He collapsed 3 minutes later and was found by the GP 15 minutes later cyanotic with respiratory difficulties. 55 minutes after the incident the patient was brought to the emergency department with apnoea, a pinkish complexion and acrostic. Due to these signs and symptoms, cyanide intoxication was suspected and the man was treated with CPR, sodium-bicarbonate and 300 mg of Co-EDTA. In a blood-sample taken before the therapy was started, the cyanide level could be measured at 5.5 mg/l (results were only available later on). Immediately after the treatment the vital signs stabilised (pulse palpable with sinus-tachycardia, systolic blood-pressure of 60 mm Hg), but the apnoea persisted. The antidote was given twice again after 10 and 15 minutes. However, this did not result in a clinical improvement. Gastric lavage and mechanical ventilation were initiated 2 hours after the exposure. 30 minutes later the patient regained spontaneous breathing. After a further hour he started hyperventilation resulting in an alkalosis (pH 7.47), hypocapnia (pCO₂ 17 mm Hg) and an increased pO₂ (95 mm Hg). The systolic blood pressure decreased from 90 to 60 mm Hg and ectopic cardiac activity could be seen in the ECG. These symptoms led to a fourth administration of Co-EDTA followed by a further drop of the blood pressure and a reactive tachycardia. 10 hours after the intoxication the antidote was given a fifth time because of a new respiratory arrest. No reaction to the therapy could be observed so mechanical ventilation was initiated again. A sixth antidote-dose was injected 12 hours after the incident and after that 9 doses had been given, one every 30 minutes. Despite of all these efforts the patient died of cardiac arrest 44 hours after the intoxication. The cyanide levels were measured 4 times during the treatment and were always below 1.0 mg/l. *Comment: The blood cyanide level on admission was 5.50 mg/l which decreased to 0.3 mg/l after administration of 1,800 mg of Co-EDTA over 12 hours. In total there was a massive overdose of Co-EDTA administered, which may have been the cause of the fatal outcome.*

In an addendum, Hillman et al (1974) reported another case of a patient admitted after ingestion of a small amount of NaCN who collapsed several minutes later but without loss of consciousness. He received one ampoule of Co-EDTA and another one 4 hours afterwards. He then developed diaphoresis, anginal pain, ventricular extrasystoles, and intense nausea and vomiting. The antidote was stopped and the symptoms disappeared. These symptoms were attributed to Co-EDTA. 36 hours afterwards, although he was well, he developed a red maculopapular rash on the thorax and the arms which cleared over the following day. This eruption was attributed to the antidote. *Comment: In this mild poisoning case the side effects of Co-EDTA were in the foreground.*

Hoang The Dan et al, 1981

A female chemical assistant (25 years old) attempted to commit suicide with KCN. 30 minutes after intake she was in deep coma, showed gasping respiration with intermittent apnea, bilateral mydriasis, pupils reactive to light and hypotonia with 80 mm Hg systolic blood pressure. She was ventilated with 100% oxygen and 2 mg of HOCO was administered. When she reached the ED 30 minutes later no reflexes could be triggered and the pupils were not reactive to light. Blood pressure and heart rate could not be detected and the ECG showed alternating brady- and tachy-cardia (cardio-circulatory instability / failure). The treatment continued with ventilation, application of 500 ml plasma (*resulting in acute heart insufficiency, treated with dobutamine*), sodium bicarbonate as well as 600 mg of Co-EDTA without glucose and infusion of STS (*20 g infused during the next 12 hours*). The blood gases showed severe acidosis (pH 7.08, anion gap 33 mmol/l). The levels normalised within 4 hours after the antidote administration. The cyanide blood levels were determined as 5.25 mg/l upon admission and 0.5 mg/l after 16 hours. The patient could be extubated after 12 hours as an in-patient and recovered completely. *Comment: A case of severe poisoning in which Co-EDTA and STS were fully effective. The very low HOCO dose was ineffective.*

Humbert et al, 1977; Braico et al, 1979

A 11-month-old child ingested from one to five 500 mg amygdalin tablets used by her father for cancer treatment and became listless and vomited after half an hour. In hospital breathing became irregular, and somnolence set in. 1 hour after ingestion, the girl was in shock and coma, and had Kussmaul (laboured and deep) respirations and acidosis. The level of cyanide in blood was determined as 0.29 mg/l. 3.5 hours after ingestion of 1.5 ml SN (3%) and 6 ml of STS (25%) were given a maximum MetHb of 20% and improving signs and symptoms was observed. However, 14 hours after ingestion respiratory arrest occurred, and the child died after 72 hours in spite of 2 more doses of SN (0.8 ml 3%) and STS (3 ml 25%). Before the respiratory arrest no cyanide and no MetHb were detectable in blood, and after the first re-treatment MetHb was only

1%. *Comment: As the initially measured cyanide level was not in a life-threatening range, and as no cyanide was detectable when the deterioration occurred, it seems doubtful that this death was indeed due to amygdalin/cyanide. Methaemoglobin can also be ruled out as cause of death.*

Humphrey, 1978

A 66 year old woman with severe hypertension received SNP infusions. After a total of 490 mg and 28 hours the patient became agitated and tachypnoeic. SNP was discontinued after lactic acidosis was found. She recovered uneventfully after bicarbonate application. *Comment: No cyanide antidote was given in this assumed cyanide poisoning from SNP.*

Hung et al, 2009

A 32 year old Taiwanese woman drank from a soda-bottle that had been contaminated with KCN, either with criminal intent or accidentally. At once she lost consciousness and tendon reflexes, and had to be intubated and ventilated. Her oxygen-saturation could be improved from 79% at the beginning to up to 100%. She exhibited alkalosis (pH 7.846, *possibly due to artificial hyperventilation and a base excess of 26 mmol/l*). 1 hour later her blood pressure dropped to 85/46 mm Hg and she became acidotic (pH 7.096, BE 21mmol/l, lactate 14 mmol/l and anion gap 37.6 mmol/l). She was given 2.5 g HOCO which led to a quick improvement, however, the antidotal therapy was repeated 1 hour later because acidosis persisted. The cyanide level in the blood was measured once at 288.6 µmol/l corresponding to 7.8 mg/l. The woman could be extubated the next day and made an uneventful recovery. *Comment: In this case of severe poisoning HOCO was effective.*

IPCS, 1993a

Four cases are specified in “Antidotes for Poisoning by Cyanide (Volume 2)” without citing the sources. 2 cases of cyanide intoxication (cyanide uptake by ingestion) were confirmed by measurements of the blood levels (9.9 mg/l and 3.7 mg/l respectively). In the third case, also cyanide uptake by ingestion, the cyanide concentration after therapy with STS was 0.26 mg/l. The fourth case (two 2.5 year old twins) was a combined cyanide and carbon monoxide intoxication with cyanide levels of 1.15 mg/l and 1.1 mg/l respectively and carbon monoxide concentrations of 5.7% and 1.4%. All cases were successfully treated with STS; no sequelae are described.

IPCS, 1993b

A 32 year old woman who suffered from a hemangioblastoma received Laetrile treatment while in Mexico and arranged for a continuing supply of Laetrile in Canada. In a suicidal attempt she ingested 9 g of the parenteral preparation of Laetrile. On admission at the hospital she revealed severe metabolic acidosis and hypoxemia. Her clinical condition and her blood gases were not stated in the paper, but the cyanide levels (3.85 mg/l) were in the lethal range. She was treated with AN by inhalation and 300 mg of SN i.v., then a further 50 mg of SN was given i.v. and 50 mg orally. The patient survived. *Comment: A case of mild poisoning in which SN treatment was effective without side effects.*

Jacobs, 1984

The lower part of a male worker's body was splashed with liquid HCN and the worker inhaled the gases evolved. When he wanted to leave the room he fell into a coma and had to be rescued by first-aid staff. The contaminated clothes were removed. He did not show any signs of spontaneous respiration, pupils were not reactive to light, pulse was weak and fast and blood pressure was low. The cyanide level in blood was 6.5 mg/l in the first evaluation rising to 16.9 mg/l after 45 minutes and dropping to 3.0 mg/l after 4 hours. The treatment consisted of ventilation and one ampoule of DMAP i.v. together with 60 ml of a 25% STS solution (12.5 g). The patient started to breathe spontaneously 2 minutes after the DMAP administration. At first he recovered quite well (pupils reactive to light again, reaction to pain positive, spontaneous breathing through tubus) but 2 hours later he developed cerebral seizure activity. Because of that he was anaesthetised and ventilated. He was diagnosed with a 'non-spontaneous transitory [psychotic] syndrome' (German: 'Aspontanes Durchgangssyndrom'). He stayed in a coma for 4 weeks and after neurological rehabilitation some permanent CNS-sequelae such as psychomotoric retardation and organic psychosyndrome remained. *Comment: In this case of severe intoxication by both dermal and inhalation exposure in which DMAP and STS initially showed very good efficacy. The author gave no reason for the secondary sequelae-development which may be due to delayed absorption of HCN through the skin.*

Jaeger et al, 1977

This is a case of a suicidal ingestion of 40 g acetonitrile by a 26 year old man. Symptoms appeared after a 3 hour delay and included vomiting and convulsions followed by coma with respiratory insufficiency, severe metabolic acidosis (pH = 6.40) and a state of shock with 2 recovered cardiac arrests. After 10 hours he was treated with oxygen ventilation, correction of shock and acidosis, and administration of the antidotes Co-EDTA, SN, STS, and HOCO. The notable later course of events was a coma for 6 days and a syndrome of multi-organ failure.

Muscle enzymes were especially elevated and histological examination showed a process of muscular destruction followed by regeneration. The very slow elimination of acetonitrile is confirmed by the persistence of thiocyanate in the urine 20 days after intoxication. The patient recovered fully after three months. *Comment: A case of severe poisoning in which the combined administration of Co-EDTA, SN, STS, and HOCO did not improve the neurological status for 6 days. Despite multi-organ failure the patient survived.*

Jamali, 1993

A 27 year old male presenting apnea episodes that was treated with oxygen, HCO₃, dobutamin and administered 4g of HOCO and 8g of STS but did not survive. Personal communication with Prof. Baud.

Jeannin et al, 1961

A 4 year old girl was found with headache, fatigue and increasing somnolence. Upon arrival in hospital, she was in a coma, sweating, breathing slowly, without palpable pulse and with tonic contractions of the extremities. Gastric lavage revealed fruit pits and a bitter almond smell. She had played with her sister and prepared a mixture of about 30 ground peach pits with sugar which she ate about 30 minutes before the onset of the symptoms. She was treated with oxygen and 50 ml methylene blue 1%, which led to full recovery. *Comment: Cyanide in blood was not analysed, but cyanide poisoning is likely due to the smell of bitter almonds. It remains unclear whether the methylene blue was effective beyond the oxygen treatment.*

Jerretin, 1963

A 24 year old man ingested 1.5 g of KCN in a suicide attempt. He arrived in hypertonic coma with mydriasis, cardio-vascular collapse (blood pressure 50 mm Hg and pulse 100/minute) and respiratory depression. He was intubated and ventilated with 100% oxygen, and underwent gastric lavage. He received 200 mg of SN 1 hour after ingestion, a second dose of SN combined with STS 5 hours and 30 minutes after ingestion as well as one ampoule (about 20 ml) of Co-EDTA six hours after ingestion. The next day he received another ampoule of Co-EDTA. The coma cleared after injecting the Co-EDTA. The patient remained amnesic until the injection of the second ampoule of Co-EDTA. He finally recovered without sequelae. *Comment: A case of severe poisoning in which despite of application of several antidotes recovery took more than a day, so efficacy of the antidotes remains doubtful.*

Johnson and Mellors, 1988

A 30 year old male laboratory supervisor presented to the ED with stupor and a rapid, irregular pulse. The patient ingested approximately 24 ounces of beer, 35 mg clorazepate dipotassium, 7.5 g of acetoaminophen, and 3 g granular NaCN. Shortly after the ingestion of the cyanide the patient vomited and called an ambulance. Ipecac was administered and he vomited again. Initial vital signs were blood pressure 90/60 mm Hg, pulse irregular at 160/min, and respirations 20/min. No focal neurological findings were observed. No bitter almond odour was detected. An arterial blood gas on 100% oxygen was pH 7.35, pCO₂ 39 mm Hg; a simultaneous venous blood gas drawn from the median antecubital vein was pH 7.35, pCO₂ 22mEq/l. An ECG showed atrial fibrillation at a rate of 160/min and ST segment depression in the inferior and lateral leads. 2 ampoules of AN were administered by inhalation, followed by 0.3 g of SN i.v. and 12.5 g of STS. A few minutes later the patient became alert and conversant. Gastric lavage was performed and magnesium citrate and charcoal were instilled. 4 hours after ingestion the serum thiocyanate level was 1.5 mg/dl, MetHb level was 3.6%, ethanol level was 56 mg/dl and the acetaminophen level was 28 µg/ml. *Comment: A case of moderate poisoning in which the 'Lilly-Kit' was fully effective.*

Johnson et al, 1989

A woman, aged 24, attempted suicide with an unknown amount of KCN after a domestic dispute. She lost consciousness, developed a metabolic acidosis and was unresponsive to all stimuli. Blood pressure, heart rate and respiration were all within the normal range. The therapeutic actions included administration of 2.0 mg of naloxone, intubation, ventilation, AN pearls, 300 mg of SN (followed by a mild decrease of the blood pressure from 124/80 to 100/80 mm Hg) and 12.5 g of STS. Methaemoglobin levels were 7.1% during inhalation of AN and 9.2% about 10 minutes after injection of SN, decreasing to 3.6% about 5 hours, and 0.2% about 18 hours, post SN. The patient became alert within 15 minutes and recovered completely (gastric lavage was additionally performed and activated charcoal was administered). 45 minutes after antidotal treatment the pH had normalised (7.4) and the MetHb was still below 10% (exactly: 9.2%). This decreased further (to 0.2%) during the following 18 hours. The cyanide level was determined once before the nitrite application as 13 mg/l (*surprisingly high in contrast to the mild clinical presentation*) and dropped to undetectable levels by the next day. The patient was extubated and transferred to the psychiatric service. She was seen in follow up three months later with no physical or neuropsychiatric sequelae. *Comment: A case of severe poisoning in which the 'Lilly-Kit' was effective and caused only minor side effects (blood pressure drop).*

Jones, 1992

A 48 year old man and a 53 year old female were found dead at their home. Both victims had vomited during the night. Circumstantial evidence indicated that the couple might have ingested ethanol and/or methanol. But analytical examinations identified acetonitrile at concentrations of 0.8 g/l in blood, 1.0 g/l in urine, and 1.3g/l in stomach contents. The cyanide concentrations in the victims were 2.4 mg/l (female) and 4.5mg/l (male), respectively. *Comment: a case of severe poisoning without the use of antidotes.*

Jouglaud et al, 1974

A 34 year old male chemist attempted suicide by ingesting 1.5 g of KCN. He arrived at the hospital in deep coma with mydriasis and hardly palpable pulse. He was given 3 g of HOCO (one quarter of the ampoule lost during its manipulation) in STS. 10 minutes later he reacted to pain stimuli. Several sudden episodes of cyanosis occurred prompting intubation and mechanical ventilation. Severe coma reappeared, so a further 4 g of HOCO were given finally prompting recovery. *Comment: A case of severe poisoning in which the combination of HOCO and STS improved the neurological status; a second dose was fully effective.*

Jourdan et al, 1993

After suicidal poisoning with 800 mg of KCN, a 28 year old was diagnosed with cerebral oedema and bilateral mydriasis. 5 hours after poisoning he received antidotal treatment combined with 100% oxygen, 0.6 g of Co-EDTA, and 4 g of HOCO (probably with 8 g of STS) followed by 10 hyperbaric oxygen treatments, but the clinical outcome was a persistent vegetative state which still persisted four years later. *Comment: A case of severe poisoning in which the neurological situation after cerebral oedema could not be improved by the antidotes.*

Kampe et al, 2000

A young man, aged 29, drank 12 bottles of beer in which he had solved 1250 mg of KCN. 10 minutes later he was in a coma, tachypnoeic (respiration frequency 35-45/min), with dilated but reactive pupils, and only three points on the GCS. The blood pressure was low (80/60 mm Hg) and the heart rate was high (95/min). He was intubated and ventilated immediately. Intravenous fluids were infused and 250 mg of DMAP was administered together with 9 g of STS. since the ambulance staff had found remains of the cyanide powder and the clinical presentation fitted to a cyanide intoxication (respiratory insufficiency without signs of cyanosis). The blood cyanide level was 6.9 mg/l (results available only hours later). The MetHb-level was measured 2 hours

after the specific treatment with the MetHb-forming agent as 9.9%. The metabolic alterations were comparatively mild (pH only 7.3). Additionally, gastric lavage was performed and activated charcoal, laxatives and furosemide were administered amongst others. The blood ethanol level was 270 mg/dl (2.7 g/l). 2 days after the suicide attempt the patient could be discharged in a good state of health. *Comment: A case of severe intoxication in which DMAP and STS was effective.*

Kasamo, 1993

A 31year old male technician in an electroplating factory, who had been suffering from temporal lobe epilepsy for 24 years and from hypertension for 2 years attempt to commit suicide by taking an unknown amount (apparently above the lethal dose) of KCN. He was treated successfully and survived without any neurological sequelae. The electroencephalograms and the nature of the seizures were not different before and after poisoning. The T₂-weighted magnetic resonance images at 9 and 51 days after poisoning showed bilateral elevation of signals in the caudate nuclei and the putamina. On the 143th and 286th days, T₂-weighted high-resonance areas were restricted to the lateral portion of the putamina. The T₁-weighted images on the 51st day showed abnormal signal elevations in both putamina, while those on the 9th, 143th and 286th days were mainly normal. Selective vulnerability of the putamen and the caudate nucleus may be due to their specific structural properties of high oxygen and glucose utilisation, and enzyme distribution. Both chronological changes of striatal damage and the absence of neurological sequelae in this patient suggest the possibility that anti-epileptics and a calcium antagonist played a neuroprotective role in the acute cyanide intoxication.

Kim et al, 1982

A 43 year old man received SNP infusion for hypertensive encephalopathy and renal insufficiency. A total dose of 2,878 mg was applied over 14 days. Peritoneal dialysis was performed initially, haemodialysis from day 9 onwards. On days 4 and 8 brief episodes of respiratory arrest occurred with spontaneous recovery. When not on SNP momentarily, a third respiratory arrest occurred on day 11. In spite of resuscitation he fell into a coma. One day later resuscitation from cardiac arrest was successful. After bilateral nephrectomy on day 15 no further SNP was required. The patient remained semi-comatose to comatose except for 4 lucid days (d 25-29), and he died on day 47. Though there were no cyanide measurements, cyanide encephalopathy was diagnosed at autopsy. *Comment: This case of encephalopathy, which the authors relate to cyanide toxicity from SNP treatment, is dubious. A cyanide antidote was not given.*

Kirk et al, 1993

Case 1: A 20 year old man was recovered from a fire scene. The paramedics found him unconscious and in convulsions. 15 percent of the body surface was burned. The heart rate was 110 beats/min. The first laboratory results showed a blood pH of 6.69, a COHb of 4.9% (3 hours later 0%) and a cyanide level in blood of 130 $\mu\text{mol/l}$ (3.48 mg/l). 10 hours later the cyanide level had dropped to 22 $\mu\text{mol/l}$ (0.59 mg/l) and another 4 hours later to 10 $\mu\text{mol/l}$ (0.27 mg/l). The patient was intubated and received 0.3 g of SN plus 12 g of STS after the laboratory results were available. 4 hours after being discovered the patient received this treatment which was repeated 6 hours later due to persistent metabolic acidosis. The highest methaemoglobin level after the antidotal therapy was 8-9% which normalised within 6 hours. The patient recovered completely. No severe side effects were reported. *Comment: In this case of severe smoke inhalation SN/STS were effective without any side effects.*

Case 2: A 39 year old man was found lifeless after exposure to smoke. He was treated with SN plus STS after he had received successful CRP. He stayed in apnoea, developed tachycardia (heart rate: 140/min) and exhibited metabolic acidosis (pH 7.17). The COHb amounted to 30% at the beginning. It normalised after 5 hours. The blood cyanide level amounted to 3.16 mg/l and was no longer detectable after 5 hours. Methaemoglobinemia peaked 2 hours after the administration of SN at 10%. The patient died 6 days later due to hypoxia of the brain. *Comment: A case of severe hypoxic brain damage causing the patient's death, probably before antidotal treatment was commenced,. The antidotes could not improve the cerebral situation.*

Case 3: A 64 year old man was found after smoke inhalation. Nothing is stated about his consciousness. The heart rate was 154/min, the respiratory rate 30/min. COHb amounted to 22% and normalised after 7 hours. Cyanide levels in blood were 0.96 mg/l at the beginning and fell to 0.5 mg/l after the next 6 hours but could no longer be detected after 13 hours. The maximum MetHb level reached 10% 1 hour after administration of the Lilly-Kit. A bundle block was seen in the ECG. The patient was put on artificial respiration after intubation. He survived and could be discharged after 5 days. *Comment: With missing data this appears to have been moderate poisoning in which the 'Lilly-Kit' was effective.*

Case 4: A 26 year old woman was found without reaction after exposure to smoke. She was tachycardic (heart rate: 106/min) and showed metabolic alkalosis (pH: 7.45). COHb was 18.3% at discovery and 7% two hours later. Ethanol in blood was measured with 79 mmol/l cyanide with 0.7 mg/l at first and was not detectable after 9 hours. Methaemoglobine reached 10% 1 hour after the application of the Lilly antidote and normalised 15 hours later. The patient recovered completely. *Comment: This was no significant cyanide poisoning, so the antidote efficacy cannot be judged.*

Case 5: A 29 year old man was exposed to a fire. 25 percent of his body surface was burned. Obviously he was not in a coma, showed tachypnoea, tachycardia and hypothermia. Laboratory results gave a carbon monoxide level of 28.7% in haemoglobin, an ethanol level of 60 mmol/l (270 mg/dl or 2.7 g/l) in blood and a cyanide level of 1.5 mg/l in blood. The patient received the nitrite antidote due to persistent metabolic acidosis and recovered completely. *Comment: A case of mild poisoning in which SN/STS seems to have been effective.*

In this case series two more patients were included. They were not treated with any antidote. One survived on supportive treatment only, the other one survived with severe neurological sequelae.

Kläui et al, 1984

Case 1: An 18 year old male ingested 300 mg of KCN and arrived in hospital in coma with a blood pressure of 100/70 and respiratory failure. After intubation he received AN. 1 hour after ingestion 0.6 g of Co-EDTA were injected, followed by another 0.3 g after which he awoke. In addition he received 10 g of STS and 0.3 g of SN. This caused a drop in blood pressure requiring catecholamines. 10 g of HOCO were given as infusion. The patient recovered fully. *Comment: Improvement was achieved by Co-EDTA, the hypotension was probably due to the SN, and HOCO did not contribute to the positive course, as given after neurological recovery. Except for hypotension from SN no side effect occurred.*

Case 2: The 16 year old friend of case 1 ingested 900 mg KCN. He had to be resuscitated upon arrival in hospital; however the pupils remained wide and fixed. High catecholamine doses could not bring the blood pressure higher than 60/30. 600 mg of Co-EDTA initially stabilised the blood pressure; after a recurring drop 300 mg of Co-EDTA were given again. This did not change the neurological status. 10 g of STS and 0.3 g of SN were given, then 10 g of HOCO. The patient had to be declared brain dead. *Comment: The 4 antidotes given could not save the patient after initial cardiac arrest.*

Case 3: A 24 year old man ingested an unknown amount of KCN. He was resuscitated after cardiorespiratory arrest by lay people until hospital arrival. He was still in respiratory arrest, in deep coma and had wide fixed pupils. In spite of catecholamine application the blood pressure could not be measured. 0.3 g of SN and 12.5 g of STS were without effect. 0.6 g of Co-EDTA normalised the blood pressure, due to electro-mechanical dissociation. An extra 0.3 g of Co-EDTA was given. No neurological improvement could be seen. 0.3 g of SN and 12.5 g of STS were given again. This patient, too, had to be declared brain dead. *Comment: The three antidotes given could not save the patient in the end. After initial cardiac arrest, SN/STS could not reverse the cardiocirculatory problems.*

Klöver and Wenderoth, 1965

A 20 year old student bought a bag of almonds which contained only bitter almonds. Though it tasted bitter she ate 30-40 pieces. A few minutes later she felt sick, developed a headache, angina pectoris and nausea. Though she felt very weak, she made her way home (1 hour) and became unconscious 40 minutes after the purchase of the almonds. She arrived at hospital by ambulance. At admission her skin was rosy, her respiration was fast and deep but sometimes stopped. The pulse rate was 92/min, blood pressure with 110/70 mm Hg, she was unconscious with seizures and smelt of bitter almonds. A gastric lavage was performed. The patient was treated with an i.v. injection of 0.3 g of SN solution (10 ml/3%) and 10 g of STS (100 ml/10%). As the respiration was insufficient, she got analeptica and had not to be intubated. 20 minutes after the antidote therapy the patient was reactive, after 50 minutes she was awake. *Comment: A case of moderate poisoning in which SN/STS was effective.*

Kreff, 1955

2 cases due to inhalation have been reported in Germany. Acetone cyanohydrin (ACH) was spilled in a plant and immediately evaporated, the condensate showed 20% HCN. A worker collapsed while trying to escape, another worker tried to help and collapsed, too. They were found 15 minutes later unconscious on the floor – one recovered fully, one died upon arrival in hospital 1 hour later. *Comment: 2 cases of severe poisoning, presumably to HCN gas, although levels of ACH or cyanide were not measured and no antidote had been given.*

The same author also reports percutaneous poisoning in a 51 year old woman. 19 litres of acetone cyanohydrin (ACH) were spilled when a glass container broke, and they splashed on her face and her clothes, which were not fully changed. 5 minutes after the accident she vomited, had dyspnoea and became comatose; 10 minutes after the incident she had convulsions. She died upon arrival in hospital. *Comment: A case of severe poisoning although levels of ACH or cyanide were not measured and no antidote was given.*

Krieg and Saxena, 1987

Case 1: A young man aged 23 accidentally drunk from a juice-bottle containing a metal cleaning solution (6.2 mg/ml cyanide). The ambulance found a comatose man with 100-120 heartbeats/min, a respiration of 30/min and a blood pressure of 80/40 mm Hg as well as a severe metabolic acidosis (pH 7.03 and pCO₂ 17 mm Hg). The oxygen-saturation was 95%. He coughed up pink sputum, the colour of his skin was pink and a pulmonary oedema could be seen in the chest-X-ray. The supportive treatment consisted of intubation and ventilation, gastric lavage, naloxone, dextrose, furosemide i.v. and morphine. The specific therapy was undertaken

with the nitrite antidote kit (0.3 g of SN and 12.5 g of STS) and was repeated after 5 hours with half the initial dose. 1 hour after the second antidote-application the patient regained consciousness and became responsive. Independently a haemoperfusion was applied for 4 hours. While it was performed the young man could be extubated. He could be discharged on the fourth day after the accident in complete remission. *Comment: A case of moderate poisoning in which SN/STS was effective.*

Case 2: A 2.5 year old girl who had drunk from a metal cleaning solution (6.2 mg/ml cyanide) was comatose, with a blood pressure of 90/30 mm Hg, tachycardia (120/min) and a tachypnoea (28/min). A smell of bitter almonds could be detected. She received AN pearls and her health state improved even before being administered a reduced dosage of the nitrite – antidote kit (76 mg of SN and 2.5 g of STS). Thereupon the patient immediately regained consciousness and was discharged with no sequelae 2 days later. *Comment: A case of moderate poisoning in which AN prompted partial improvement and SN/STS were fully effective.*

Küpper, 2009

Case 1: In this case of hydrogen cyanide intoxication (skin absorption and inhalation), the cyanide level prior to therapy was 1.5 mg/l. 4 hours after application of 12.5 g of STS the cyanide level was 0.12 mg/l and 24 hours later the intoxication cyanide was not detectable in the patient's blood. The patient recovered clinically very fast and uneventful after therapy. Neither side effects of the therapy nor sequelae were observed. *Comment: As the clinical signs have not been reported, it is unclear what degree of poisoning was effectively reversed by STS alone.*

Case 2: Acetone cyanohydrin was taken up by inhalation and skin absorption in the second case. Initially the cyanide blood level was 3.5 mg/l. 5 hours after therapy with 12.5 g of STS, the level decreased to 0.69 mg/l and 24 hours after intoxication it was lower than 0.1 mg/l. After application of STS clinical parameters, the patient's health, improved quickly. Neither side effects nor sequelae were observed after the therapy. *Comment: As clinical signs are not reported, it is unclear what degree of poisoning was effectively reversed by STS alone.*

Case 3: Likewise the exposure to a combination of acetone cyanhydrin and hydrogen cyanide was the cause for the third intoxication case. The initial cyanide level was 2.07 mg/l. 30 minutes after therapy with 12.5 g of STS the level decreased to 0.42 mg/l. Then 125 mg of DMAP were administered. Measurements of the cyanide levels performed subsequently showed levels lower than 0.2 mg/l. The therapy was followed by a quick improvement of the clinical status. No side effects of the therapy or sequelae were observed. *Comment: As clinical signs are not reported, it is unclear, what degree of poisoning was effectively reversed by STS and DMAP.*

Kurt et al, 1990

A two year old child (15.8 kg) ingested 5 to 10 ml of liquid containing 84% acetonitrile. In the ED she was asymptomatic and was discharged. 12 hours later she awoke restless, vomiting and this was followed by 3 seizures 2 hours later. Upon arrival in hospital she was comatose with a respiratory rate of 36 and pulse rate of 136, blood pressure 104/89 and marked acidosis. She received oxygen and the 'Lilly-Kit' (3.3 ml SN 3%, 17 ml STS 25%). She awoke within minutes and fully recovered. *Comment: A case of severe poisoning in which SN/STS were fully effective.*

Lam and Lau, 2000

Case 1: A young lady, aged 19, who was working as a secretary in a chemical-trading company, stayed in a room in which HCN had been liberated. After being in there for 4 hours, she collapsed and was brought to the ED. There she presented a blood pressure of 77/65 mm Hg, a heart rate of 120/min and only 3 points on the GCS. The BGA showed severe acidosis (pH 7.07), a bicarbonate concentration of 11 mmol/l, a base excess of 18.4 mmol/l and an anion gap of 42.6 mmol/l. Treatment included oxygenation, intubation, application of sodium bicarbonate (to combat the metabolic acidosis) and administration of the nitrite antidote kit (AN for inhalation and 50 ml of a 25% STS solution (12.5 g) i.v.). As the blood pressure was quite low SN was withheld. A short time after the antidote administration the patient developed cerebral seizure activity that could be controlled with midazolam and phenytoin. The patient regained consciousness soon afterwards but short-term memory and retentiveness problems persisted. *Comment: A case of severe intoxication in which STS (and AN) showed efficacy although, the patient retained memory problems for unknown reasons. No BCN determination was reported.*

Case 2: A physicist (the owner) of the company for which Case 1 was working rescued her and stayed in the environment for 10 minutes. He was admitted to the ED with severe dizziness, near syncope, tachycardia (heart rate 110/min) and tachypnoea (AF 25/min), with a GCS of 14. Under oxygenation with 100% oxygen the blood pH was 7.58. As he developed these symptoms after one of his employees had been intoxicated by cyanide gas he was preventively given the nitrite antidote kit (AN for inhalation, 10 ml of a 3% solution of SN and 50 ml of a 25% solution of STS). He recovered without any complications and was discharged home three days later. *Comment: A case of mild poisoning in which the intense antidotal treatment consisting of the whole nitrite antidote kit was actually not indicated although the overtreatment resulted in a prompt and complete recovery and does not seem to have caused any adverse effects.*

7 cases of hydrogen cyanide gas poisoning which occurred in an industrial building in Hong Kong are presented here. 2 of them were more severely injured and required specific antidotal treatment. The other 5 were mild and responded to supportive treatment alone. All

patients except one recovered completely. Cyanide poisoning was reported to be relatively uncommon in urbanised area, so high index of suspicion is important for early diagnosis and treatment. The authors believed that prevention of cyanide poisoning can be achieved by proper storage of chemicals, and by enforcing rescuers to wear special chemical protective clothing to avoid systemic poisoning because of dermal absorption of hydrogen cyanide gas. They recommended that as there are newer and safer cyanide antidotes available, each emergency department should have a stock of updated products such as HOCO.

Lang and Stintzy, 1960

Case 1: This is a French report on a 19 year old man having worked with acetone cyanohydrin and developing headaches and nausea/vomiting at work. He went home, and after further deterioration (opisthotonos) he was taken to hospital. 3 hours after leaving work he arrived there in deep coma with cyanosis, hypotension, and slow, deep respiration. Later tonic-clonic seizures set in but recovery followed. *Comment: No specific treatment was done, as the possibility of acetone cyanohydrin poisoning was stated only one day after the incident, when the patient was already recovering. No measurements of cyanide in blood were done.*

Case 2: A further case has been reported in a 23 year old laboratory worker filling acetone cyanohydrin from a rail tanker into steel drums. One of his cotton gloves was drenched, which he did not notice. He put the glove into his pocket. 5 minutes later nausea set in, vomiting occurred, and 10 minutes afterwards he lost consciousness and collapsed. AN and artificial respiration led to short improvement, but then coma and convulsions set in. The use of SN and STS also improved the status for a few minutes only. After the glove had been found, clothing had been changed, and a decontamination bath had been performed, SN and STS led to an immediate and persistent improvement [Cited by Thiess and Hey, 1969]. *Comment: The initial non-satisfactory efficacy of AN/SN/STS was due to continued absorption. After decontamination, the antidotes SN/STS were fully effective.*

Lasch and El Shawa, 1981

Two episodes of cyanide poisoning occurred in children after ingestion of apricot kernels. The first episode involved eight children who exhibited typical signs and symptoms of cyanide poisoning two hours after having ingested a large amount of apricot kernels. Seven children recovered. One died soon after admission. The second episode involved 16 children who had eaten a sweet prepared from such kernels. The symptoms and signs were identical with those in the first group but appeared one-half hour after the ingestion and were much more severe. Thirteen children recovered, two died shortly after admission, and a third child died two hours later. Apricot kernels contain a cyanogenetic substance called amygdalin, which after hydrolysis,

liberates hydrocyanic acid. This activation usually occurs only after ingestion. In the second instance hydrolysis probably occurred during the preparation of the sweet, explaining the short interval between the ingestion and the appearance of the signs of poisoning.

Lawson-Smith et al, 2010

Case 1: One patient was removed from a fire in a house. The COHb in blood was 40%, the lactate 8.2 mmol/l. Ethanol could be found in blood. He was intubated, sedated and ventilated and brought to hospital. On admission the following laboratory parameters could be determined: pH 7.33, COHb 17%, lactate 4.2 mmol/l, cyanide 58 μ mol/l = 1.5 mg/l. The patient received an unspecified dose of HOCO prior to 5 cycles of hyperbaric oxygen-treatment. After the first turn of hyperbaric oxygen therapy the COHb had dropped to 2.8%. Cyanide blood levels had gone down to 28 μ mol/l (0.76 mg/l) and to 23 μ mol/l (0.62 mg/l) after 24 hours respectively. The patient was extubated on day five and recovered completely.

Case 2: A severely injured patient was rescued from a burning apartment. He had 2-3 degrees of burns and only a GCS of 3. He was intubated and ventilated. On admission to hospital the COHb amounted to 33.5% and blood lactate was up to 9.0 μ mol/l. An unknown dose of HOCO was administered. Prior to hyperbaric oxygen treatment the blood pH was 7.32, COHb 10.2%, lactate 4.0 mmol/l and the blood cyanide level 39 μ mol/l (1.0 mg/l). After the first hyperbaric oxygen cycle pH was 7.3, COHb 3.9%, lactate 2.5 mmol/l and cyanide 14 μ mol/l (0.38 mg/l). Despite all this therapy the patient died 5 days after the incident in septicaemia, MOF, hyperthermia with irreversible cardiac arrest. *Comment: In both patients the COHb were not critical, and the cyanide levels comparable. However, the outcome was contradictory (death vs. recovery). As the HOCO dose and the clinical course of its application have not been reported, the efficacy cannot be assessed.*

Lazarus-Barlow and Norman, 1941*

Case 1: A woman aged 23 showed a pulse rate of 132 and respiration of 26. She was drowsy, the pupil were slightly dilated. After stomach lavage she was unconscious. Her respirations became gradually slower and more laboured, the pulse rate had fallen to 45, the pupils were fully dilated, and trismus was present. She died within 1 hour. Postmortem analysis showed traces of HCN and SNP.

Case2: A young surgeon was found dead. He had taken a large dose of arsenic before SNP. It was suggested, that death must have taken place very quickly and that the arsenic played no role. *Comment: In both cases of severe poisoning the individuals were found dead and therefore no antidotes had been administered.*

Lee et al, 1982

An 18 year old man with schizophrenia and cancerophobia following an alleged nutritional and metabolic antineoplastic diet and health foods had ingested 30 tablets of Laetrile (3 grams) in a suicide attempt. He arrived in hospital deeply comatose, cyanotic, diaphoretic, with dilated pupils and hypotension. Gastric lavage was performed with activated charcoal, and he underwent mechanical ventilation with 100% oxygen. 5 hours after admission his status improved and he recovered without sequelae. Cyanide levels in blood were not measured. *Comment: No specific antidote was given.*

Lee-Jones et al, 1970

Case 1: A 25 year old man arrived at the accident and emergency department at midnight, deeply unconscious. He was pulseless and apnoeic, but the skin was warm and the mucous membranes were pink. He was in cardiac asystole and, because cardiac action was not restored by external cardiac massage and intravenous isoprenaline a transvenous bipolar pacemaker catheter was introduced into the right ventricle. Cardiac response to the pacemaker impulse was intermittent and he died 30 minutes after admission. Forensic examination showed cyanide in blood and tissues. *Comment: This case will not be part of the antidotal statistics as no specific antidote was given*

Case 2: A 35 year old man was found unconscious in the street. On admission in hospital he was deeply unconscious, with fixed dilated pupils and absent deep tendon reflexes. The blood pressure was unrecordable, the heart rate 40/min. He was not cyanosed, respiration was shallow and infrequent. Ventilation was manually assisted via a cuffed endotracheal tube. As the diagnosis was unclear at the time, lumbar puncture was performed. The patient went into asystole. Resuscitation with external cardiac massage, isoprenaline and a transvenous pacemaker restored circulation only temporarily. As at that time a suicide note with links to cyanide poisoning was found, 0.3 g of SN and 12.5 g of STS were given i.v. Nevertheless asystole recurred and the ventricle failed to respond to the pacemaker impulse. *Comment: The time that had passed between the intoxication and the specific treatment was not stated explicitly. It might have been long as the patient died a short time after the antidotal therapy because of cardiac arrest. This case describes fatal cyanide intoxication as treatment started too late. It points to the reality that after cardiac arrest reanimation does not seem possible even if nitrite is used. This may differ from DMAP where resuscitation under these circumstances was possible in four cases but not that did not occur without severe sequelae.*

Case 3: A 25 year old technician accidentally drank from a bottle about 1 g of KCN. An 18-minute treatment was started. The patient was drowsy, his heart rate was 120/min and regular, blood pressure 110/80 mm Hg, tachypnoea: 30/min. He was treated with inhalation of AN plus

0.3 g of SN, 12.5 g of STS i.v. Aspiration and lavage were then performed and a further 12.5 g of STS were left in the stomach. The patient survived uneventfully. *Comment: No cyanide levels nor MetHb levels after treatment and no blood gases are given. This makes the determination of the severity difficult. In terms of effectiveness of the antidote used it is not conclusive if nitrite changed the outcome, but early treatment is imperative.*

Case 4: A 14 year old girl either intentionally or accidentally drank a solution which was acquired for electroplating. The exact amount of cyanide in it stayed unknown. Twelve minutes later the girl was admitted to hospital. She was fully conscious with a regular heart rate of 120/min and a blood pressure of 110/70 mm Hg. She was treated with AN by inhalation, 0.3 g SN, 12.5 g STS i.v. and after gastric lavage with the same amount STS orally. The outcome was uneventful. *Comment: The only conclusion that can be drawn from this case is that the treatment did not do any harm.*

Leor et al, 1986

After i.v. injection of ampoules of 3 g each of amygdalin, followed by oral application of a third ampoule on the next day, a 65 year old woman developed diarrhoea two hours later, followed by progressive stupor and convulsions. In hospital she was in deep coma, and received oxygen 100% and antidotal treatment with SN (0.6 g) and STS (dose not specified). Within eight hours she improved and regained consciousness. However, she died 11 days after admission of liver failure, the underlying disease being hepatocellular carcinoma and cirrhosis. Cyanide in blood was 0.23 mg/l two hours after admission, so cyanide poisoning is certain. *Comment: The antidotes were primarily effective; death was due to hepatic necrosis on the basis of underlying disease and hypoxemia.*

Liebowitz and Schwartz, 1948

After taking an estimated 3-5 g KCN in a suicide attempt, a 60 year old chemist vomited once half an hour later and arrived in hospital one hour after ingesting the poison. At that time no history about the poisoning was available. He was comatose, did not react to painful stimuli and showed generalised muscular rigidity. Blood pressure was 120/80, pulse rate 140 and respiratory rate 40/min. Respiration was stertorous [heavy snoring], deep and rapid, with an expiratory grunt [deep guttural sound]. The skin was pink, cold and clammy, and the pupils were dilated. The pulse was very weak and the heart sounds were inaudible. Gastric lavage was performed immediately and an infusion of 1000 ml of 5% glucose was started. Half an hour later the patient regained consciousness. Two hours after admission the CO₂ combining power of the plasma was 25 volumes per 100 ml, and rose to 41 volumes on the third day. Blood cyanide determinations revealed 20 mg HCN/100 ml leading to an estimated total of 1.2 g of HCN in the total blood

volume. Thirteen hours later the tests revealed the presence of 0.25 mg HCN/100 ml of blood. A qualitative test for MetHb was negative, as well as the blood thiocyanate determinations on the first, second, fourth and fifth day. Free cyanide in the second twenty-four hour specimen of urine was found to be 0.116 mg in 1160 ml of urine. Eight hours after admission he was alert, well orientated and cooperative. His only complaints were weakness and nausea. The patient was discharged on the fifth day. *Comment (WM): After ingestion of an unusually large amount of KCN, the patient recovered promptly and uneventfully without any specific therapy.*

Litovitz et al, 1983

A 23 year old female chemistry student intentionally took a teaspoon of KCN and was brought in with seizure activity, tachycardia, mild hypotony (blood pressure 92/50 mm Hg) and clonic spasms of the extremities. The reaction to painful stimuli was preserved. Therapy was initiated about 30 minutes after the intake of the poison with oxygen, naloxone, dextrose, and as cyanide antidotes AN pearls were crushed and SN (0.3 g) as well as STS (12.5 g) were injected. A metabolic acidosis could be seen in the blood gases (determined immediately after treatment had begun, and moreover after 20 and 30 minutes) with a pH of 7.17, 7.26 and 7.36 respectively. One ampoule of sodium bicarbonate was given between the later two blood gas examinations. The pCO₂ also increased from 18 mm Hg via 33 to 45 mm Hg. Due to the dextrose infusion she developed hyperglycaemia with a blood sugar of 720 mg/dl. The blood cyanide level was measured once with 6.1 mg/l. Despite this high level and the clinical and metabolic alterations she recovered within two hours and later only complained about nausea and vomiting. *Comment: The Lilly-Kit was effective in this moderate poisoning case.*

Nineteen months later the same young woman was seen with tonic seizures after a collapse. She developed respiratory arrest and mechanical ventilation was initiated. The heart rate increased from 10/min (at arrival) to 110/min 17 minutes after admission. During this time span the cyanide antidote kit (AN pearls, 0.3 g SN and 12.5 g STS) had been administered. An odour of bitter almond could not be detected by several medical persons. Due to the circulatory distress, cyanosis and dilated, fixed pupils, she was intubated and CPR was performed. Dopamine was injected and caused the blood pressure to rise (80/50 mm Hg) and respiration recovered. Acidosis (pH 7.19) could be seen in the blood gas analysis. The coma persisted, so that the patient received multiple cycles of hyperbaric oxygen. The initial blood cyanide level (ca. 40 minutes after cyanide ingestion) was 683 µg/dl (68.3 mg/l) and dropped down to 12 µg/dl (1.2 mg/l) cyanide (two hours after the hyperbaric treatment began) with a corresponding MetHb level of 14.3%. Based on the fact that the results of the blood samples were only available eight hours later, a second dose of STS (12.5 g) was administered two hours after the hyperbaric treatment had begun. Despite all these efforts the woman died four days later because of multiple brain damages, as the authors suppose. *Comment: This severe poisoning case with respiratory*

arrest could not be reversed with the Lilly-Kit, and hypoxic brain damage was probably the cause of death.

Lorz, 1950

The death of a child was due to lice treatment with an acrylonitrile product. During autopsy a strong bitter almond smell was noticed. *Comment: As this child was found dead, this case cannot contribute to antidote evaluation.*

Losek et al, 1991

After ingestion of approximately 60 ml Super Nail Off (containing acetonitrile) a 23-month-old boy vomited three times post ingestion, but he did not seem to be ill. His blood pressure was 96/60 mm Hg, his pulse rate 144 beats/min, and his respiratory rate 40 breaths/min, blood pH 7.43 and anion gap 13. Eight litres/min of oxygen by mask was initiated and the patient was admitted. Twenty-four hours post ingestion the patient began having staring episodes and was not responding to his mother. Vital signs were heart rate 130 beats/min; blood pressure 110/50 mm Hg, respiratory rate 30 breaths/min. Oxygen saturation by pulse oximetry was 93% on 8L/min of oxygen mask. The low oxygen saturation prompted the administration of amyl nitrite, but without success. An arterial blood gas analysis revealed the following values: pH 7.43, PCO₂ 26.4 mm Hg; pO₂ 167.6 mm Hg, HCO₃ 17 mm Hg, oxygen saturation 99%. Lactic acid concentration was 50.1 mg/dl and MetHb level was 0.7 g/dl. A later treatment included activated charcoal with magnesium citrate, intravenous sodium thiosulphate 1.65 ml/kg every four hours. After two doses of STS, the lactate level decreased to 14.2 mg/dl (normal 4.0 to 15 mg/dl) and arterial blood gas values were pH 7.42, PCO₂ 33.8 mm Hg, pO₂ 435.8 mm Hg HCO₃ 21.8 mEq/l and oxygen saturation 99.9%. During his hospital stay, the patient received five doses STS every four hours. The patient was discharged on the third hospital day. → The blood cyanide levels were measured to be 2.1mg/l.

Lundquist, 1992

Case 1: A 28 year old young man tried to commit suicide by drinking 10 to 50 ml of a metal cleaning solution that contained 9% cyanide. He collapsed, vomited and fell in a coma. At admission to the ED one hour later he showed dilated pupils, the heart rate was 90/min, the systolic blood pressure 160 mm Hg and the blood gas analysis showed metabolic acidosis with a pH of 7.12 and a BE of -22.8mmol/l. He was intubated and ventilated and gastric lavage was performed. Moreover, sodium bicarbonate and STS (95 mmol) were applied i.v. This was followed by a tachycardia (140/min) with elevation of the ST segment on the ECG. About

30 minutes after he had ingested the poison he received 0.3 g Cobalt-EDTA. Although the drug was not combined with glucose as it is generally recommended, the patient did not develop any adverse effects, but became responsive 15 minutes after the administration of the antidote. The cyanide level was measured in a whole blood sample which contained 105 $\mu\text{mol/l}$ (according to 2.84 mg/l) on the same day and 4.5 $\mu\text{mol/l}$ (0.12 mg/l) on the next day. The plasma levels were 1.8 $\mu\text{mol/l}$, corresponding to 0.05 mg/l on the first day and 0.6 and 0.7 $\mu\text{mol/l}$, corresponding to 0.02 mg/l on the second and third day post exposition. Eleven hours after the incident he was extubated and recovered completely. *Comment: This moderate or severe case (depth of coma is unclear) was effectively treated with Co-EDTA and STS.*

Case 2: A 28 year old man was found comatose and convulsing. On admission at the ED one hour later he was intubated and ventilated because of insufficient respiration. Moreover, he showed dilated pupils, hypotonia (systolic blood pressure 65 mm Hg) and hypothermia of 34°C. The blood pH was 7.11 and the BE -16.5 mmol/l. The patient was given naloxone and cortisone, though without any positive reaction. A brain CT showed small bleeding in basal ganglia and probable brain oedema (confirmed in a second CT, 12 hours later). About six hours after the incident KCN powder was detected in the man's flat. The blood cyanide level was 139 $\mu\text{mol/l}$ (3.76 mg/l) at this time (samples were analysed on day 11 after the incident). Gastric lavage was performed and 120 mmol of STS (corresponding to about 30 g) were administered i.v. This resulted in prompt recovery, the patient regained consciousness and could be extubated, but 15 hours after the intoxication he developed – for whatever reason – diffuse cerebral oedema and died after five days. *Comment: This was a severe case of poisoning; however, the patient was – initially – effectively treated with STS alone. Cyanide levels were probably even higher shortly after ingestion, and the brain oedema had begun to develop early in the clinical course. The developing brain oedema could obviously not be reversed and progressed.*

Lurie, 1953

A man, aged 47, entered a room which had been fumigated with hydrocyanic acid. Two or three minutes after exposure, he temporarily lost consciousness but 15 minutes after the incident he was presented at the EC fully conscious. However, he was pale, bradycardic, with weak peripheral reflexes and complained about shortness of breath and nausea. He was given methylene blue, nitrites and calcium thiosulphate. Despite this treatment he lost consciousness again and became cyanotic a short time after the antidote administration. ACTH was then applied, resulting in a prompt improvement of his clinical condition. Three hours later the ACTH therapy was repeated and 12 hours after the intoxication the patient was completely orientated. According to the author no changes were detectable in the blood parameters. *Comment: The patient was initially given antagonistic antidotes (methylene blue and nitrites/calcium thiosulphate). Perhaps this was why no changes in his condition could be seen. The author then attributes the outcome of the case to ACTH, but he does not explain how this substance should be*

able to antagonise cyanide (ACTH increases the amount of suprarenal hormones). As the ACTH was administered only a short time after the other antidotes, it might be suspected that the effect of methylene blue had regressed until then and that the nitrites could develop their efficacy. As nitrites were the main 'accepted' cyanide antidotes, the case report is cited in this chapter. It seems as if the patient was orientated most of the time – though the author explicitly states the awareness at the end of the case report – so it could be discussed whether it was a mild or a moderate intoxication (I decided on a moderate one).

Lutier et al, 1971

Case 1: A 23 year old man lost consciousness and developed seizures and intermittent apnoea after the ingestion of an unknown amount (1-10 g?) of KCN dust during 1.5 hours. The initial therapy consisted of intubation and ventilation. As the patient developed circulatory instability without measurable blood pressure, aramine and sodium bicarbonate were injected. This resulted in spontaneous respiration but the coma persisted. Therefore he was given 4 g HOCO in a solution of STS. A systolic blood pressure of 130 mm Hg could be established. Four and a half hours after intoxication 10 ml methylene blue, vitamin, vitamin B1 (200 mg) and one ampoule Co-EDTA were administered after the patient had recovered. *Comment: HOCO/STS were effective in this probably severe poisoning, as Co-EDTA was given after recovery.*

Case 2: One hour after ingesting 1 g KCN a chemical engineer presented with profound coma, rigidity with trismus, bilateral mydriasis, and polypnoea with respiratory pauses. The systolic blood pressure was 140 mm Hg and the pulse was 140/minute. He was intubated and ventilated with 100% oxygen. After nitrite and STS and one ampoule Co-EDTA (300 mg in 20 ml) his condition had not improved and he received an ampoule with 4 g HOCO in STS. In the middle of the infusion, the neurological state improved, the gasps disappeared, and consciousness was regained at the end of the infusion, about one hour after the beginning of the treatment. *Comment: The application of HOCO/STS was obviously effective, though it is unclear, whether the other antidotes given before HOCO may have contributed to the outcome.*

Case 3: A 58 year old man intoxicated by KCN presented profound coma with the appearance of bouts of hypertension. He received 600 mg of Co-EDTA (2 ampoules) combined with AN inhalation. Because of the lack of improvement, he received 4 g of HOCO/STS a quarter of an hour later, and hyperbaric oxygen therapy during one hour. The coma persisted. He was extubated at the 14th hour and complete recovery occurred by 24 hours after poisoning. *Comment: HOCO/STS slowly led to recovery. But other antidotes (dicobalt-EDTA, AN) were used, which makes it difficult to assess HOCO efficacy with certainty.*

Case 4: A patient was hospitalised, following ingestion of KCN, in a profound coma with axial hypotonia, clonic movements, and mydriasis. He presented polypnoea and then more and more

significant apnoeic episodes. The hemodynamic state was initially conserved, but then degraded with unobtainable blood pressure. He received 4 g of HOCO in solution in STS. Spontaneous movements reappeared at the conclusion of the infusion; the patient regained his reflexes and spontaneous respiration. Four hours and 30 minutes after poisoning he received: 10 ml of methylene blue, 1 g of vitamin C, and 300 mg of Co-EDTA which were well tolerated. The blood pressure was 120/80 mm Hg. Recovery was complete two hours after receiving Co-EDTA. *Comment: The HOCO/STS mixture prompted partial recovery in this severe case, Co-EDTA contributed to the final recovery.*

Further cases: A man of unspecified age was found by the ambulance in a deep coma with his respiration turning from tachy- to apnoea and vice versa but with sufficient circulation (respiration rate 140 mm Hg systolic). He was intubated, ventilated and received SN in an uncharted quantity, two doses of STS (amount not mentioned either) as well as 600 mg Co-EDTA (twice the recommended dose). No significant improvement occurred, so he was given 4 g HOCO and 8 g STS. He recovered completely within a short time. *Comment: HOCO/STS seem have been most effective in this severe case, although the polypragmatic approach including Co-EDTA, SN and STS made the assessment difficult.*

This consideration can also be applied to another case of KCN poisoning (patient's age and sex, as well as respiratory or circulatory parameters were not mentioned). No supportive treatment was established (unless it was simply not reported). The antidotal therapy consisted of 600 mg cobalt-EDTA, AN, STS, 4 g HOCO and hyperbaric oxygen. This second patient also recovered from the incident. *Comment: Due to lack of data, it is unclear, which antidote (combination) was effective. The severity also is unclear.*

Mannaioni, 2002

An 80 year old diabetic patient reached the ED after a sudden collapse. His son suspected that his father might have drunk from an open bottle from the family's jewellery-factory and containing a potassium-cyanide-solution. The patient was in a coma, he was non-reactive to painful stimuli, gasping and sweating. His skin was pale, although the face was slightly pinkish. The vital signs were within the normal range and both pupils were reactive to light. In the blood samples a lactate of 9.2 mmol/l was detected, but the patient was taking oral diabetes medication. The blood gases showed a severe acidosis (pH 7.15) with a high pCO₂ (52 mm Hg) and a low pO₂ (28.3 mm Hg). The first therapeutic actions were mechanical ventilation with 100% oxygen, the infusion of liquids and sodium-bicarbonate as well as the application of activated charcoal. As cyanide-antidotes the doctors used 1 g of HOCO together with 0.3 g SN, both about 40 minutes after his admission. This did not result in a better clinical condition, so 15 minutes later a second dose of nitrite, now in combination with STS (3 g) was applied. Furthermore, 3 g of HOCO were infused within the next 24 hours. Under this treatment the elderly man's

condition improved steadily, and he started breathing spontaneously. Twelve hours after admission, he was fully conscious and he did not develop any neurological damage. During the course of his recovery different parameters were measured in the blood samples: The pH fell from 7.15 at the time of admission to 7.03 10 minutes later but could be increased and stabilised at 7.38 during the next two hours with supportive and specific treatment. Lactic acid increased from 9.2 mmol/l to 15 mmol/l 20 minutes later and then decreased to 7 mmol/l at the time of the second antidote administration. The cyanide level was 27.2 $\mu\text{mol/l}$ (corresponding to 0.74 mg/l) at the beginning and dropped down to 5 $\mu\text{mol/l}$ (0.14 mg/l) within 24 hours and to 2.5 $\mu\text{mol/l}$ (0.068 mg/l) after 48 hours. The levels of thiocyanate rose from 280 $\mu\text{mol/l}$ at the first analysis up to 310 $\mu\text{mol/l}$ 24 hours later and then fell to 175 $\mu\text{mol/l}$ during the following 24 hours. Ten minutes after the second antidote-dose the MetHb amounted to 4%, 10 minutes later to 9% and to 10% one hour after the nitrite application. Three days after the accident the patient could be discharged without any sequelae. *Comment: This was a severe case of cyanide intoxication. The rise of the thiocyanate levels indicates the detoxification of the cyanide in the liver. The combination of several antidotes was effective, no assessment as to which was the main cause for recovery can be made.*

Marbury, 1982

A 58 year old female with congestive heart failure and aortic valvular insufficiency received SNP to lower her blood pressure. This was effective, but SNP dosage had to be increased. On the third days she became confused and combative. SNP doses were lowered, other antihypertensives added to the regimen. Two days later she was lethargic with Cheyne-Stokes respiration. Two more days later her cyanide blood level was 5 mg/l, thiocyanate 24 mg/l. This prompted the application of SN 300 mg and STS 12.5 g. Haemodialysis was started, but it removed thiocyanate only, not cyanide. The total dose of SNP had been 1 g over six days. The patient survived. *Comment: SN/STS were effective in a case of moderate cyanide poisoning.*

Martín-Bermúdez, 1997

A 19 year old woman presented a coma (GCS: 7 points) and with mydriasis on both sides, which was non-reactive to light. She developed a hemodynamic instability (treated with i.v. fluids and dopamine), metabolic acidosis (with a lactate level of 5.6 mmol/l thirty hours after admission, therefore bicarbonate was administered), ARDS (oxygenation was started), renal failure with polyuria and hypernatremia as well as a rhabdomyolysis. Additionally, gastric lavage was performed and the patient received activated charcoal and the nitrite antidote kit (SN 0.3 mg and STS 12.5 g). Blood had been taken before starting the therapy and six hours later the results came in: the cyanide level was 3.04 mg/ml (*note: The units of mg/ml cited may be an error as usually the dimension unit is mg/l rather than mg/ml and 3.04 mg/ml would correspond to 3040*

mg/ and it is not possible to achieve such a value). The specific treatment was repeated (because of the blood cyanide level, whatever that means for this amount) and the patient could be discharged seven days later without any sequelae. *Comment: Severe intoxication (the patient developed diverse complications), SNS/STS were effective.*

Martinelli et al, 2008

A 51 year old female inadvertently ingested 2500 mg of Laetrile that she was using as cancer remedy. In the ED where she went, when realising she had overdosed Laetrile, she suffered diaphoresis, apnoea, bradycardia, and generalised seizures. After 5 g HOCO she quickly recovered. Whole blood cyanide was 0.4 mg/l. *Comment: This was a moderate poisoning case, in which HOCO was effective.*

Mascarenhas et al, 1969

An 18 year old man was brought to the ED 45 minutes after ingesting a calcium cyanide insecticide. He was comatose, tachycardic, with gasping respiration. He shortly suffered several tonic-clonic seizures. AN pearls were administered until a solution of 3% SN was available. 10 ml were administered, followed by 20 ml of 10% STS. His cyanosis deepened. MetHb concentration was 5.9 g/dl or about 40%. Metaraminol was administered to maintain the blood pressure over 100 mm Hg (unreported drop from the original 130/30 mm Hg). Seventy-five minutes after SN therapy, 20 ml of methylene blue were administered. Fifteen minutes later, the relatively profound cyanosis began to abate. Three hours after methylene blue, the MetHb was 0.5 g/100 ml. On admission, BCN was measured at 2.6 mg/l. The patient was discharged from hospital on day three without sequelae. *Comment: This was at least a moderate case of poisoning. The Lilly-Kit led to cyanosis, however it probably also led to recovery.*

Maxwell, 1978

Out of curiosity on what would happen a 60 year old man ingested twice a double dose of his daily Laetrile[®], and in both cases he went to hospital with weakness, light-headedness, palpitations and headaches beginning one hour after each ingestion. Upon arrival (half an hour transport) he was well in both cases. Cyanide measurement on the second presentation showed 0.6 mg/l blood. *Comment: Mild poisoning without antidote application.*

McKiernan, 1980

A male worker was contaminated with a KCN solution on 8% of his body surface. His clothes were put off and the man was washed. Although he was fully conscious, the physicians administered 20 ml of Co-EDTA i.v. The patient developed larynx oedema and vomited. He was discharged after three days in hospital and no chronic damages persisted. *Comment: In this case of mild cyanide intoxication Co-EDTA was administered without indication, resulting in larynx oedema and vomiting as adverse effects.*

MacRae, 1974

A 42 year old woman had hypotensive anaesthesia with 0.01% SNP solution. Blood pressure fell slower than usual. Over 90 minutes 250 mg were infused. During surgery the respiratory rate increased and became laboured. Extrasystoles appeared before the end of surgery. Her lips and ears were pink; the skin was warm and sweating. After cessation of the last anaesthetics spasticity in the arms and flaccid legs were noted. Lumbar puncture was normal. Bicarbonate infusion improved the symptoms. Recovery was complete. No cyanide was measured. *Comment: Potentially mild poisoning, though unproven. No antidotes administered.*

Mellino, 1980

A 52 year old female patient received SNP during treatment of myocardial infarction. The total dose was 1093 mg over 34 hours. After 32 hours laboured respiration was noted, nausea, vomiting and irritability set in. She became unconscious, and had severe lactic acidosis and a thiocyanate level of 25 mg/l. SNP was discontinued, bicarbonate was given, which prompted recovery. *Comment: Mild poisoning. No antidotes administered.*

Merryfield, 1974

A 20 year old male died after surgery with hypotensive anaesthesia. 750 mg SNP had been infused over five hours. Tachyphylaxy developed. In the recovery ward his respiration rate slowed and blood pressure decreased. Cardiac arrest in asystole followed. External cardiac compression was successful, but periods of tachycardia and asystole occurred repeatedly until he died 32 hours after cessation of anaesthesia. No cyanide was found in urine, serum or brain, and thiocyanate was less than 10 mg/l. Severe acidosis may have been the cause of death. *Comment: Probably not cyanide poisoning.*

Messing, 1991

A 29 year old student of chemistry took 50 ml of a 1% potassium cyanide solution (500 mg) in attempted suicide. He became comatose, mydriatic and was admitted to hospital in an apneic state. He woke up after seven hours and developed Parkinsonism in the following weeks. This regressed slowly in the second month after the poisoning apart from dysarthria, bradykinesia of the upper limbs and very brisk monosynaptic reflexes. Three weeks after the intoxication, CCT was largely normal, and there was CSF-dense hypodensity in both putamina after five months. Sharply delimited signal elevation in T2 corresponding to the two putamina was detected in the MRI eight weeks and five months after ingestion of the poison

Michaelis et al, 1991

A previously healthy young male (body weight 60 kg) ingested 5 ml of acetonitrile without severe symptoms of intoxication. After a STS bolus application the cyanide blood levels decreased rapidly to 10% of the initial value. *Comment: Mild poisoning, in which STS was effective.*

Michenfelder and Tinker, 1977

Twelve hours after cessation of SNP infusion in a 13 year old boy who had become obtunded and tachypnoeic and rapidly deteriorated further, until he finally died, a cyanide level of only 0.18 mg/l was found. Therefore cyanide intoxication had been ruled out. *Comment: No cyanide intoxication, no antidote treatment.*

Miller and Toops, 1951

A 62 year old drug dealer tried to commit suicide with 6 g KCN (quite a high quantity) after he had been arrested. Only three minutes later he lost consciousness, developed seizure activity and a shock. Ten minutes after taking the poison he became cyanotic, bradypnoeic (respiration frequency 4/min) and hemodynamically instable (heart rate and blood pressure could not be detected). Fifteen minutes after the incident gastric lavage was performed and four ampoules of AN were crushed over his face. He received 50 ml of STS i.v. (the concentration was not reported, so it is not known whether 12.5 g or 25 g had been injected), coramine and oxygen. Half an hour later his conditions had stabilised (respiration frequency 10/min and a weak pulse palpable). As the patient convulsed more than once, the antidote was repeated in the same quantity and dextrose was administered so that he recovered within 30 minutes. *Comment:*

Severe intoxication (hemodynamic instability, no pulse/blood pressure measurable), treatment with AN and STS was effective and successful.

Mlingi et al, 1992

An extensive outbreak of acute intoxications, with nausea and vomiting as major symptoms, occurred after ingestion of bitter cassava in a drought stricken district in southern Tanzania in 1988.

Case 1 : A 7 year old girl was admitted with abdominal pain and vomiting and was released the following morning without specific treatment. She recieved antibiotic treatment and recovered in less than 24 hrs.

Case 2 : A 4 year old girl was admitted after intensive vomiting. On admisison she was semi-conscious, dehydrated without fever, and routne neurological examination was normal.

Case 3 : A 10 year old boy was admitted unconscious after sudden onset of intoxication symptoms. Due to unavailability of antidotes he was treated with dextrose saline infusion and corticone but died 3 hours later. *Comment: No antidote was given in any case.*

Montoliu et al, 1979

A 42 year old male developed malignant hypertension with renal insufficiency and cardiac failure. SNP infusion was started. After 90 minutes and a total of 13 mg SNP, his blood pressure dropped and the infusion was stopped. However, the blood pressure fell further in spite of treatment with fluids and catecholamines. In parallel severe metabolic acidosis developed, and the patient became obtunded. About two hours after the infusion was stopped, a refractory cardiac arrest occurred. *Comment: No cyanide measurement was done, and no antidote was applied.*

Morse et al, 1979

In Mexico, a treatment with Laetrile® via i.v., intramuscular, oral and rectal routes in a 48 year old woman resulted in cold sweats, headaches, nausea, lethargy and dyspnoea. Upon hospital arrival she had Kussmaul respiration, but improved after fluid replacement. A combined cyanide blood sample (2 ml at admission, 4 ml 6 hours later), showed 1.16 mg/l, which makes cyanide poisoning certain. *Comment: A mild poisoning case without antidote use.*

Morse et al, 1981

A 32 year old woman used Laetrile for four years. In an emotional crisis she ingested 9 g of the parenteral preparation of Laetrile in an attempt at suicide. She was immediately taken to hospital, where a gastric lavage was performed. AN was administered by inhalation and 300 mg of SN were given i.v., then 50 mg of STS were given i.v. and another 50 mg was administered through nasogastric tube (Eli Lilly Kit). After her admission metabolic acidosis and hypoxemia were observed; the serum cyanide level was 3.85 mg/l. Twelve hours later the cyanide serum level was 3.5 mg/l and the thiocyanate serum level was 0.22 mmol/l (1.3 mg/dl). She made a full recovery. *Comment: The repeated combination of AN and STS was effective, however the severity of poisoning is unclear.*

Moss et al, 1981

A 32 year old woman with von Hippel-Lindau's disease ingested 9gms of Laetrile in a suicide attempt. She was immediately hospitalised and underwent gastric lavage. AN was administered and 300mg SN and 50mg STS i.v. then a further 50mg STS by nasogastric tube (Lilly kit). At administration she had severe metabolic acidosis and hypoxemia and a blood cyanide level of 143µmol/L (385µg/l). 24 hrs later it had dropped 13 µmol/L (35µg/l) and the serum thiocyanate 0.22 µmol/L (1.3µg/l). After correction of her metabolic abnormalities she recovered without sequelae.

Motin et al, 1970

A 54 year old man ingested 1 g of KCN in a suicide attempt. While initially showing only mild signs of poisoning, upon arrival in hospital he had lost consciousness, but he still reacted to pain. Trismus and mydriasis were present, while breathing was irregular. Mechanical ventilation was used after intubation and gastric lavage. Then 300 mg SN, 2 g STS and 300 mg Co-EDTA were given. As this did not improve the status 4 g HOCO in 500 ml 10% STS (50 g) were given, which led to rapid and full recovery. *Comment: One more case of recovery after HOCO/STS application, while the other antidotes given before had not been effective.*

Mueller and Borland, 1997

A 39 year old woman was admitted after swallowing 25 g of acetonitrile two hours previously in a suicide attempt. She vomited and physical examination was unremarkable. Eleven hours after ingestion, she became nauseated, confused, sweaty and tachycardic (140 beats/min). She developed Kussmaul respiration and became rapidly comatose. She was given 20 ml of 3% SN

and 200 ml STS intravenously as bolus dose. Grand mal fits occurred, she was intubated and ventilated. Two hours later the methaemoglobin level was 15%. On day 2 she was stable but 32 hours following the ingestion, she became hypotensive and tachycardic and a further 30 ml of SN and 3% STS were administered. A continuous infusion of 3% SN at a rate of 2.5 ml/min was administered from day 4 to day 5. She was finally discharged 26 days following admission. Blood levels of acetonitrile (max. 640 mg/l) and cyanide (max. 2.7 mg/l) were measured. The halftime of acetonitrile was 36 h and of cyanide 44 h and the harmful blood cyanide level persisted for over 24 hours after ingestion. *Comment: The delayed occurrence of symptoms after ingestion of acetonitrile and the delayed reoccurrence after the first treatment required an intensive observation of the patient during two days. The repeated i.v. administration of SN and STS was successful.*

Muraki et al, 2001

Neither acetonitrile nor cyanide could be measured in a new case of suspected, yet unproven, acetonitrile poisoning describing the development of massive rhabdomyolysis and acute renal failure. *Comment: From the report it is not clear, whether and to what degree an uptake of acetonitrile from the described exposure had occurred.*

Musshoff et al, 2002

Case 1: A 38 year old goldsmith was found dead in a large pool of blood. In the closet a plastic vial labelled “Fabvergoldungsbad” (gold plating solution) was found. The analysis identified pure NaCN. An autopsy was performed three days later. The chemical analysis of different specimens yielded the following cyanide concentrations: blood 80.9 mg/l, gastric content 1.26 mg/l, lung 46.3 mg/kg, brain 4.8 mg/kg, kidney 0.89 mg/kg, liver 17.6 mg/kg, bile 21.3 mg/l. The distribution of cyanide in the specimens examined indicated oral ingestion. *Comment: Patient was found dead, no treatment.*

Cases 2 and 3: A physician and his wife, both 71 years old, were found dead. The examination of two vials found besides them yielded positive results of cyanide. Analysis of the blood samples quantified cyanide levels of 6.1 and 8.6 mg/l, respectively, which indicated severe intoxication and provided adequate evidence to attribute both deaths to cyanide toxicity and to be a suicide. *Comment: The patients were found dead, no treatment.*

Case 4: The 70 year old wife of a chemist had ingested a spoonful of substance which had earlier been described by her husband as highly toxic. A few minutes later she experienced convulsions and, despite immediate attempts at resuscitation, she died two hours later. The powder in the

vials taken in at the scene proved to be NaCN. The cyanide levels were: blood 4.2 mg/l; brain 2.54 mg/l; gastric contents 1.2 g/l. *Comment: Patient was found dead, no treatment.*

Case 5: A 29 year old woman, who worked as a laboratory technician in a metal processing factory experienced seizures and collapsed, after drinking a cappuccino with her colleagues. A few minutes later she was already comatose. Four hours afterwards the woman was admitted to hospital, specimens of blood and stomach were obtained for toxicological examination. The cyanide blood level was 3 ml/l. Death occurred four days later. The colleague who had given her the cappuccino was convicted of murder. *Comment: Patient died; no antidote treatment.*

Mutlu, 2002

An 83 year old retired biochemistry professor and owner of a small chemical firm was found unresponsive in the bathtub by his wife (he was dressed in his pyjamas and his arms were crossed on his chest). The ambulance men saw a hypopnoeic patient, with a barely palpable pulse and no measurable blood pressure. During transportation to hospital pulseless electric activity occurred for which he received CRP in combination with epinephrine and atropine (1 mg each). Spontaneous circulation could be re-established within 10 minutes of resuscitation. In the ED the weak pulse and no measurable blood pressure persisted; moreover, no spontaneous respiration was detectable, the patient was hypothermic (32°C) and both pupils were mydriatic and non-reactive to light. The ECG showed atrial fibrillation (history of ischemic cardiomyopathy and a stroke) and the cardiac enzymes were elevated (hinting to myocardial infarction). Pulmonary oedema was seen in the chest X-ray. Blood gases revealed a pH of 7.1, a venous oxygen saturation of 62 mm Hg and a lactate of 6.2 mmol/l (whether the pH and lactate were also measured in venous blood is not clear). The patient was intubated and started on catecholamines. This resulted in a stabilisation of the blood pressure, but after transfer to the coronary care unit, he had become flaccid, unresponsive to external stimuli (sound, light, pain) and had lost the cranial nerve functions. His relatives found a suicide note in the bathroom, together with an empty potassium-cyanide bottle. Notifying this he was administered STS (dose not reported on) with a subsequent stabilisation of the blood pressure. Eighteen hours after the suspected cyanide intoxication the blood cyanide level was 1.0 mg/l. *Comment: Severe and finally fatal intoxication in which the delayed application of STS alone was not effective. Although the blood level is not explicitly given, it seems to be probable that the antidote was applied at the same time (18 hours post ingestion), that means much too late. As the cranial nerve function could not be re-established, nor could the peripheral reflexes, the patient was declared brain-dead the next day. Severe intoxication, died under STS treatment only.*

Nagler et al, 1978

Three workers suffered from hydrocyanic acid intoxication because of the open addition of a CN^- -salt to an acid.

Case 1: The first worker, a 42 year old man, was standing about 1 metre away from the tank that contained the solution. He became semi-comatose a short time after inhalation of the gas had begun. As the application of one ampoule of AN resulted in tachycardia (120/min), he was administered Co-EDTA together with 20 ml of a 30% glucose solution. This antidote was repeated a few minutes later (as the first dose had not shown any efficacy). This resulted in a prompt improvement of his status (he regained consciousness) but adverse effects also occurred (the patient vomited, became agitated and developed facial and peripheral oedema for 24 hours). Moreover, the blood pressure rose to 200 mm Hg systolic. *Comment: A case of moderate cyanide poisoning with effective application of both AN and Co-EDTA. However, the latter one prompted adverse effects.*

Case 2: A second worker was standing about 3 metres away from the toxic solution. He later complained about a sore throat and was given one ampoule of AN for inhalation. Two hours later he became bradycardic and one dose of the cobalt-EDTA/glucose combination was administered. He immediately complained about nausea, retro-sternal chest pain and oedema. He was discharged 3 days after the incident and treatment. At this time all adverse effects had gone. *Comment: In this case of mild cyanide intoxication the CO-EDTA was actually not required. Altogether, the treatment showed good effectiveness, however, adverse effects occurred.*

Case 3: A 46 year old worker was standing 10 metres away from the tank. As he also developed mild symptoms of cyanide poisoning (headache, nausea and a sore throat), he was given one ampoule of AN and one dose of Co-EDTA as well. He suffered from adverse effects of this antidote as did his workmates: chest pain, vomiting, tachycardia and oedema. He also had to stay in hospital for 3 days. *Comment: Another case of mild intoxication that was treated with cobalt-EDTA without indication. The patient also developed adverse effects of the antidote but recovered in the end.*

Nakatani et al, 1992

A 31 year old man presented deep coma and seizure activity. The respiration frequency decreased within a few minutes from 15/min at admission to 6/min. He was intubated and ventilated. The BGA showed acidosis (pH 7.049, BE -20 mmol/l). Gastric lavage was performed and activated charcoal administered. Half an hour after admission an empty bottle that had formerly contained KCN was found in his flat. So specific treatment for cyanide poisoning was initiated: 3 ampoules of AN for inhalation every 3 minutes (result: mild drop of systolic blood

pressure to 80 mm Hg) and 10 g of STS i.v. over 10 minutes. Half an hour later he had completely recovered. The MetHb was always below 2%, the pH increased from 7.02 to 7.5 five hours later together with normalisation of the lactate that had been up to 12.4 mmol/l before. The blood cyanide level was measured twice: 1.2 mg/l after 2 hours and 0.1 mg/l after 3 hours. *Comment: This was a severe case of poisoning, so the cyanide levels analysed 6 months later from stored samples are not reliable. AN/STS were effective.*

Naughton, 1974

Case 1: A 35 year old prisoner was seen while taking a capsule containing cyanide salt. He fell in a coma and developed tachycardia 10 minutes after the intake. The reflexes could be triggered normally and he was responsive to painful stimuli. He was intubated, ventilated and given i.v. fluids together with sodium bicarbonate to counteract the metabolic acidosis, recognisable in the BGA (pH 7.25, pCO₂ 15 mm Hg and BE -2 0mmol/l). Blood cyanides accounted for 1.0 mg/l. About one hour after intoxication the patient was given 11 pearls of AN for inhalation (despite of this high dose, the author does not report any adverse effects such as hypotonia or peripheral vasodilatation). This was followed by 600 mg of Co-EDTA (twice the recommended dose) together with 50 ml of a 50% glucose solution, 1 hour later. This resulted in a flush, atrial fibrillation and ventricular extra systoles. Therefore digitalis was administered to him. The patient regained consciousness within the next 2.5 hours and could be extubated afterwards. 4.5 hours after the incident he was completely responsive and made an uneventful recovery. The blood cyanide was 0.6 mg/l. *Comment: AN and Co-EDTA were effective in this moderate poisoning, however, the latter had significant side effects.*

Case 2: A man (age not mentioned) tried to commit suicide after he had found his wife in a comatose status (she had tried to kill herself with an overdose of sedatives). The ambulance found an unconscious, tachypnoeic patient. As he smelt of bitter almonds and an empty bottle with the same odour was found beside him, cyanide intoxication was suspected. In the blood gas analysis, mild acidosis (pH 7.3), hypocapnia (pCO₂ 16 mm Hg) and a BE of -16mmol/l were detected. He was intubated, ventilated with 100% oxygen and gastric lavage was performed together with the application of activated charcoal. Furthermore, he was given sodium bicarbonate and 300 mg of Co-EDTA twice (20 and 45 minutes after admission), each together with 50 ml of a 50% glucose solution. Two hours after the suicide attempt he was conscious again and orientated; he only complained about facial oedema (*side effect of the double antidote dose*). The cyanide amounted to 0.2 mg/l in the blood sample. He recovered completely. *Comment: Co-EDTA was effective in a moderate poisoning, but prompted side effects.*

Nicoletta et al, 2007

A 58 year old man with a long history of hypertension was admitted with aortic dissection, and put on SNP infusion. The dosage had to be increased until he developed psychomotor agitation, hallucinations and speech impairment. SNP infusion was stopped. No cyanide measurements were done, nor is the total or the maximum SNP dose reported. *Comment: Cyanide poisoning was unproven, no antidotes were given.*

O'Brien et al, 2005

A 32 year old woman ingested 6 amygdalin tablets and was found shortly thereafter by her father with a rapid progression from abnormal behaviour to gait disturbance and writhing on the floor plus somnolence. She recovered fully over 8 hours with symptomatic treatment and without antidote application, though nephrogenic diabetes insipidus had occurred. No cyanide measurement was made, thiocyanate in blood was 445 µmol/l, which indicates probable cyanide poisoning. *Comment: Mild poisoning, no antidotes applied.*

Ortega and Creek, 1978

A 3 year old boy was given Laetrile® enemas for 3 days, each with 3.5 g Laetrile®. After the second application vomiting and diarrhoea occurred, after the third dose the child became lethargic, unresponsive, tachypnoeic and cyanotic. He recovered under oxygen therapy. The blood cyanide level was 2.14 mg/l, so cyanide poisoning is certain. *Comment: Moderate poisoning, no antidotes given.*

“Patient S.T. was first admitted at the age of 2¹⁰/₁₂ years with a two-month history of constipation and back pain. Physical examination revealed a hard, nontender spherical mass in the right upper quadrant of the abdomen. Initial evaluation revealed a normal complete blood count, urinalysis, chest radiograph, skeletal survey, bone, liver and spleen scans, and bone marrow. An i.v. pyelogram (urography) revealed an extra-renal calcified mass in the right flank close to the kidney. Abdominal exploration resulted in the excision of a tumor immediately below the hilum of the right kidney. Pathology examination revealed neuroblastoma with invasion of the capsule and regional lymph nodes. Ten days after surgery, chemotherapy consisting of vincristine, cyclophosphamide, and dimethyltriazene imidazole carboxamide was initiated. However, after receiving three doses of the latter and one dose of cyclophosphamide, the patient was lost to follow-up. It was subsequently learned that the patient was receiving treatment with Laetrile, 500 mg daily by mouth and 3.5 g i.v., daily.

Because of increasing technical difficulty in administering the drug i.v., the mother was advised to use the parenteral form of Laetrile as an enema. Following administration of the second daily enema of 3.5 g Laetrile in 10 ml, the patient developed vomiting and diarrhoea which persisted into the third day of treatment. Shortly after receiving the third Laetrile enema, the patient, who previously had been irritable, became progressively lethargic and unresponsive, tachypnoeic, and cyanotic. He was then taken to a local hospital where O₂ therapy and i.v. hydration were rapidly initiated.

Because of the history of Laetrile administration, the possibility of cyanide poisoning was suggested and, five hours after admission, cyanide blood concentration levels were obtained and reported as 214 µg/dl. (The lethal range is 260 to 3,000 µg/dl). The child recovered slowly over the next 12 hours. Subsequent evaluation revealed recurrence of intra-abdominal disease with several bone metastases.”

Orušev et al, 1972

A report from Yugoslavia described a 35 year old worker exposed to acrylonitrile vapours in the open for one hour who developed some neurological symptoms like intention tremor, but recovered without specific therapy. Blood levels for acrylonitrile or cyanide are not reported. *Comment: Mild poisoning, but it is unclear whether cyanide was involved. No antidotes given.*

Pardee, 1947

There is a report from 1947 on a boy dying after ingestion of large amounts of choke (wild) cherries for days including kernels. One morning he complained about epigastric pain, vomiting and convulsions and died within an hour. *Comment: Cyanide poisoning unclear, but probable. Fatal case, no antidotes given in time.*

Patel, 1986

Case 1: A 62 year old woman had surgery for coronary bypasses and cardiac aneurysmal resection. Her blood pressure required SNP therapy after surgery. The SNP dose had to be increased progressively. In parallel her clinical status deteriorated. On day 3 she became unresponsive to pain stimuli. Total SNP had been 721 mg over 80 hours. SNP infusion was stopped and a blood sample taken. Cyanide levels were 0.585 mg/l on day 3 and 0.8 mg/l on day 4. STS was given at doses of 12.5 and 6.25 g. This prompted improvement. *Comment: In spite of relatively low cyanide levels this may have been severe poisoning, but this is unclear. STS was effective.*

Case 2 in text (4 in table): A 65 year old man with myocardial infarct had quadruple coronary bypasses and mitral valve replacement. SNP (and dopamine) were required with an increasing need for SNP. After 30 hours and 157 mg SNP total he became restless and did no longer respond to commands. SNP was discontinued and a blood sample for cyanide and thiocyanate taken. Lactic acidosis was seen. Heart rhythm and hemodynamics became unstable. Cardiac arrest occurred late on day 2 and could not be reversed. Cyanide level had been 0.5168 mg/l and thiocyanate 16 mg/l. STS had been given, though the dose is not reported. *Comment: In spite of a low cyanide level, there was cardiac arrest. STS was not effective.*

Paulet, 1965

Case 1: A 32 year old chemical engineer unintentionally swallowed a small amount of a solution containing potassium- and copper-cyanide while working. A few minutes later he developed nausea, palpitations and paraesthesia. 300 mg Co-EDTA were administered 20 minutes after the incident. This resulted in vasodilation with a flushed face, in accordance with the fact that cobalt-EDTA, when being administered without strict indication (as in this case) may cause adverse effects, even in the recommended doses. All the symptoms and adverse effects subsided within half an hour. *Comment: Mild intoxication treated with Co-EDTA, which was effective, but caused mild side effects.*

Case 2: A male patient, aged 24, had taken 1.5 g of KCN in a suicide attempt. Forty-five minutes later he was seen in the ED with pink-coloured, cool skin and dilated pupils. He was in a deep coma, respiration was shallow and weak and the systolic blood pressure was only 50 mm Hg together with a tachycardia heart rate of 100/min. The treatment consisted in 200 mg of SN, STS (dose not mentioned) and – as this nitrite-antidote treatment did not show any positive reaction – 300 mg of Co-EDTA were given twice (solved in 20 ml each time), 5 hours after admission as well as on the next day. The patient regained consciousness straight after the first dose of Co-EDTA (Co-EDTA) and made an uneventful recovery. *Comment: In this severe case of poisoning SN/STS were obviously not effective, while Co-EDTA was. The reason for the application of the second dose is not clear.*

Pentore et al, 1996

In Italy a 56 year old woman came to hospital with nausea, vomiting, respiratory problems and sleepiness. Her pupils were dilated. As she rapidly fell into a deep coma, artificial respiration was conducted, she became conscious again, but remained confused for 14 days. Later she developed signs of a parkinsonian syndrome, retrobulbar neuritis and sensory-motor neuropathy. MRI showed abnormal signal intensities involving the basal ganglia. She had eaten choke cherries steeped in alcohol (cyanide in cherries up to 15 mg/kg, in the spirit 45 mg/kg).

Comment: Cyanide levels in blood were not measured; neither antidotes, nor oxygen probably, were administered.

Perschau et al, 1977

A 14 year old boy was scheduled for a Harrington rod correction of a thoracolumbar scoliosis in association with spastic paraplegia below the level of T-6. Under anaesthesia, sodium nitroprusside (0.1 mg/ml in 5% dextrose) was infused. A total dose of 130 mg nitroprusside was administered over five hours. As the arterial systolic blood level fell below 30 torr, a cardiopulmonary resuscitation was performed. Because of the patient's severe intrapulmonary shunting, hypotension, and arterial hypoxemia, not to administer inhalation or i.v. nitrites, in order to avoid further compromising the patient by inducing elevated MetHb levels and reducing available oxygen. Therefore it was decided to administer 150 mg/kg STS (25%) and thereafter 75 mg/kg every 3 hours. By 12 hours after the cardiovascular collapse, he was responding appropriately to command. *Comment: This seems to have been a severe poisoning, STS was effective, although delayed, which would have been expected.*

Peters et al, 1982

A 31 year old chemist wanted to commit suicide by cyanide and informed his wife at 5.30. One hour later he was found in his laboratory and brought to hospital at 7.20. In hospital he showed central and peripheral cyanosis, a respiratory rate of 2/min, a heart rate of 72/min, and an arterial blood pressure of 140/70 mm Hg. He was intubated and respired with 100% O₂. At 8.00 he arrived at the ICN still comatose and cyanosed. Bilateral leg clonus and dilated pupils were observed. The patient was again ventilated with 100% oxygen and received 0.3 g SN and then 25 g STS ('Lilly-Kit') i.v. Blood samples were taken for estimation of cyanide, MetHb and other biochemical/haematological parameters. The metabolic acidosis was corrected by 200 ml 8.4% sodium bicarbonate. At admission the patient had severely displaced: pH: 7.07 ↓, pO₂: 41.2 kPa ↓, BE: -21 mmol/l ↓↓, bicarbonate: 9 mmol/l ↓↓, pCO₂: 4.1 kPa ↓↓. In the recovery phase which started at 9.45, the patient was hyperventilating, had seizures which had to be treated by diazepam and even by neuromuscular relaxation. At 10.15 the patient got a second dose of SN, and STS because of the presence of cyanide in the gastric aspirate. In addition 150 g STS in 300 ml water were infused in the stomach. As no MetHb could be measured SN/STS treatment was repeated together with AN. Twenty-four hours after admission neuromuscular blockade was reversed and spontaneous respiration established. The total amount of KCN taken was 8 ml of 5% solution = 400 mg. The blood cyanide levels were 2.6, 1.9 and 9.7 mg/l at 10.5, 14.5 and 26.5 hours after ingestion, respectively. The authors explained the last increase by incorrect sample collection. *Comment: Severe intoxication, treated with SN/STS.*

MetHb formation could not be sufficiently achieved for unknown reasons in spite of repeated dosing. However, in the end SN/STS was probably effective, but the patient survived with sequelae.

Pijoan, 1942

Case 1-3: One person, 15 years old, came to hospital with assumed choke cherry ingestion and stupor/somnolence and fibrillar muscular twitching. He recovered after gastric lavage. So did a person, 54 years old, with slight stupor, listlessness and fibrillar muscular twitching, and another person, 14 years old, with somnolence, dizziness and fibrillar muscular twitching. All three had pink cyanosis; initially all had had vomiting and epigastric pain. *Comment: In these cases cyanide poisoning can be only regarded as possible, no antidotes were given.*

Posner, 1977

In an oliguric patient (male, 66 years old) with hypertensive crisis, SNP was given at a total dose of 238 mg over 38 hours. Hypotension persisted and the patient died 24 hours later. Maximum cyanide level was 2.99 mg/l. *Comment: No antidotes were given.*

Potter, 1950

Case 1: A 52 year old man worked in a HCN-processing company. He inhaled hydrocyanic acid while trying to clean a conduct that led to a tank containing HCN. Actually he was wearing gloves and a respirator but he had to take the latter off because the fresh air flow was not working properly. A short time after he had left the room he became dizzy, dyspnoeic, developed a headache and a collapse. As first line treatment he was given oxygen and one ampoule of AN for inhalation. Five minutes after the incident the man presented a deep coma with irregular, laboured breathing, tachycardia of 110/min, non-reactive pupils and convulsions. Therefore he was administered the cyanide antidote kit consisting of 300 mg of SN and 50 ml of a solution containing 50% STS (25 g). His breathing normalised immediately, he showed spontaneous movements, murmured and the pupils were again reactive to light. Ten minutes after the antidote application he was fully conscious and made a good recovery. The author reports that the cyanide level in the blood samples was elevated (however, the exact amount is not given). *Comment: In this severe case the nitrite antidote kit showed a prompt and satisfying efficacy.*

Case 2: A man, aged 41, was employed in a company working with HCN. When he was trying to clean a gas-pipe for HCN he was hit by liquid prussic acid on his hand, as he had put off the gloves he had worn before to be able to work better. Two minutes after the accident the man complained about dizziness and dyspnoea, 5 minutes after the exposure he lost consciousness.

He was given oxygen at once and one ampoule of AN was crushed over his face. His respiration was irregular, his face was flushed, he was tachycardic (heart rate 100/min), his pupils were fixed and dilated and no deep tensor reflexes could be triggered. As 2 more ampoules of AN and coramine injection did not influence his conditions but he rather became cyanotic, 300 mg of SN and 50 ml of a 50% STS solution (25 g) were administered i.v. After this special treatment his clinical status improved quickly, he moved spontaneously, murmured, the respiration normalised and the tachycardia ceased. *Comment: In this severe poisoning AN obviously did not prompt improvement, while SN/STS were effective.*

Prieto, 2005

A middle aged woman (30 years old) injected herself with a cyanide-solution s.c. in a suicide attempt. She was found lying on the floor with only 3 points on the GCS, bilateral mydriasis responsive to light and hypotonia. Local necrosis could be detected at the points of injection. In the emergency department the patient was intubated and i.v. fluids were administered, in combination with dopamine and noradrenaline. In the blood gases a pH of 6.74 (indicating severe acidosis) was seen, with a lactate of 19.3 mmol/l and an anion gap of 33mmol/l. Sodium bicarbonate was infused and increased the blood pH (7.13). The blood cyanide level was determined at 4.6 mg/l (which shows that the resorption after s.c. injection occurred quickly and completely). A further treatment consisted of haemodialysis (lasting four hours) resulting in a pH of 7.3 and a rise in the pCO₂ (from 19.4 mm Hg to 32.8 mm Hg). A few hours later the woman reached 15 points on the GCS and could be transferred to the psychiatric unit days days later. *Comment: This is an example of severe cyanide intoxication which could be controlled with supportive treatment only.*

Queisser, 1966

This is a cassava poisoning report from an island off the coast of Tanzania. Half an hour after a meal of fresh cassava, which tasted bitter, two children (aged 7 and 9) fell ill with nausea, vomiting, chest oppression and dizziness. Half an hour later one child died after becoming unconscious. A further 30 minutes later the second child died, too, and a third child developed dizziness and nausea. Three qualitative hydrocyanic acid tests applied during *post-mortem* examination gave positive results for the gastric content of both children. *Comment: In these two severe and one mild poisonings no antidotes were given.*

Quinlan et al, 2008

A 14 year old girl had renal transplantation and developed hypertension after surgery. SNP was given, but increasing doses were needed. A total of 200 mg was given over 56 hours. Blood cyanide was found to be 3.1 mg/l and SNP was discontinued. 0.3 g SN and 19 g STS were applied and haemodialysis to remove thiocyanate. Cyanide fell quickly, and the patient recovered fully. *Comment: In spite of the cyanide level this obviously was mild poisoning, in which SN/STS may not have been required, but was effective.*

Rachinger et al, 2002

A 35 year old laboratory worker tried to commit suicide with a cyanide salt. Ten minutes after the ingestion she presented unconscious and was intubated and ventilated. DMAP and STS were given in unspecified doses. The patient developed an intermittent akinetic mutism with agitation and an irreversible cerebral ischemia. *Comment: Clinical details were only partially communicated. This seems to have been moderate poisoning, so that it is unclear, why brain ischaemia developed. DMAP plus STS have at least not been fully effective.*

Racle et al, 1976

A 68 year old man ingested 5 g of KCN in a suicidal attempt. Upon arrival in hospital 30 minutes later he was in a deep coma, had convulsions and mydriasis as well as cardiovascular collapse and tachypnoea. Mechanical ventilation was installed after intubation, and he was given 4 g HOCO in an 8 g STS solution. This led to recovery. *Comment: HOCO/STS were effective and saved the patient. The cyanide level was not measured, but the clinical course was typical.*

Ram et al, 1989

A 53 year old with intracerebral haematoma, which was removed surgically, needed SNP to control his blood pressure postoperatively. After initial improvement he became drowsy on day 7, had metabolic acidosis, and fell into a coma. Under the assumption of cyanide poisoning 250 mg DMAP and 12.5 g STS were given, and SNP was discontinued. Haemodialysis and oxygen ventilation were started. The patient's condition improved; neither cyanide nor thiocyanates were measured. *Comment: This is possible, but unproven, cyanide poisoning, probably moderate. DMAP and STS in combination with haemodialysis and oxygen were effective.*

Ruangkanchanasetr et al, 1999

Nine hours after ingestion of boiled cassava, two children, a 4 year old girl and a 1.5 year old boy, vomited and became comatose. They required intubation and ventilatory support ($\text{FiO}_2 = 1$). The girl needed antidotes (SN 3% 4 ml, and STS 2.5% 250 ml), the MetHb was measured at 3.95%, gastric lavage with application of activated charcoal and liquid paraffin was also done. The cyanide level before antidote application was 0.56 mg/l (19 hours after ingestion). The boy could be kept on supportive treatment only, though his circulation was initially instable. He had a bitter almond odour on breath, and the cyanide level in blood (23 hours after ingestion) was 0.32 mg/l. Metabolic acidosis and lactic acidemia were observed in both children. *Comment:* Two cases of moderate poisoning, in one case SN/STS, in the other case supportive treatment, were effective.

Rauscher, 1978

A 27 year old female with renal transplant developed a gall bladder rupture requiring surgery. Hypertension developed, which was difficult to control and required SNP therapy. Over 48 hours 1,120 mg were given. Then consciousness shifted between agitation and coma, acidosis was found, the hemodynamic status became unstable. In spite of intensive treatment, cardiac arrest in asystole occurred. Resuscitation and bicarbonate were successful, so the patient survived. Cyanide was not measured. *Comment: It is unclear, whether this was cyanide poisoning. Anyhow, no antidote was given.*

Riudavets et al, 2005

A 17 year old man drunk a beverage laced with KCN by another individual. He became unresponsive within a few minutes of ingesting the cyanide and suffered cardiorespiratory arrest. Resuscitation efforts were implemented. Then he was taken to a local hospital, where he received antidotes for cyanide. A blood test revealed the presence of thiocyanate (25.5 mg/l). He remained in a coma, on ventilator, and was pronounced brain dead 4 days later. *Comment: After cardiac arrest the patient succumbed to cyanide poisoning. The antidotes given were not specified, so this case cannot be evaluated.*

Robinson, 1959

Case 1: A young woman, who accidentally took 1.04 grains of hydrogen cyanide walked 150 yards, was found unconscious 5 minutes later, became comatose within 20 minutes, and died within 30 minutes of taking the poison.

Case 2: A student nurse died 2 hours after the start of symptoms with nausea, followed by coma and convulsions. 2.5 grains of SNP and 54 mg of cyanide were found in the stomach.

Case 3: A worker fell down and was found in coma with fixed, small pupils. He died in the hospital receiving room 35 minutes later. A suicide note was found and 170 mg of KCN were found in the stomach.

Comment: In all three fatal cases no antidote was given.

Roedelsperger, 2009

One case was probable inhalation by a worker working close to an acetone cyanohydrin leakage, who complained about a bad taste. Again, cyanide in blood was below the limit of detection, and except for the taste there were no symptoms. 2.5 g HOCO were given.

A skin contact (about 1 palm size) with a mixture of acetone cyanohydrin and HCN did not result in symptoms either. Cyanide in blood was not determined. 2.5 g HOCO were given.

A similar case occurred (2 palms area affected); cyanide in blood was about 1 mg/l and decreased quickly after antidote (HOCO 2.5 g) application (1.5 h: 0.15 mg/l). This worker had no symptoms either.

Comment: Antidotes were applied in all cases. However, as their application does not seem to have been required, the cases are not contributive for assessment of antidote efficacy. In one case a cyanide level of about 1 mg/l was found, however, this patient, too, was asymptomatic, and the applied antidote efficacy cannot be assessed either [personal communication].

Rosenberg et al, 1989

A 46 year old man ingested 1,500 mg of KCN in a suicide attempt. He survived, but later on developed a severe parkinsonian syndrome. The MRI revealed multiple areas of low-signal intensity in the globus pallidus and posterior putamen. A 6-fluorodopa PET study revealed bilateral decreased uptake in the basal ganglia. This evidence of functional impairment of dopaminergic nigrostriatal neurons is related either to direct toxicity of cyanide or to the effects of cerebral hypoxia secondary to cyanide intoxication.

Rubino and Davidoff, 1979

A 49 year old woman with lymphoma ingested 20-40 apricot pits as a surrogate lunch, not for cancer treatment. After 30 minutes, headache, weakness, disorientation, nausea and vomiting set in. In hospital she was given AN, SN and STS. Her cyanide blood level was 3.2 mg/l. Recovery was complete. *Comment: In this mild poisoning case, SN/STS were effective, but would probably not have been required.*

Sadoff et al, 1978

After having injected 12 g of Laetrile i.v. per day for a month without problems, a 17 year old girl ingested a 10.5 g dose orally, as injection was not possible. Shortly after the ingestion, she developed headache and dizziness, collapsed, had tetanic contractions of the hands, and generalised convulsions. Breathing was laboured; she became comatose after 10 minutes, and died 24 hours later in hospital after gastric lavage and mechanical ventilation. No antidotes were given. Cyanide in blood was normal, but the sample was taken 36 hours afterwards, and the analysis was delayed by one month. Though evidence was circumstantial only, cyanide poisoning was suspected to be the cause of death. *Comment: In this fatal supposed cyanide poisoning no antidotes were given.*

Saincher and Swirsky, 1994

A chemistry student (aged 23) took 100 mg KCN in a suicide attempt. Three hours later he became pale, incontinent, disorientated, showed uncoordinated movements and murmuring. The vital and biochemical parameters were as follows: heart rate 135/min, accelerated respiration 36/min, pH 7.02 (7.26 with oxygen), pCO₂ 24 mm Hg (34 mm Hg under oxygenation), BE -24 mmol/l and anion gap 33 mmol/l. The blood cyanide was measured at 4.65 mg/l. However, the patient was not comatose and because of this no antidote was applied though the patient reported cyanide poisoning. He was only given oxygen and 150 mmol sodium bicarbonate. A forced alkaline diuresis was performed and pneumonia treated with antibiotics. The young student became alert and responsive. The acidosis cleared (pH 7.26) and the pO₂/pCO₂ normalised. An intermittent weakness of the legs and the pulmonary affection were both reversible. *Comment: Mild intoxication in spite of high cyanide level. No antidotes were applied.*

Sarikaya, 2006

Parkinsonian syndrome developed in a young male following a suicide attempt with cyanide. MRI was reported as showing bilateral decreased signal on the T1 image and increased signal on

the T2 image in both lentiform and caudate nuclei as well as a minimally increased signal on T1 image in the cerebellar white matter consistent with hypoxic-ischemic changes. F-18 FDG PET/CT scan of the brain demonstrated absent FDG accumulation in both putamens. There was also diminished FDG uptake in both caudate nuclei and the cerebellum. The CT scan demonstrated hypodense areas representing encephalomalacic changes both in the putamens as well as in the caudate nuclei. Similar areas were also present in the cerebellum.

Sartorelli, 1966

A 22 year old chemist was exposed to acrylonitrile vapours (distillation) in a small room for two hours. He developed headaches, nausea and dizziness, followed by vomiting, dyspnoea, tremor, incoordination, and convulsions without loss of consciousness. No specific treatment or blood measurements for acrylonitrile or cyanide were performed, the patient recovered fully. *Comment: Alleged cyanide intoxication, symptoms might be due to acrylonitrile, too. No antidotes applied.*

Schulz et al, 1979

A 42 year old man was treated with a combination of SNP and STS. During long-term infusion of SNP the patient developed toxic cyanide levels (3.6 µg/ml erythrocyte cyanide concentration). The patient was treated successfully with STS; erythrocyte cyanide levels decreased to 0.5 µg/ml within 7 hours. *Comment: Effective lowering of intra cellular cyanide level by STS.*

Schulz et al, 1982

Nuremberg Case: This is a short note in a table on the death of a 29 year old female after a dose of 480 mg SNP.

Cologne Case: See below, a total dose of 28 mg SNP.

Schulz and Roth, 1982

A 10-day-old infant received SNP at 2-5 µg/kg for increasing hypertension. After 30 hours, the blood pressure dropped and acidosis developed. An erythrocyte cyanide level of 11.3 mg/l was found, STS applied and SNP infusion suspended. SNP was recommenced after three days as infusion together with STS (100 mg/kg). Cyanide levels remained low. However, two days after SNP/STS were started again, the child died from respiratory failure. *Comment: A case of*

moderate poisoning, STS (and stop of SNP) was effective. Later the combination STS/SNP was given, so that no further cyanide poisoning can be assumed. Reason for respiratory failure is unclear.

Scolnick et al, 1993

Two workers were accidentally exposed to propionitrile fumes from a waste slurry. One worker collapsed after seven hours of exposure, and had to be resuscitated on site. Upon arrival in the hospital he was in coma with eye deviation, and later developed seizures. A case of poisoning with carbon monoxide, cyanide, or hydrogen sulphide was suspected and the cyanide antidote kit with 300 mg SN and 12.5 g STS was applied i.v. Thereafter the seizures stopped and the patient regained conscience. A further treatment with hyperbaric oxygen was started. Forty-eight hours later the patient was dismissed with a resolving pneumonia. Laboratory results arrived later and revealed a cyanide concentration in blood of 5.0 mg/l. The patient complained of severe headaches and dizzy spells over 30 more days. *Comment: Effective treatment with SN/STS in severe poisoning.*

The other worker complained of headache, nausea, and dizziness after two hours exposure. He vomited and then lied down in the cafeteria, where he was found five hours later in a confused and disoriented state. His nausea persisted, as did his headache. The blood cyanide level was 3.5 mg/l. He also received the cyanide antidote kit described above, and was discharged 25 hours later. *Comment: Mild poisoning, in which SN/STS was effective.*

Selden, 1990

A man, 31 years old, took a capsule containing KCN in a suicide attempt. He vomited at once, was unresponsive to external stimulation except pain, and showed bradycardia of 44/min and tachypnoea of 32/min. Twenty-two minutes after ingestion of the poison he showed cyanosis, although he was ventilated with 100% oxygen. The patient was mildly acidotic (pH 7.29) and was behaving himself. He was intubated, ventilated and received AN for inhalation. Moreover, 0.3 g SN and 12.5 g STS were administered twice within four minutes. This caused the MetHb to rise to 9.8% after eight minutes. The maximum MetHb was measured after 90 minutes with 13.4%. A short time after application of the antidotes the man was alert and had to be sedated because of his aggressive behaviour. Before being extubated 75 minutes after admission, gastric lavage was performed. The cyanide levels were determined 30 minutes, 60 minutes, 120 minutes, 3.5 hours and 6 hours after the incident at 5.12 (dissociation between high blood cyanide and mild clinical presentation) – 6.93 – 3.09 – 0.67 and 0.18 mg/l. The man recovered completely. *Comment: Moderate intoxication in spite of high blood cyanide level. AN/SN/STS were effective, though a double dose of SN/STS was given.*

Shragg et al, 1982

A 67 year old woman with colon cancer took 4-5 ground bitter almonds. Thirty to forty-five minutes later she became light-headed and felt nauseous, started vomiting and had abdominal cramps. After the symptoms subsided overnight she crushed and ingested 12 bitter almonds. Fifteen minutes later she had severe abdominal cramps and collapsed. Upon arrival in hospital she was in deep coma and incontinent. As bitter almond smell was detectable she was given AN, SN (0.3 g) and STS (12.5 g), and gastric lavage was done, a cathartic and activated charcoal were given, intubation was done for oxygenation. Twenty minutes after the antidote application the patient became responsive again, however pulmonary oedema developed. She made a full recovery. Later on the cyanide content of one bitter almond was determined as 6.2 mg and cyanide was highly positive in a qualitative check of the gastric contents. *Comment: A case of severe poisoning, in which SN/STS were effective.*

Singh et al, 1989

A 24 year old man was working in a silver processing plant. While cleaning a tank that had contained silver cyanide he fell into it and lay in the remains of the toxic solution for an unknown time. The cyanide concentration in the room was 200 ppm. On arrival in the ED he was in coma, apnoeic and a smell of bitter almonds could be detected. Furthermore, he was tachycardic (120/min), the blood pressure could not be measured, both pupils were fixed and dilated and he did not react to painful stimulus. So intubation was performed and 0.3 Co-EDTA together with 50 ml 50% glucose solution were administered. The antidote was given twice within one minute. The treatment resulted in spontaneous respiration but this only persisted for seven minutes. So a third dose of Co-EDTA was applied, now leading to an oedema of the whole body. Cardiac ischemia was seen in the ECG and the BGA showed a severe acidosis of 7.08. It was only now that the patient was decontaminated, the stomach was washed and he was given i.v. fluids. As vasopressors were continuously administered the systolic blood pressure could be established at 100 mm Hg. Despite all these efforts the young patient died 24 hours later.

During his hospitalisation, different biochemical parameters were monitored and are summarised in the following table:

Time (h) post exposure	0,5	2	4	12	18	24
CN ⁻ (μmol/l) / (mg/l)	804/21.7	819/22.1	608/16.4	23/0.62	15/0.41	15/0.41
SCN ⁻ (μmol/l)	-	147	172	345	281	267
pH	7.21	7.23	7.27	7.33	7.40	7.41
Lactate (mmol/l)	17.5	13.0	5.4	3.8	2.4	2.2

Comment: In this severe cyanide intoxication case the Co-EDTA was not able to rescue the patient though it had been given 3 times. The antidote was not effective and it even prompted adverse effects.

Sipe et al, 2001

A 78 year old woman was involved in a motor vehicle collision. At the scene she had a decreased level of consciousness and was intubated. At the hospital she was started on esmolol and nitroprusside doses in the range of 4.4 $\mu\text{mol/kg/minute}$ to control her blood pressure. The Ativan (lorazepam) infusion was discontinued because of her altered mental status. She received a total of 1300 mg nitroprusside over the five days of infusion. A cyanide level was checked on post-trauma day 4 (2.0 mg/l or 80 $\mu\text{mol/l}$). At this time the nitroprusside infusion was discontinued and 4 mg/kg (10 ml) of 3% SN and 50 ml of 25% STS were administered. The same regimen was repeated nine hours later. She did not improve neurologically. An electro-encephalogram showed theta waves slowing down. The family decided to withdraw support on post-trauma day 20. The patient died on post-trauma day 21. *Comment: It is unclear, whether this was primarily cyanide poisoning. The antidotes given (SN/STS) were not effective, which may well be due to a non-significant cyanide involvement in this case.*

Smith, 1977, 1978

Case 1: A 48 year old woman with lymphoma diagnosed in 1965 began taking laetrile (6 mg i.v. each week and 500 mg tablets orally three times a day) in February 1977. On April 25 she was admitted to hospital with fever, malaise, headache and severe abdominal cramps. She had a diffuse macular erythematous rash, marked lymphadenopathy and hepatosplenomegaly, and abdominal tenderness without peritoneal signs. Laetril therapy was discontinued and after two days her symptoms cleared. Against advice she resumed the laetrile therapy on June 5 and was readmitted on June 20 with the same syndrome. Blood cyanide level was 1mg/dl; skin biopsy was consistent with drug eruption. Symptoms resolved within 48 hours of discontinuing laetrile therapy.

Case 2: A 46 year old man was found to have large cell anaplastic carcinoma of the lung metastatic to the left temporal-parietal area of the brain. In September 1976 he began taking laetrile, 500 mg orally each day. In March 1977, he was admitted with progressive neuromuscular weakness of both lower and upper extremities as well as bilateral ptosis. Laetrile therapy was discontinued; he has not recurrence of the above symptoms.

Comment: Both patients recovered without antidotal treatment.

Steffens et al, 1998; Steffens, 2002

A worker was splashed on the face, neck and chest. Assuming cyanide involvement he received STS (10 ml 10%) twice and was started on an N-acetylcysteine infusion, which caused an allergic reaction and had to be stopped. The maximum acetonitrile concentration in blood in a whole series of biomonitoring measurements was 6.9 mg/l, while cyanide could not be detected at any point in time. Either there had been no cyanide formation, or the cyanide formed with a latency had been efficiently captured by the applied STS. *Comment: No proven cyanide poisoning, so antidote efficacy cannot be assessed.*

A worker was steam cleaning a filter contaminated with propionitrile and thus inhaling it. One hour later he was feeling unwell and then developed nausea and dizziness, followed by somnolence and collapse. During the injection of 20 ml STS 10% he regained consciousness, after which he remained free of symptoms except for an anterograde amnesia for some time. The initial cyanide blood level had been 3.16 mg/l, and it dropped quickly (1.25 hours) by more than 50% after the STS injection, which was once repeated. In parallel the propionitrile concentration in blood dropped from an initial 12.74 mg/l by 90%. Nevertheless a second dose of 20 ml STS 10% was applied. *Comment: Moderate poisoning effectively treated with STS alone. The immediate positive reaction to STS (during the injection) indicates a rapid onset of antidotal effect.*

Steffens, 2003a,b

A 48 year old worker was exposed without protection equipment near a heat exchanger under repair, which was gassing out a.o. acrylonitrile. Approximately 30 minutes later he was vomiting and experiencing somnolence. Upon decontamination, he was given an i.v. application of N-acetylcysteine (150 mg/kg bw) during 15 minutes. Initially he showed increased nausea; then, he recovered. Biomonitoring later showed a low acrylonitrile concentration, but also possible life-threatening cyanide concentration in the whole blood (3.1 mg/l). There was a marked decrease of blood cyanide after application of N-acetylcysteine alone.

Three years later, the same worker accidentally inhaled acrylonitrile. There were few symptoms, only slight nausea. Being aware of the first incident, he was immediately given an i.v. application of the full N-acetylcysteine regimen (a total of 300 mg/kg bw as for paracetamol poisoning), and an additional i.v. application of 10 ml STS 10% approximately 2.5 hours later. The therapy was well tolerated except for skin itching and generalised erythema. Later biomonitoring showed again a potentially critical cyanide concentration in the whole blood (3.4 mg/l). The acrylonitrile level was low again. The marked decrease of the blood cyanide concentration was due to application of N-acetylcysteine alone, well before the injection of STS.

Steffens et al, 2003

The authors described two cases of life-threatening hydrocyanic acid (HCN) gas poisoning. In at least one case the uptake was strictly percutaneous, as a self-contained breathing apparatus had been used. A stream of gas containing 19% HCN was released from a leaking flange. One worker collapsed and was treated on site with 2 ml DMAP. At the same moment a fire fighter wearing a self-contained breathing apparatus collapsed, was unconscious and had foam on his mouth and nose. He also received 3 ml DMAP immediately. Upon arrival in the emergency room the cyanide levels were 5.3 and 6.75 mg/l respectively in the whole blood (after the application of DMAP). MetHb was 5.5 and 16.2% respectively. After decontamination both patients received 50 and 150 mg/kg bw N-acetylcysteine respectively, as parallel acrylonitrile poisoning had to be suspected. In addition they were given 30 and 40 ml STS 10% respectively. In both cases the cyanide levels dropped significantly to 0.035 and 0.036 mg/l on the following day, and both patients survived without sequelae. Conclusions: HCN gas can easily penetrate the skin and lead to possible fatal cyanide poisoning. If work in contaminated areas is necessary, a chemical protection suit has to be used, normal fire fighting equipment has been shown to be insufficient protection. The immediate application of DMAP at about 50% of the recommended dose has been shown to be life-saving, even though MetHb levels remained relatively low. The high cyanide levels after DMAP treatment show that there is no definitely lethal cyanide concentration. It might be possible, that the rapid increase of cyanide levels may play a greater role. *Comment: The application of N-acetylcysteine plus STS seems to be an effective antidote treatment. Gaseous HCN was absorbed through intact skin.*

Steffens, 2010

Two cases of life-threatening hydrocyanic acid (HCN) gas poisoning when a stream of gas containing 19% of HCN was released from a leaking flange.

Case 1 : A worker collapsed and was treated on site with 2ml 4-dimethylaminophenol (4-DMAP). Upon arrival in the emergency room the cyanide level was 5.3/l in whole blood (after the application of 4-DMAP) and methemoglobin was 5.5%. After decontamination he received 50 and 150 mg/kg b.w. N-acetylcysteine (NAC) respectively, as parallel acrylonitrile poisoning had to be suspected. In addition 30 and 40mL sodium thiosulfate 10% respectively were given. Cyanide levels dropped significantly to 0.035 on the following day, and the patient survived without sequelae.

Case 2 : A fire fighter wearing SCBA in attendance collapsed, was unconscious and had foam in front of his mouth and nose. He also immediately received 3ml 4-DMPA. Upon arrival in the emergency room his blood cyanide level was 6.75 mg/l in whole blood (after the application of 4-DMAP) and Methemoglobin was 5.5%. After decontamination he received 50 and 150 mg/kg

b.w. N-acetylcysteine (NAC) respectively, as parallel acrylonitrile poisoning had to be suspected. In addition 30 and 40mL sodium thiosulfate 10% respectively were given. Cyanide levels dropped significantly to 0.036 mg/l on the following day, and the patient survived without sequelae.

Comment: HCN gas can easily penetrate the skin and give rise to possibly fatal cyanide poisonings as in at least one case the uptake was strictly percutaneously, as a self-contained breathing apparatus (SCBA) had been used. The immediate application of 4-DMAP at about 50% of the recommended dose has been shown to be life-saving, even though methemoglobin levels remained relatively low. The high cyanide levels after 4-DMAP treatment show, that there is no definitely lethal cyanide concentration. It might be possible, that the rapidity of increase of the cyanide levels may play a greater role. The application of NAC plus sodium thiosulfate seems to be an efficient treatment of cyanide poisonings and should be evaluated further.

Stickel, 2008 (Zilker and Eyer, 2005; Zilker and Schweizer, 1987)

Stickel describes eight moderate and severe intoxications with HCN. All were treated with DMAP. Five of the persons survived with DMAP therapy, three died despite being treated with the antidote. Zilker and Eyer (2005) reported the same cases in a historical overview.

Case 1: A 34 year old analytical chemist had taken KCN in a suicide attempt. Twenty minutes later he was found comatose with loss of peripheral muscle tonus. He was then intubated, ventilated and given sodium bicarbonate, Ringer's solution and, as specific cyanide antidotes, 0.25 g DMAP together with 10 g STS. Ten minutes later he was alert and responsive again. After gastric lavage the patient was taken to hospital, where ventilation was continued, the gastric lavage was repeated and a second dose of DMAP was administered. The blood cyanide level was 2.4 mg/l two hours after ingestion. The MetHb reached 37.7% within five minutes after the second antidotal treatment. The patient developed mild haemolysis (bilirubin 4.7 mg/dl and Hb decrease from 13.3 g/dl to 10.3 g/dl) as side effect of the DMAP-application. Altogether he made an uneventful recovery and could be discharged a few days later. This case is also published by Zilker and Schweizer (1987). *Comment: Moderate intoxication with efficacy of DMAP and STS applied. Side effects (haemolysis) were probably due to the second dose, the need for which is unclear.*

Case 3: A 28 year old female chemical laboratory assistant committed suicide with cyanide. After 10 minutes she was found in a coma, with tachycardia, tachypnoea, but satisfying circulation. She was intubated, ventilated and 500 mg DMAP together with 25 g STS were injected. Furthermore she was given sodium bicarbonate and sedatives. Two hours later she was admitted to hospital being in a non-reactive coma. Nevertheless she regained consciousness a

few days later (despite severe haemolysis with bilirubin of 9.3 mg/dl, Hb fall from 13.6 mg/dl to 9.0 mg/dl and LDH maximum of 648 U/l) and recovered completely. The blood cyanide levels were 25, 31 and 0.6 mg/l 20 minutes, 2 and 11 hours after poisoning, respectively. The MetHb-level was measured 110 minutes after the antidotal treatment and was still within the therapeutic level (33.4%). *Comment: **Severe poisoning** with very high cyanide blood level. Recovery took several days, so it is not fully clear, whether this was due to DMAP and STS. The high dose of DMAP given prompted adverse effects.*

Case 4: A 39 year old goldsmith ingested 5 g of KCN. Ten minutes later he lost consciousness and became cyanotic with gasping respiration and tachycardia. The circulation remained quite stable. The ambulance staff intubated and ventilated the patient and administered 250 mg DMAP in combination with 10 g STS (the latter was given twice). This treatment resulted in spontaneous respiration, sweating and hyper-salivation. One hour after poisoning the blood cyanide level reached its maximum value of 1.46 mg/l. The patient remained in a coma for one day and on the respirator for two more days. Although he developed severe haemolysis (Hb decreased from 15.8 mg/dl to 9.3 mg/dl) he recovered completely. *Comment: **Moderate intoxication**, the effectiveness of DMAP and STS is not clear, as recovery took relatively long. Significant adverse effect of DMAP at normal dose.*

Case 5: A 56 year old goldsmith had inhaled NaCN gas while at work. Only a few minutes later he collapsed and exhibited convulsions. His circulation was stable and he breathed spontaneously, but with difficulties. The patient was intubated and 1,000 mg of DMAP (i.e. 4 times the recommended dose) were administered, together with 25 g of STS. This resulted in severe cyanosis with a MetHb-level of 73% within 90 minutes after the antidotal treatment. Therefore blood was transfused and the amount of MetHb could be reduced to 46.3% within two hours and 20 minutes after the overuse of DMAP (33.8% after three hours and 40 minutes). The MetHb level further decreased to 32.6, 21.6 and 2%, after 3, 4 and 5 days post exposure, respectively. The cyanide level in blood was determined once (at admission) with 37 mg/l. The overdosage of DMAP caused massive haemolysis (bilirubin 8.9 mg/dl, Hb decrease from 13.9 to 7.3 g/dl, LDH 6600 U/l), and intermittent renal failure. In the end the patient recovered completely. *Comment: **Severe cyanide intoxication** could be rescued with DMAP (and STS), which thus were effective. The overdosage of this antidote resulted in serious side effects (e.g. changes in the blood count, and renal failure). The blood cyanide level was extremely high (37 mg/dl), perhaps because of the cyanide bound to MetHb. No explanation is given for the administration of 1,000 mg of antidote.*

Case 6: A 17 year old schoolboy ingested KCN in a suicide attempt. Within 20 minutes he developed cardio-respiratory arrest (the cyanide level in the blood sample counted 34 mg/l but it was available after a few hours only). The ambulance staff did not have any cyanide antidote with them so 750 mg DMAP and 17 g STS could only be administered 1.5 hours later

(intermediate actions not described). Circulation stabilised straight after this treatment, but no EEG activity could be detected. The patient developed cerebral oedema, diffuse cerebral bleedings and acute renal failure. He died five days after the intoxication because of circulatory failure. *Comment: Severe intoxication with antidotal treatment being established too late and thus remaining ineffective. Multi-organ failure had already been initiated. Also the DMAP dose was very high.*

Case 7: A woman (aged 36) had ingested KCN. It took 90 minutes before the diagnosis of cyanide intoxication was established. During this time the treatment consisted of supportive actions like intubation, ventilation and application of catecholamines, as the patient was in a coma, apnoeic and showed circulatory failure. After being diagnosed correctly the woman received 250 mg DMAP and 10 g STS. This resulted in sufficient circulation (blood pressure 150/100 mm Hg, tachycardia), but no EEG activity could be reconstituted and she died five days later. The cyanide levels were 10.96 mg/l two hours later, and approximately 1 mg/l 11 and 24 hours post exposure. The MetHb-level reached 19% within 15 minutes after the antidotal treatment. *Comment: Severe poisoning, late initiation of antidotal treatment, which improved circulatory, but not neurological status and overall was only partially effective – probably due to treatment delay.*

Case 8: A 40 year old woman collapsed after cyanide ingestion. The ambulance arrived 45 minutes later and found a comatose patient in cardiac arrest. She was intubated, ventilated and received 250 mg DMAP and 10 g STS. This led to satisfying circulation, but a few hours later irreversible ventricular fibrillations occurred and the patient died. The cyanide level in blood was 14 mg/l and MetHb 14.8%, both 45 minutes after the antidotal treatment. *Comment: Severe intoxication with very high cyanide level. DMAP and STS were only temporarily effective to improve circulation.*

In conclusion, Stickel reports on eight cases of cyanide poisoning, all treated with DMAP. The antidote worked well for five patients (moderately and severely poisoned ones). But three patients with severe intoxications died as the antidote was not effective at all. Mild haemolysis was regularly seen, even under normal dosage; nevertheless this adverse effect could have been easily managed. The MetHb level became life threatening for one patient who had been given an antidote dose that was too high. However this could be treated and the patient survived without sequelae. The case reports show that DMAP is an effective antidote even in severe cases. It must be admitted that after cardiac arrest had taken place, DMAP was not as effective as in patients whose circulation was stable. Under the recommended dose the adverse effects are mild and can be managed without any problems. Severe side effects normally occur only if DMAP is overdosed. All patients should receive STS to support the cyanide elimination.

Suchard et al, 1998

A 41 year old woman ingested about 30 chewed apricot kernels as an alleged health food. Twenty minutes later weakness developed, followed by dyspnoea; the paramedics found her in a deep coma and unresponsive to painful stimuli. Cyanide poisoning was assumed and later confirmed, the application of AN, SN and STS (Eli Lilly Kit) was successful and led to immediate improvement. As four hours later cyanide in blood was still or again about 1 mg/l, a second half dose of the antidote kit (SN and STS) was given. After the first dose MetHb was 7.3%, and 10.5% after the second dose. *Comment: In this severe case of poisoning AN/SN/STS were fully effective.*

Sutter et al, 2010

A 34 year old female, suspected to have cyanide poisoning because of decreased oxygen extraction from arterial to venous side and profound metabolic acidosis, had a cardiac arrest and was resuscitated. Under the suspicion of cyanide poisoning HOCO 5 g followed by STS 12.5 g were given. This led to rapid improvement, although a dramatically elevated lactic acid level persisted. Haemodialysis was attempted, but could not be performed due to false 'blood leak' alarm of the machine, which was caused by the plasma discoloration from HOCO. *Comment: HOCO and STS were effective in this severe case.*

Tartière et al, 1992

After ingestion of KCN, a 32 year old man was found sweating and with convulsion-like movements, though awake. Oxygen was given; then 5 g HOCO as infusion for over three hours, in spite of circulatory stability and only slight somnolence. He fully recovered after the treatment. *Comment: In this mild poisoning case, HOCO led to recovery and was effective.*

Tassan et al, 1990

A 15 year old girl tried to kill herself by intake of 2.5 g KCN. One and a half hour later she presented comatose, with hemodynamic instability and severe metabolic acidosis. In the ED gastric lavage was performed, mechanical ventilation initiated and sodium bicarbonate and beta-mimetics administered. She received 4 g of HOCO as specific cyanide antidote and 8 g of STS. Upon admission, the cyanide level was 494 $\mu\text{mol/l}$ (13.3 mg/l, high). The girl recovered completely, despite of the delay in the therapeutic intervention and the severity of the intoxication. *Comment: Very high cyanide level, yet full recovery after HOCO/STS in this severe case of poisoning.*

Thiess and Hey, 1969

A further case of acetone cyanohydrin poisoning has been reported in a 23 year old worker filling acetone cyanohydrin from a rail tanker into steel drums. One of his cotton gloves was drenched, which he did not notice. He put the glove into his pocket. Five minutes later nausea set in, vomiting occurred, and ten minutes afterwards, he lost consciousness and collapsed. AN and artificial respiration led to short improvement, but then coma and convulsions set in. The use of SN and STS also improved the status only for minutes. After the glove had been found, clothing was changed, and a decontamination bath was performed. *Comment: SN and STS led to an immediate improvement and were effective. Cyanide was not measured.*

Thomas and Brooks, 1970

A 19 year old boy who was working in a photography shop had his face contaminated with KCN powder. A short time later, he complained of stomach cramps, was pale and lost consciousness. Twenty minutes after the incident he arrived at the ED. Once there his blood pressure went down to only 90/70 mm Hg. His stomach was washed and he was given the nitrite antidote kit (AN pearls, SN 300 mg and STS 12.5 g). After the antidotes had been administered, he quickly regained consciousness. As pain in his stomach, agitation and a weakness of the upper extremities persisted, the therapy was repeated two hours later (half the initial dose) and again ten hours afterwards (whole dose because of a new opisthotonos). The patient recovered completely. *Comment: This was moderate cyanide poisoning effectively treated with AN/SN/STS. Whether the third dose had really been necessary might be discussed.*

Townsend and Boni, 1975

A 34 year old man ingested 48 apricot kernels, both raw and in a milk shake, and developed vomiting, headache, flushing, perspiration, dizziness and faintness. After induced vomiting he recovered quickly. *Comment: No cyanide measurements were done, no antidotes were applied.*

Trapp, 1970

Trapp reported the case of a 30 year old male who crouched in a tank full of silver cyanide to escape police after a robbery. He fell and immediately lost consciousness with seizures despite AN and i.v. STS. He received one session of hyperbaric oxygen at 3 ATA which resulted in prompt woke up with a good urine flow and correction of metabolic acidosis with IV sodium bicarbonate. Therefore, the pressure was slowly reduced to 2 ATA. The pH was measured during the hyperbaric oxygen therapy at 7.2. He remained alert, volunteered his name, and said he had

swallowed ‘gallons’ of tank solution. He was discharged without sequelae but with burning sensation of his right eye and scrotum. *Comment: A combination of oral-dermal-inhalation intoxication can be assumed. Cyanide levels in the blood were not measured, clinical data are incomplete. AN/STS were not effective in this case, but the doses were not given.*

Turchen and Manoguerra, 1989; Turchen et al, 1991

A 39 year old woman ingested 59 ml of Super Nail Off (containing 99% acetonitrile) in a suicide attempt. Admission vital signs were blood pressure, 108/60 mm Hg; 124 beats/min; respiration 28 breaths/min; compensated metabolic acidosis with a normal arterial pH and a base deficit of 17. The whole blood cyanide level was 313 µg/dl and the ethanol level 150 mg/dl. The patient underwent gastric lavage. Following a latent period of approximately 12 hours, the patient developed cyanide poisoning with severe metabolic acidosis, seizures, and shallow respirations. She responded to the administration of SN (0.3 g) and STS (12.5 g), although the administration of nitrite produced bradycardia and hypotension. She developed several relapses over the course of her hospitalisation and each time responded to STS administration (at 26.5 hours 12.5 g STS, at 34.5 hours 0.3 g SN plus 12.5 g STS were given i.v.). Nine hours later she was given another 12.5 g STS with a rapid response. Her cyanide level peaked 59 hours post-ingestion at 1.281 mg/dl. Laboratory tests showed a continuing metabolic acidosis and mild hypernatraemia. She was tachypnoeic with a respiration rate of 28 breaths/min. Sixty-five hours after ingestion, her cyanide level was 1049 µg/dl and thiocyanate was 12 µg/dl. Despite persistently high cyanide levels her vital signs improved on the fourth day. She was discharged the morning of her sixth hospital day. *Comment: The delayed onset of symptoms is due to delayed cyanide formation from acetonitrile. The administration of STS was more effective than SN which exacerbated the patient’s hypotension as side effect.*

Uitti, 1985

An 18 year old man ingested 975 to 1,300 mg of KCN in a suicide attempt. In the ED he was comatose, not responding to painful stimuli. Blood pressure was 160/70 mm Hg, and pulse was 108/min. He was treated with gastric lavage, endotracheal oxygen, and i.v. SN and STS. He regained consciousness in 7 hours, but he had severe Parkinson syndrome, characterised. He died 18 months later after he had taken an overdose of imipramine and alcohol. *Comment: In this severe poisoning SN/STS were effective with some delay. The reason for the Parkinson syndrome is not clear.*

Valenzuela et al, 1992

A 16 year old man ingested 1 g KCN. On admission he was comatos, his blood pressure was undetectable, and he had deep respiratory depression. His heart rate was 70 beats/min. Besides cardiopulmonary resuscitation, he received i.v. STS. Three hours later he was fully conscious. After 24 hours he was neurologically intact and the results of general physical examination were normal. A few days later he developed a severe dystonia syndrome. He had a positive response to an apomorphine test and showed improvement with levodopa treatment. A 21-year follow-up showed minimal neurological sequelae; CT showed bilateral putaminal lucencies. *Comment: The administration of STS was successful, but a delayed severe dystonia syndrome occurred.*

Zaknum (2005) + Rachinger (2002), 2005

A 35 year old female laboratory assistant attempted suicide by taking an unknown amount of cyanide. 10 minutes later she was found comatose with stable cardiovascular function and was immediately given emergency treatment. Under controlled respiration she was administered DMAP and STS, which resulted in a linkage with cyanide, and antioedema treatment was administered. After withdrawal of sedatives and respirator she exhibited agitation and akinetic mutism. A dopamine agonist (amantadine sulphate) was administered i.v. and later orally. She received extensive physio-, logo- and ergotherapy but CSN necrosis and sequelae persisted.

van Dijk et al, 1987 / van Heijst et al, 1987

A 21 year old male was found deeply comatose. His sister found NaCN pellets nearby and called for help. He arrived at hospital about 1 hour after ingestion. He was cyanotic and deeply comatose with bradycardia and first-degree AV block, which descended into cardiac arrest. He was resuscitated, but remained cyanotic. He received STS 20 g, then about 1 hour later received 300 mg SN. Fifty minutes after dosing, his MetHb was 'only' 3.6%, so he received 250 mg 4-dimethylaminophenol (DMAP). Twenty-five minutes later he was so deeply brown-blue cyanosed that the physician administered empirically 160 mg toluidine blue. A MetHb obtained shortly thereafter revealed 69% methaemoglobinemia. Toluidine blue 250 mg was given and the MetHb dropped to 46%, then later to 4.4%. Because this MetHb level was considered too low, DMAP was injected again, the physician this time giving only 80 mg, which resulted in a MetHb of 52%. Over the next 10 hours, DMAP was administered 5 times with close monitoring of MetHb. During the whole treatment, the patient remained deeply comatose. Thirty hours after admission, ventricular fibrillation occurred and he died.

van Heijst et al, 1987 / Douze et al, 1987

All the patients reported by van Heijst received combinations of different antidotes together with HOCO in a sub-therapeutic dose. The low doses of HOCO applied in these cases are not considered to be usable for assessment.

Case 1: A 32 year old male who attempted suicide by ingesting 3 g of KCN. He was found unconscious with inadequate respirations, receiving mouth to mouth respiration. He was extremely bradypnoeic on admission (respiration rate = 2), with heart rate 30 beats/min and blood pressure 110/80 mm Hg. A smell of bitter almonds led clinicians to administer 10 ml SN and 50 ml STS. Convulsions were treated with diazepam and acidosis with sodium bicarbonate. Because he remained unconscious, the treatment with SN and STS was repeated. One hour after admission, he awoke, complaining of shortness of breath. His skin colour was brownish blue. The MetHb was 58%. He received methylene blue 30 ml of a 1% solution. He was transferred the next day from the ICU to psychiatry in good physical condition. *Comment: No blood levels determined, but serious clinical presentation this seems to have been moderate poisoning. SN/STS at double dose was effective, but caused high MetHb levels.*

Case 2: A 23 year old male student took KCN. The dose could not exactly be determined. In addition he tried to inhale some HCN-gas by heating KCN. He was brought to hospital approximately 20 minutes after the event. On admission the patient was semi-comatose and in convulsions. The heart rate was 100/min, the blood pressure 120/60 mm Hg, the respiratory frequency was normal. He was given 20 mg of HOCO i.v. followed by intubation and artificial ventilation with 100% oxygen. 100 ml of 25% STS were administered i.v. As the dose of HOCO was insufficient DMAP was administered i.v. By mistake 1,000 mg were given. The MetHb rose to 70%. This was treated by toluidine blue. Three hours later MetHb came down to 25%. The patient developed jaundice caused by massive haemolysis. The highest bilirubin was 7.2 mg/dl, the lowest Hb was 7.7 g/dl. The anaemia was treated with blood transfusions. The patient survived. *Comment: Patient treated with DMAP as main antidote (highest dose).*

Vogel et al, 1984

A 24 year old officer on a cargo ship was sprayed with acrylonitrile, when a valve burst. He had dizziness, flushing, nausea and vomiting and soon began to hallucinate. The cyanide level was 0.13 mg/l upon admission and reached a maximum of 0.184 mg/l one day later. The patient was treated with 15 (!) doses of the “cyanide kit”, each containing 10 ml 3% SN and 50 ml 25% STS solutions, over 72 hours. Even though the highest MetHb level reached was 14.5%. When the level fell below 10%, the patient started hallucinating again, convulsions occurred once. Additionally oxygen and HOCO were given at an unspecified dose. Extra-corporal dialysis was attempted influencing the hallucination ascribed to cyanide. Later the laboratory values indicated

rhabdomyolysis, which was treated by hydration. The patient made a full recovery. *Comment: The measured cyanide levels were insignificant and do not explain even the mild to moderate course. This is in line with the inefficacy of the cyanide antidotes. It can be disputed, whether this was acrylonitrile, and not cyanide poisoning.*

Vogel, 1991

A chemist, 30 years old, tried to commit suicide with KCN which he had taken from his lab. He was seen in the emergency department in a comatose state, with a tachycardia of 120/min and with gasping respiration (20/min). He was unresponsive to painful stimuli and the corneal reflex could not be triggered. The anion gap was 30 mmol/l and the base excess at 17.8 mmol/l. The remaining biochemical parameters were within the normal range. He was intubated, ventilated and gastric lavage was performed. Moreover he got activated charcoal and forced diuresis was induced. No specific antidotal treatment was applied. Nevertheless the patient recovered with supportive care only. He became alert three hours later and could be extubated. Six hours after the suicide attempt he admitted the intake of cyanide and the blood cyanide level was then measured (0.23 mg/l but that is not valuable because of the long delay). *Comment: Obviously a severe poisoning, survival with supportive treatment only.*

Wananukul and Kaojarern, 1992

A 19 year old male accidentally ingested KCN at an electroplating shop. He vomited shortly after, turned red, had a generalised seizure and became unconscious. In the ED he was stuporous, with shallow respirations. Initial blood cyanide concentration was 12 mg/l. He received 10 ml of SN some 3.5 hours after admission then, STS. He was described as having fully recovered six hours after treatment. *Comment: This moderate poisoning was effectively treated with SN/STS.*

Weng et al, 2004

A man, aged 51, took an uncertain amount of KCN solved in a milk-emulsion (3 g of KCN per litre milk) in a suicide attempt. Twenty minutes later he had become non-reactive, collapsed and lost consciousness. He developed convulsions and stayed in coma. The blood pressure was 77/60 mm Hg one and a half hour after the ingestion of the poison. He was tachycardic (147/min) and exhibited 7 points on the GCS. The blood gases showed a metabolic acidosis (pH 7.22) with a lactic acid of 12 mmol/l. The cyanide level in the blood was 2.9 mg/l. The patient was intubated, ventilated and was given activated charcoal. It took 2.5 hours before 5 g of HOCO were administered i.v. The blood cyanide level increased to 4.2 mg/l one hour after the antidote application. The patient's clinical conditions improved step by step and three hours later

he could be extubated and he could be discharged after five days in complete remission. *Comment: This is a severe poisoning case effectively treated with HOCO, even though this was applied late, and though the cyanide level initially increased after antidote application.*

Werner, 1979

This is a dissertation on the treatment of poisoning with hydrocyanic acid, its salts and derivatives with 4 DMAP. All cases included in the dissertation are included in the publication by Zilker and Eyer, 2005.

A 30 year old woman ate 35 pieces of bitter almonds as she wanted to commit suicide. She felt sick and dizzy. The vital parameters were stable as was the consciousness. She was administered 250 mg DMAP and 10g STS and recovered completely. *Comment: A very mild poisoning not requiring treatment. So the antidote efficacy/effectiveness cannot be assessed.*

The second case reports on another woman, aged 38, who had ingested bitter almonds. She complained about dizziness, nausea and a sore throat. She smelled like bitter almonds. The blood pressure was 115/70 mm Hg, HF 100/min. She was administered 500 mg DMAP together with 15 g STS. The treatment caused severe haemolysis and the patient reported a headache. The side effects could be treated within a few days. *Comment: Mild poisoning, no antidote indication, so that efficacy/effectiveness cannot be evaluated. Instead significant adverse effects occurred at double dose.*

Wesseon, 1985

A 59 year old scientist was found comatose with a suicide note thereby. He was given fluids i.v. and naloxone. When coming to the emergency department he showed spontaneous respiration, the pupils were reactive to light but reflexes could not be triggered. The blood pressure was 140/80 mm Hg, the heart rate tachycardic (136/min) and the respiration fast (32/min). The pH decreased from 7.13 at admission to 6.99 after half an hour, despite bicarbonate infusion. The lactate was very high with 29.2 mmol/l (but it seemed to be relatively well compensated). He was intubated, the stomach was washed and activated charcoal was administered. At this time the poison was not known so a haemodialysis was performed (also for treating the acidosis). After some time a friend of the patient's suspected cyanide intoxication and the nitrite antidote kit was then applied (SN 300 mg and STS 12.5 g). The man became alert after about five hours and made an uneventful recovery. The cyanide level was determined at the beginning (0.4 mg/l), shortly before the antidotes were given (2.1 mg/l) and at the end of the dialysis (0.1 mg/l). *Comment: In this (probably) moderate poisoning SN/STS were effective in spite of late application.*

Wolfsie, 1951

Wolfsie reports on 12 cases that had been accidentally poisoned by cyanide either via the respiratory or via the dermal route. All were male workers between 29 and 48 years. Two patients had suffered from mild intoxications without loss of consciousness and ten from moderate poisonings with coma but sufficient circulation. The mild and four of the moderate intoxications were treated with AN, the other six with SN (0.3 g) and STS (12.5 g). Three of these patients received the antidote only after spontaneous clinical recovery. All patients were ventilated and decontaminated when the exposure route had been the dermal one. The men recovered completely. *Comment: 12 mild or moderate intoxications were treated with nitrites, which showed good efficacy. Diagnosis by the patients' colleagues might not always have been reliable. In at least in 3 cases antidote (given after recovery) were not required.*

Wright and Vesey, 1986

A 42 year old woman drank 100 ml of a potassium-gold-cyanide-solution ($\text{KAu}(\text{CN})_2$, corresponding to 540 mg cyanide and 2.0 g gold) in a suicide attempt. She vomited at once, but did not go to the hospital before two hours had elapsed. Once there she was responsive, with a heart rate of 110/min. Gastric lavage was performed as first line therapy. Two hours later the patient became drowsy, tachypnoeic and hypotensive (blood pressure 90/60 mm Hg), oxygen and 2 ampoules of AN were given per inhalation. As specific antidote 300 mg of Co-EDTA together with 4 g of glucose in 20 ml were infused. She reacted anaphylactic to this treatment in the form of a flush and peri-orbital oedema. In the further course the patient developed more and more hypotonia and tachycardia as well as she became anuric. Steroids, adrenaline, antihistaminics, i.v. fluids and oxygen were administered, however, her clinical conditions deteriorated. She had to be intubated 9.5 hours after the intake of the poison (five hours after antidotal treatment). She died 3.5 hours later because of cardiac arrest. The blood gases were determined as well as the cyanide levels in the plasma and in the erythrocytes (Table below).

Blood measurements by Wright and Vesey, 1986

Time (h)	Event/parameter	Remark						
16.30	Ingestion of KAu(CN) ₂							
19.00	Admission to hospital							
21.00	Administration of antidote							
			pH	Base excess (mmol/l)	Blood CN (μmol/l)		Blood CN (mg/l)	
					Plasma	Erythrocytes	Plasma	Erythrocytes
21.35					2.1	188.5	0.056	5.1
22.05	7.30	−6.7						
23.50					0.5	42.9	0.013	1.2 ^a
1.00.	7.30	−11.2						
2.30	7.15	−17.0						
2.40					0.14	5.5	0.004	0.15
3.10	7.20	−10.5						
5.15	7.18	−8.7						

^a The gold level was 14.2 mg/l

Comment: Initially the intoxication did not seem to be very serious, but in spite of antidote application (AN and Co-EDTA) it had a fatal outcome. This could have been because of the delayed treatment, or a combination of gold and cyanide intoxication and the patient died most likely due to the gold poisoning for which there is no real antidote. The cyanide antidote Co-EDTA caused severe side effects in the woman, although it had been administered in the recommended dose.

Wu et al, 2001

A 27 year old attempted to commit suicide by ingesting 5 ml gold KCN solution ($\text{KAu}(\text{CN})_2$). He developed vomiting and abdominal pain three hours later. At the hospital gastric lavage and activated charcoal was administered. He was referred to another hospital (based on the suspicion of cyanide intoxication). On arrival, he was conscious and complained of vomiting and showed hyperamylasaemia, and hepatic dysfunction. The blood cyanide test was reported as undetectable (<0.1 mg/dl). A cyanide antidote was not given. The authors mentioned that aurocyanide is extremely stable; so that very little cyanide should be liberated (formation constant is 10×10^{38}).
Comment: Not a significant cyanide poisoning and no antidotes required.

Würzburg, 1996

Thirty-seven employees were exposed to cyanide between 1956 and 1985. One was found dead. Thirty-six employees were treated; most were given nitrite and oxygen. Some received oxygen alone. All recovered completely. One-third of these employees were unconscious. One was convulsing. Most were discharged home at 6 h post-exposure. Some employees remained at the plant to work an additional shift. AN and/or oxygen were the only agents used with 33 employees. Forced oxygen was administered to the unconscious, apnoeic employees. Three employees were given SN and STS i.v. Treatment generally began within 3 min. In 5 to 20 minutes all of the unconscious employees reacted positively to the use of forced oxygen and forced AN or SN. There were no residual effects except headache and transient loss of appetite. AN and oxygen have been effective tools in the treatment of cyanide intoxication at this plant site. There have been no intercurrent or residual drug effects that outweigh the life-saving capacity of these agents. SN should be employed if the use of oxygen and AN fails to improve the cardiovascular status/level of consciousness in 5-10 min.

Yacoub et al, 1974

Case 2: A chemical analyst (50 years old) who was working in an electro-plating company tried to commit suicide by taking approximately 1 g of KCN. He complained about stomach pain immediately and he smelt of bitter almonds. His sons took him to the ED. There he presented somnolent and cyanotic about 30 minutes after the ingestion. He had vomited more than once, was shivering and showed a both-sided miosis (*in contrast to the mydriasis generally seen in cyanide poisoning*). The blood pressure was 90/60 mm Hg and the heart rate 140/min. He was acidotic (pH 7.25) and the cyanide level in the blood was 15 mg/l for the free ion and 42 mg/l for the whole blood cyanide (very high for comparatively mild symptoms). After gastric lavage 50 ml of a 10% STS solution was instilled, parallel he was treated with 2 pearls of AN per inhalation, glucose and sodium bicarbonate i.v. As cyanide antidotes he received 0.3 g of Co-EDTA and 2 g of STS, followed by 4 g of HOCO. Within a few minutes he developed an allergic reaction (urticaria, Quinke's oedema) that could be treated with steroids and antihistaminics. He showed a pink coloured skin because of the HOCO-infusion. Five hours and a half after the incident the patient was conscious, a bit drowsy and tachypnoeic (respiration frequency 24/min). He could be discharged at the fifth day in complete remission. The free blood cyanide level before treatment with antidotes were 15 mg/l (total 42 mg/l) and after treatment (second day) 0.18 mg/l (total 50.8 mg/l). *Comment: The blood-cyanide level was extremely high, maybe erroneously high in comparison to the relatively mild clinical presentation. The need for combined antidotal treatment with AN, Co-EDTA and HOCO is not obvious. Significant side effects (allergy) were seen, which can have been due both to CO-EDTA and especially HOCO.*

Case 1: M. S. Lorenzo, aged 27, working in a factory using cyanide, attempted to commit suicide by ingesting an unknown quantity of potassium cyanide on 14 June 1973. He was driven to hospital and taken to the intensive care unit. Upon arrival the patient was in cardiac arrest with bilateral mydriasis, sweating and cyanosis. Resuscitation attempts, with external cardiac massage, intubation and assisted ventilation, administration of isoprenaline and THAM proved ineffective. Electrocardiogram confirmed the existence of a cardiac arrest that remained irreversible. An anatomical examination performed 48 hours later shows the existence of a strong odor of cyanide at the opening of the body, a massive congestion of all organs, "buff" looking liver, the presence of liquid in the bronchi and hemorrhagic staining of the gastric mucosa.

Yatziv and Simcha, 1969

In Israel a two year old girl ingested a large amount of apricot pits. A few minutes later the first symptoms appeared, and she quickly became unconscious. This coma lasted for 45 minutes. Gastric lavage and oxygen led to rapid improvement, and the girl recovered fully within eight hours. *Comment: No antidote was given.*

Yeh et al, 1992

A 75-year old non-smoking monk was admitted to the hospital with altered mental status and apparent seizure activity. He had no significant past history, but he did possess Laetrile. Prior to admission, the patient had been alert and weak, with abdominal discomfort. An episode of emesis was followed one hour later by coma. On admission the patient had a tremor and was unresponsive to painful stimuli. Vital signs were pulse 114/min with atrial fibrillation, respirations 28/min, blood pressure 157/67 mm Hg. The pupils were equal with diminished reactivity to light. Both gag and deep tendon reflexes were decreased. The admission arterial blood gas value on 6 litres nasal oxygen/min was pH 7.31, pCO₂ 16 mm Hg, pO₂ 154 mm Hg, oxygen saturation 99%. Initial therapy was lorazepam, later a Swan-Ganz catheter was placed and administration of dopamine and dobutamine was started. Seven hours later, the patient became alert and revealed that he had ingested 10 Laetrile tablets. He received 25g of sodium thiosulphate and gradually improved, and was discharged within 5 days. Subsequently, his cyanide level on admission was found to have been markedly elevated at 180 micromol/L. *Comment: The administration of thiosulphate was successful.*

Yen et al (1995)

21 cases of cyanide poisoning were identified at the Taipei Veterans General Hospital from 1985 to 1992. No individual case details are provided. This group of 21 was selected using specific

criteria from among 138 cases of cyanide poisoning seen in the hospital. Fifty-three of the 138 received the cyanide antidote kit, of whom 38 (72%) survived. Among the 21 selected cases, 5 were identified as having no or mild effect. The remaining 16 were identified as having severe effects. Two of 5 in group 1 received the cyanide kit. Apparently, all but two in Group 2 received the kit. Six patients died in that group in spite of the kit for an overall survival in selected patients of 71%. Two groups are differentiated. 5 exhibited mild effects two of which got antidote treatment with the nitrite antidote kit. All survived and would have probably survived without any antidotal treatment. 16 cases were graded as severe. 15 of them had lost consciousness, showed respiratory suppression, 4 apnoea, 5 bradycardia, 8 had hypotension, 4 a cardiovascular collapse, 13 metabolic acidosis. 8 of the survivors had antidotal treatment, 6 of the none-survivors had antidote treatment, but two had no such therapy. The survival rate in the treatment group was 57%. The survival rate in the group which did not get the antidote was zero if only the severe cases are taken into consideration. *Comment: The study points to a positive effect of the treatment with the nitrite antidote, but the number of severe cases which had no treatment and died was very small (2).*

Zaknum, 1905

A 35 year old female technician ingested a lethal dose of KCN in a suicide attempt at a hospital's laboratory. DMAP and STS were promptly administered. On admission to the ICU 15 minutes after intoxication she was in coma but pupils were not dilated, blood pressure was 120/70, heart rate of 110/min, mixed crepitate rales on the base of the lungs were reported. Following artificial coma she presented with an agitative state for several days followed by akinetic mutism, buccofacial and ideomotoric aphasia. Severe rigid-akinetic syndrome, dysarthria, dysphagia and generalized dystonia developed weeks later. MRI revealed lesions in the caudate and lentiform nuclei, precentral cortex, and cerebellum. SPECT by (123-I)2 β -carbomethoxy-3- β -(4-iodophenyl)-Tropan on two occasions revealed progressive loss of dopamine transporter suggestive of nigral neuronal apoptosis. Striatal and frontal hypometabolism and hypoperfusion were found by FDG-PET and HMPAO-SPECT. *Comment: Successful administration of DMAP and STS (doses were not reported) after ingestion of a lethal amount of KCN, but significant neurological sequelae.*

Zavotsky et al, 2004

A 52-years-old man came in contact with a liquid chemical believed to be 2,4,5 trichloro-6-carbopentoxyphenyl oxylate. Approximately two to four hours after exposure, he felt nauseous and began vomiting. Upon arriving at the emergency department, the staff decontaminated the patient. The vital signs were 109 heart beats/min, 20 breaths/min, blood pressure 110/38 mm Hg and SaO₂, 98%. He was still vomiting and complained of mild upper abdominal pain. Based on

laboratory findings and the patient's presentation, he was evaluated and treated for acute biliary disease and possible pancreatitis. After 18-20 hours after exposure the patient became acutely confused with BP of 80/30 mm Hg, and increasing heart and respiratory rates to 120 beats/min and 40 breaths/min. The patient's agitation became unmanageable, so he was sedated and intubated. BUN and creatinine were now 6 mg/dl, and 2.3 mg/dl, respectively, with associated oliguria, and a lactate level of 10.06 mmol/l indicating a profound metabolic acidosis. In a urine screen a thiocyanate level of >64 mg/l was found. After a further request to the employer a MSDS of acetonitrile was provided. 300 mg SN were administered intravenously over 10 minutes followed by 12.5 g STS intravenously over 10 minutes. Immediately the patient's condition began to improve. The patient was monitored for 72 hours. He had an acute episode of confusion shortly after beginning transferred to a medical-surgical floor and was treated again with SN and STS as a precaution. He was discharged home after six days.

Zilker, 2009

A female patient had ingested cyanide in a suicide attempt. She lost consciousness but was cardiorespiratory stable. 250 mg of DMAP were administered and she recovered completely. Cyanide was detected in the blood, quantity not given. *Comment: The case was reported on personally. DMAP showed good efficacy with no adverse effects. Although it was not explicitly cited it seems likely that the patient was given additional STS as this is recommended in cyanide poisoning.*

Zilker and Eyer (2002)

Seven intoxications with hydrogen cyanide through inhalation or inhalation together with skin contact are reported on. Five patients suffered mild intoxications (not even in coma). These persons were given 250 mg DMAP between 12 minutes and 5.5 hours after exposition and all recovered completely. The antidotal treatment did not lead to any adverse effects. The other two patients had been poisoned more seriously. They presented in deep coma. They also received 250 mg DMAP each and made an unremarkable recovery (one showed an elevated MetHb-level of 40% but without any symptoms of haemolysis). 5 mild intoxications, 2 moderate intoxications. All seven patients made a full recovery under DMAP treatment, but whether it was actually indicated in the mild poisoned patients could be discussed.

Zilker and Eyer, 2005

Zilker and Eyer reported 23 cases of intoxications with cyanide in a historical overview in 2005. Of the 23 cases 1 was an accidental administration of DMAP, 13 were also published by Werner (1979) in his thesis, 1 by Daunderer (1974) and a further 8 by Zilker and Stickel (2008).

Zilker, 2010

Personal communication. A 64-year female patient ingested an unknown amount of KCN showing signs and symptoms of moderate cyanide poisoning. After administration of 250 mg DMAP and 10 g STS she made a full recovery.

APPENDIX D: EFFECTS OF HOCO ON LABORATORY MEASUREMENTS AND PARAMETERS

Tissue / Parameter	Apparatus	HOCO (mg/l)	Interference (Change of concentration)	Remark	Reference
Plasma <i>in vitro</i>					
AST	Boehringer Mannheim/Hitachi 736 Analyzer	100, 500, 1,000	Decrease	At all three HOCO concentrations	Curry et al, 1994
GGT			Decrease		
Phosphorus			Increase		
Triglycerides *			Increase		
Human blood <i>in vitro</i>					
Total Hb	Co-oxymeter IL 482, CCD 270, AVL 912, OSM3, ABL 520	192.5 – 798.4 mg/l (143 - 593 μmol/l)	Increase	Overestimation of all three parameters, the magnitude of which depends on the concentration of HOCO	Gourlain et al, 1995
COHb			Increase		
MetHb			Increase		
LDH	Vitros® 750 XRC	31 - 1,125	Inaccuracy	At > 600 mg HOCO/l	Vest, 2002
Bilirubinaemia (total and conjugate), iron, phosphorus, creatinine, CPK, AST, protein, glucose, lactate			Inaccuracy	At all HOCO concentrations	
Bilirubinaemia (total and conjugate), creatinine, CPK	Cobas® Integra 700	31 - 1,125	Inaccuracy (≥ 10%)	At ≥ 124 mg HOCO/l	Vest, 2002
Total Hb	Co-oxymeter GEM Premier 4000s	0, 0.25, 0.5, 1, 2 g/l	Increase	Up to 2.6 g/dl	Pamidi et al, 2009
HbO ₂	IL 682		Decrease	2 - 20% bias	
COHb	Radiometer ABL 735		Increase	0 - 6% bias	
MetHb	Siemens RapidPoint 405		Increase	0 - 15% bias	

Tissue / Parameter	Apparatus	HOCO (mg/l)	Interference (Change of concentration)	Remark	Reference
Rabbit** blood <i>in vivo</i>					
Total Hb	Co-oxymeter	165 mg/kg i.v.	1.784 mM	Overestimation (2 - 3%)	Lee Ann, 2007
HbO ₂		1.784 mM	0.892 mM	Underestimation (62 - 8%)	
COHb		0.892 mM	0.892 mM	Overestimation (2 - 16%)	
MetHb		0.382 mM	0.892 mM	Overestimation (1 - 2%)	
Swine blood <i>in vitro</i> (haemolysed, oxygenated, de-oxygenated)					
MetHb	Spectrophotometer Varian Cary 100	X = 317.46		Overestimation from 0.1X to 10X	Denninghoff et al, 2008
		0.1X		The clinical relevance of this study is questionable	
		0.33X			
		2X			
		3.3X			
		6.66 X			
		10X			
		20X			

* HOCO had no effect on the concentration of Albumin, Alkaline Phosphatase, ALT, Amylase, BUN, Calcium, Cholesterol, Cholinesterase, CPK, Creatinine, Glucose, Iron, LDH, Magnesium, Total bilirubin, Total protein, Uric acid

** New Zealand strain

MEMBERS OF THE TASK FORCE

F. Baud (co-Chairman)	University Paris 7, Hôpital Lariboisière F - Paris
W. Steffens (co-Chairman)	Bayer D - Monnheim
S. Borron (Corresponding) Part-time	Texas Tech University USA - El Paso, TX
U. Küpper	Evonik Industries D - Wesseling
D. Pallapies ^a Part-time	Formerly BASF D - Hanau
R. Rödelsperger	Evonik Industries D - Darmstadt
H. Smet	Poisons Centre Brussels B - Brussels
T. Zilker	Technical University Munich D - Munich
W. Mayr (Technical Editor)	D - Wuppertal
H. Vrijhof (Scientific Secretary)	ECETOC B - Brussels

^a Resigned when he moved to Ruhr Bochum University.

MEMBERS OF THE SCIENTIFIC COMMITTEE*(Peer Review Committee at the time that the report was reviewed)*

F. Lewis (Chairman) Global Platform Lead	Syngenta UK - Bracknell, Berkshire
B. van Ravenzwaay (Vice Chairman) Senior Vice President - Experimental Toxicology	BASF D - Ludwigshafen
R. Bars Team Leader, Toxicology Research	Bayer CropScience F - Sophia Antipolis
D. Farrar Occupational Health Business Manager	Ineos Chlor UK - Runcorn
A. Flückiger ^a Chief Occupational Health Officer	F. Hoffmann - La Roche CH - Basel
H. Greim Institute of Toxicology and Environmental Hygiene	Technical University München D - München
G. Malinverno Global Governmental and Regulatory Affairs	Solvay B - Brussels / I - Milano
L. Maltby Professor of Environmental Biology	University of Sheffield UK - Sheffield
S. Marshall Environmental Science Leader	Unilever SEAC UK - Bedford
M. Meisters Manager Health and Environmental Sciences EMEA	DuPont de Nemours B - Mechelen
C. Money Industrial Hygiene Adviser, Europe	ExxonMobil B - Machelen

^a Steward responsible for primary peer review


MEMBERS OF THE SCIENTIFIC COMMITTEE (cont'd)

M. Pemberton ^a Director	Systox UK - Wilmslow
C. Rodriguez Principal Toxicologist, Corporate Central Product Safety	Procter and Gamble B - Strombeek-Bever
L. Rushton Principal Research Fellow	Imperial College London UK - London
D. Salvito Vice President, Environmental Sciences	RIFM USA - Woodcliff Lake
J. Snape Principal Scientist	AstraZeneca UK - Brixham
G. Swaen Senior Epidemiologist	Dow Chemical NL - Terneuzen
J. Tolls Director Environmental Safety Assessment	Henkel D - Düsseldorf
S. van der Vies Professor of Biochemistry	VU Medical Center NL - Amsterdam
C. van Leeuwen Principal Scientist	KWR Watercycle Research Institute NL - Nieuwegein
H.-J. Wiegand Product Stewardship, Corporate Environment, Safety, Health, Quality	Evonik D - Essen

^a Steward responsible for primary peer review

ECETOC PUBLISHED REPORTS

The full catalogue of ECETOC publications can be found on the ECETOC website:
<http://www.ecetoc.org/publications>



Responsible Editor:
Dr Alan Poole
ECETOC AISBL
Av. E. Van Nieuwenhuysse 2 (box. 8)
B-1160 Brussels, Belgium
VAT: BE 0418344469
www.ecetoc.org
D-2013-3001-231

Established in 1978, ECETOC (European Centre for Ecotoxicology and Toxicology of Chemicals) is Europe's leading industry association for developing and promoting top quality science in human and environmental risk assessment of chemicals. Members include the main companies with interests in the manufacture and use of chemicals, biomaterials and pharmaceuticals, and organisations active in these fields. ECETOC is the scientific forum where member company experts meet and co-operate with government and academic scientists, to evaluate and assess the available data, identify gaps in knowledge and recommend research, and publish critical reviews on the ecotoxicology and toxicology of chemicals, biomaterials and pharmaceuticals.