

JACC Report

No 25

**1-Chloro-1,2,2,2-tetrafluoroethane
(HCFC 124)
CAS No. 2837-89-0**

July 1994

ISSN-0773-6339-25

Joint Assessment of Commodity Chemicals No. 25

**1-Chloro-1,2,2,2-
tetrafluoroethane (HCFC 124)**

CAS No. 2837-89-0

July 1994

LIST OF ECETOC PUBLICATIONS (continued inside back cover)

MONOGRAPHS

No.	Title
No. 1	Good Laboratory Practice. Oct 79
No. 2	Contribution to Strategy for Identification and Control of Occupational Carcinogens. Sep 80
No. 2	Definition of a Mutagen, for 6th Amendment. Sep 80
No. 3	Risk Assessment of Occupational Chemical Carcinogens. Jan 82
No. 4	Hepatocarcinogenesis in Laboratory Rodents : Relevance for Man. Oct 82
No. 5	Identification and Assessment of the Effects of Chemicals on Reproduction and Development (Reproductive Toxicology). Dec 83
No. 6	Acute Toxicity Tests, LD ₅₀ (LC ₅₀) Determinations and Alternatives. May 85
No. 7	Recommendations for the Harmonisation of International Guidelines for Toxicity Studies. Dec 85
No. 8	Structure-Activity Relationships in Toxicology and Ecotoxicology: An Assessment. Feb 86
No. 9	Assessment of Mutagenicity of Industrial and Plant Protection Chemicals. Jun 87
No. 10	Identification of Immunotoxic Effects of Chemicals and Assessment of their Relevance to Man. Aug 87
No. 11	Eye Irritation Testing. Jun 88
No. 12	Alternative Approaches for the Assessment of Reproductive Toxicity (with emphasis on embryotoxicity/teratogenicity). Nov 89
No. 13	DNA and Protein Adducts: Evaluation of their Use in exposure Monitoring and Risk Assessment. Oct 89
No. 14	Skin Sensitisation Testing. Mar 90
No. 15	Skin Irritation. Jul 90
No. 16	Mutation Research, Special Issue: Early Indicators of Non-Genotoxic Carcinogenesis. Jun 91
No. 17	Hepatic Peroxisome Proliferation. May 92
No. 18	Evaluation of the Neurotoxic Potential of Chemicals. Sep 92
No. 19	Respiratory Allergy. Aug 93
No. 20	Percutaneous Absorption. Aug 93

JACC REPORTS

No.	Title
No. 1	Joint Assessment of Commodity Chemicals, Melamine. Feb 83
No. 2	Joint Assessment of Commodity Chemicals, 1,4-Dioxane. Feb 83
No. 3	Joint Assessment of Commodity Chemicals, Methyl Ethyl Ketone. Feb 83
No. 4	Joint Assessment of Commodity Chemicals, Methylene Chloride. Jan 84
No. 5	Joint Assessment of Commodity Chemicals, Vinylidene Chloride. Aug 85
No. 6	Joint Assessment of Commodity Chemicals, Xylenes. Jun 86
No. 7	Joint Assessment of Commodity Chemicals, Ethylbenzene. Aug 86
No. 8	Joint Assessment of Commodity Chemicals, Methyl Isobutyl Ketone. May 87
No. 9	Joint Assessment of Commodity Chemicals, Chlorodifluoromethane. Oct 89
No. 10	Joint Assessment of Commodity Chemicals, Isophorone. Sep 89
No. 11	Joint Assessment of Commodity Chemicals, (HFA-132b) 1,2-Dichloro-1,1-Difluoroethane. May 90
No. 12	Joint Assessment of Commodity Chemicals, (HFA-124) 1-Chloro-1,2,2,2-Tetrafluoroethane. May 90
No. 13	Joint Assessment of Commodity Chemicals, (HFA-123) 1,1-Dichloro-2,2,2-Trifluoroethane. May 90
No. 14	Joint Assessment of Commodity Chemicals, (HFA-133a) 1-Chloro-2,2,2-Trifluoromethane. Aug 90
No. 15	Joint Assessment of Commodity Chemicals, (HFA-141B) 1-Fluoro 1,1-Dichloroethane. Aug 90
No. 16	Joint Assessment of Commodity Chemicals, (HCFC-21) Dichlorofluoromethane. Aug 90
No. 17	Joint Assessment of Commodity Chemicals, (HFA-142b) 1-Chloro-1,1-Difluoroethane. Feb 91
No. 18	Joint Assessment of Commodity Chemicals, Vinylacetate. Feb 91
No. 19	Joint Assessment of Commodity Chemicals, Dicyclopentadiene. Jul 91
No. 20	Joint Assessment of Commodity Chemicals, Tris-/Bis-/Mono-(2-ethylhexyl)phosphate. May 92
No. 21	Joint Assessment of Commodity Chemicals, Tris-(2-butoxyethyl)-phosphate. Mar 92
No. 22	Joint Assessment of Commodity Chemicals, Hydrogen Peroxide. Jan 93
No. 23	Joint Assessment of Commodity Chemicals, Polycarboxylate Polymers as Used in Detergents. Nov 93
No. 24	Joint Assessment of Commodity Chemicals, (HFC-125) Pentafluoroethane. May 94
No. 25	Joint Assessment of Commodity Chemicals, (HCFC-124) 1-Chloro-1,2,2,2-Tetrafluoroethane. Jul 94

Joint Assessment of Commodity Chemicals No. 25

**1-Chloro-1,2,2,2-
tetrafluoroethane (HCFC 124)**

CAS No. 2837-89-0

July 1994

ISSN-0773-6339-25

ECETOC Jacc Report No. 25

© Copyright - ECETOC (European Centre for Ecotoxicology and Toxicology of Chemicals), 4 Avenue E. Van Nieuwenhuyse (Bte 6), 1160 - Brussels, Belgium.

All rights reserved. No part of this publication may be reproduced, copied, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise without the prior written permission of the copyright holder. Applications to reproduce, store, copy or translate should be made to the Director. ECETOC welcomes such applications. Reference to the document, its title and summary may be copied or abstracted in data retrieval systems without subsequent reference.

The content of this document has been prepared and reviewed by experts on behalf of ECETOC with all possible care and from the available scientific information. It is provided for information only. ECETOC cannot accept any responsibility or liability and does not provide a warranty for any Abuse or interpretation of the material contained in the publication.

THE ECETOC SCHEME FOR THE JOINT ASSESSMENT OF COMMODITY CHEMICALS

This report has been produced as part of the ECETOC programme for preparing critical reviews of the toxicology and ecotoxicology of selected existing industrial chemicals.

In the programme, commodity chemicals, that is those produced in large tonnage by several companies and having widespread and multiple uses, are jointly reviewed by experts from a number of companies with knowledge of the chemical. It should be noted that in a JACC review only the chemical itself is considered; products in which it appears as an impurity are not normally taken into account.

ECETOC is not alone in producing such reviews. There are a number of organisations that have produced and are continuing to write reviews with the aim of ensuring that toxicological knowledge and other information are evaluated. Thus a Producer, Government Official or Consumer can be informed on the up-to-date position with regard to safety, information and standards. Within ECETOC we do not aim to duplicate the activities of others. When it is considered that a review is needed every effort is made to discover whether an adequate review exists already; if this is the case the review is checked, its conclusions summarised and the literature published subsequent to the review assessed. To assist ourselves and others working in this field we publish annually a summary of international activities incorporating work planned, in hand, or completed on the review of safety data for commodity chemicals. Interested readers should refer to our Technical Report No. 30 entitled "Existing Chemicals: Literature Reviews and Evaluations".

This document is a revision of JACC report No. 12 originally published in May 1990 on 1-chloro-1,2,2,2-tetrafluoroethane (HCFC 124; CAS No. 2837-89-0). The text revisions introduce new environmental and toxicology data.

1-Chloro-1,2,2,2-tetrafluoroethane (HCFC-124)

CONTENTS

SECTION 1.	SUMMARY AND CONCLUSIONS	1
SECTION 2.	IDENTITY, PHYSICAL, AND CHEMICAL PROPERTIES, ANALYTICAL METHODS	3
2.1	IDENTITY	3
2.2	PHYSICAL AND CHEMICAL PROPERTIES	3
2.3	ANALYTICAL METHODS	4
SECTION 3.	PRODUCTION, STORAGE, TRANSPORT AND USE	5
SECTION 4.	ENVIRONMENTAL TRANSPORT, DISTRIBUTION, TRANSFORMATION, AND IMPACT	6
4.1	SOURCE	6
4.2	ENVIRONMENTAL DISTRIBUTION	6
4.3	ATMOSPHERIC LIFETIME	7
4.4	OZONE DEPLETING POTENTIAL	7
4.5	GLOBAL WARMING POTENTIAL	7
4.6	TROPOSPHERIC OZONE FORMATION	8
4.7	DEGRADATION MECHANISM AND PRODUCTS	8
4.8	CONTRIBUTION OF DEGRADATION PRODUCTS TO ENVIRONMENTAL CHLORIDE, FLUORIDE AND TRIFLUOROACETATE AND TO THE ACIDITY OF RAINWATER	9
4.9	BIODEGRADATION	10
SECTION 5.	ENVIRONMENTAL LEVELS AND HUMAN EXPOSURE	11
SECTION 6.	EFFECTS ON ORGANISMS IN THE ENVIRONMENT	12
SECTION 7.	KINETICS AND METABOLISM	13
7.1	ANIMAL STUDIES	13
7.2	HUMAN STUDIES	13

SECTION 8.	EFFECTS ON EXPERIMENTAL ANIMALS AND <i>IN VITRO</i> TEST	
	SYSTEMS	14
8.1	SINGLE EXPOSURE	14
8.2	REPEATED EXPOSURE	14
8.3	LONG-TERM EXPOSURE	16
8.4	SKIN AND EYE IRRITATION/ALLERGIC SENSITIZATION	16
8.5	SPECIAL STUDIES: CARDIOVASCULAR AND RESPIRATORY	
	EFFECTS	17
8.6	DEVELOPMENTAL TOXICITY/REPRODUCTIVE PERFORMANCE	17
8.7	MUTAGENICITY	18
8.8	CARCINOGENICITY	18
SECTION 9.	EFFECTS ON HUMANS	19
BIBLIOGRAPHY		20
MEMBERS OF THE TASK FORCE		22
MEMBERS OF THE SCIENTIFIC COMMITTEE		23

SECTION 1. SUMMARY AND CONCLUSIONS

1-Chloro-1,2,2,2-tetrafluoroethane (chlorotetrafluoroethane), a non-flammable colourless gas at room temperature, is currently under development as a chlorofluorocarbon alternative and is not yet available in commercial quantities.

The atmospheric degradation of chlorotetrafluoroethane will occur mainly in the troposphere by reaction with naturally occurring hydroxyl radicals. Its overall atmospheric lifetime is approximately 6.9 years and about 95% of the degradation will occur in the troposphere. Chlorotetrafluoroethane has a low calculated ozone depletion potential (ODP) of 0.022 and a global warming potential of 0.10 relative to a reference value for both indices of 1.0 for trichlorofluoromethane (CFC-11).

In limited studies in rats to evaluate metabolic fate and pharmacokinetics, chlorotetrafluoroethane appears to undergo oxidative metabolism resulting in the excretion of trifluoroacetic acid and fluoride ion in the urine. Studies to assess the metabolic fate of inhaled tetrafluoroethane in rats, mice, and hamsters are currently in progress.

Chlorotetrafluoroethane has a low order of acute inhalation toxicity with a 4-hour LC_{50} in rats between 1,283,400 and 1,674,000 mg/m^3 (230,000 and 300,000 ppm). The main toxicological action of this fluorocarbon during exposure is weak anaesthesia. As with many other halocarbons and hydrocarbons, inhalation of high concentrations of chlorotetrafluoroethane, followed by an intravenous epinephrine challenge to simulate stress, can induce a cardiac sensitization response in dogs. In these experimental screening studies, cardiac sensitisation (life-threatening arrhythmia) was seen at concentrations of 146,000 mg/m^3 (26,000 ppm) and above. The no-observable-effect-level was 55,800 mg/m^3 (10,000 ppm).

In a 2-week inhalation toxicity study, rats exposed to 558,000 mg/m^3 (100,000 ppm) chlorotetrafluoroethane showed no adverse effects. In 90-day studies at concentrations as high as 279,000 mg/m^3 (50,000 ppm), rats and mice showed minimal toxic effects such as slight central nervous system depression and minor blood chemistry changes. No-observable-effect level in these 90-day studies was 27,900 mg/m^3 (5,000 ppm) for male rats whilst for male mice a NOEL was not achievable. For female rats and mice a NOEL of 15,000 ppm was established. A lifetime inhalation toxicity/carcinogenicity study in rats is currently in progress.

No evidence of embryotoxicity or teratogenicity of chlorotetrafluoroethane was seen in developmental studies by inhalation in rats and rabbits at exposure levels as high as

279,000 mg/m³ (50,000 ppm). Minimal evidence of maternal toxicity was seen at concentrations of 83,700 mg/m³ (15,000 ppm) and above in each of these studies.


Chlorotetrafluoroethane was not mutagenic either in *in vitro* or in *in vivo* studies using bacteria or yeast or mammalian cell lines or the mouse micronucleus assay.

There are no reported effects of chlorotetrafluoroethane in man. An occupational exposure limit (8-hour average) of 1,000 ppm (5,580 mg/m³) is recommended by AIHA.

SECTION 2. IDENTITY, PHYSICAL, AND CHEMICAL PROPERTIES, ANALYTICAL METHODS

2.1 IDENTITY

Chemical Structure:



Chemical Formula:



Common Name:

1-Chloro-1,2,2,2-tetrafluoroethane
Chlorotetrafluoroethane

IUPAC Name:

Ethane, 2-chloro-1,1,1,2-tetrafluoro-

Common Synonyms:

1,1,1,2-Tetrafluoro-2-chloroethane
Fluorocarbon 124
HFA-124; HCFC-124¹

CAS Registry Number:

2837-89-0

EINECS Number:

220-629-6

Conversion Factors:

1 ppm = 5.58 mg/m³ at 25°C
1 mg/m³ = 0.179 ppm at 25°C

2.2 PHYSICAL AND CHEMICAL PROPERTIES

Chlorotetrafluoroethane is a nonflammable colourless gas. Some physical and chemical data are given in Table 1.

¹ Abbreviations mean: Hydro-Chloro-Fluoro-Carbon C₂HClF₄

First figure = number of C-Atoms minus 1	1
Second figure = number of H-Atoms minus 2	2
Third figure = number of F-Atoms	4
= HCFC 124	

2.3 ANALYTICAL METHODS

A method of analysis for chlorotetrafluoroethane has been described using gas chromatography with dual flame ionization detection (Malley, 1991a).

Table I Physical and chemical properties of chlorotetrafluoroethane*

Molecular Weight	136.5
Physical form	Gas
Colour	Colourless
Boiling point, °C at 1013 hPa	-12
Liquid density at 25°C, g/ml	1.36
Vapour density (air = 1)	4.7
Vapour pressure, hPa at 25°C	3,850
% Volatiles by volume at 20°C	100
Solubility in water at 25°C, g/l at 1013 hPa	1.45
Solubility in organic solvents	Miscible with acetone, ethanol, petroleum solvents
Log P _{ow}	1.9-2.0
Log K _{oc}	1.7-1.9 (calculated)
Flammability	Nonflammable

* From E. I. duPont de Nemours & Company, Inc.; Material Safety Data Sheet - HCFC124; January 1993.

SECTION 3. PRODUCTION, STORAGE, TRANSPORT AND USE

Processes for the production of chlorotetrafluoroethane are still under development. Possible routes are hydrofluorination of tetrachloroethylene or hydrodechlorination of 1,1-dichloro-1,2,2,2-tetrafluoroethane (CFC-114a). The grade of purity obtained is greater than 99%.

So far, chlorotetrafluoroethane is produced only on a small scale. Like other hydrochloro-fluorocarbons, chlorotetrafluoroethane is regulated under the Montreal Protocol and is thus scheduled for virtual phase-out by 2020. Chlorotetrafluoroethane is being developed as a substitute for fully halogenated chlorofluorocarbons. Its main potential applications are believed to be as a component for refrigerant blends and as a blowing agent for polystyrene and polyolefin foams.

SECTION 4. ENVIRONMENTAL TRANSPORT, DISTRIBUTION, TRANSFORMATION, AND IMPACT

4.1 SOURCE

There is no known natural source of chlorotetrafluoroethane.

The figure of 25 kt/y is adopted here to represent a conservative upper limit to possible future annual emissions, in order to assess certain aspects of the potential environmental impact of chlorotetrafluoroethane.

4.2 ENVIRONMENTAL DISTRIBUTION

On the basis of the following physical properties, chlorotetrafluoroethane may be expected, when released to the environment, to partition almost exclusively into the atmosphere:

- it is a gas at room temperature and atmospheric pressure, boiling at -12°C ;
- its Henry's Law constant for dissolution in water is 1.45 g/l.bar at 25°C ; for an atmospheric concentration of 7 ppt (v/v) (i.e., the calculated steady-state concentration resulting from emissions of 25 kt/y), the equilibrium concentration in cloud and surface waters would thus be 0.01 ppt (w/w);
- any chlorotetrafluoroethane which might be present in aqueous waste streams discharged directly into rivers or lakes would be expected, by analogy with similar compounds, to have a half-life with respect to volatilization of days or a few weeks at the very most.

Moreover, any chlorotetrafluoroethane present in surface or ground waters would have little tendency to partition onto biota or soil. $\log P_{ow}$ is 1.9-2.0, indicating a low potential for passive bioaccumulation. From various correlations, $\log K_{oc}$ is estimated to lie in the range 1.7-1.9, which means that chlorotetrafluoroethane would be moderately mobile in soils. The atmospheric lifetime of chlorotetrafluoroethane (about 7 y, see below) is much longer than either the intrahemispheric or interhemispheric mixing times. Therefore this compound will become more or less uniformly distributed in the atmosphere on a global scale (at a concentration <7 ppt (v/v), see above).

4.3 ATMOSPHERIC LIFETIME¹

The atmospheric degradation of chlorotetrafluoroethane will occur mainly (about 95%) in the troposphere, being initiated by attack of naturally occurring hydroxyl radicals. The lifetime with respect to this process is estimated to be 7.3 y. Additionally some chlorotetrafluoroethane will be transported to the stratosphere and be degraded there by reaction with hydroxyl radicals, oxygen atoms and by photolysis. The overall atmospheric lifetime is estimated to be 6.9 years (WMO, 1991).

4.4 OZONE DEPLETING POTENTIAL

Ozone Depleting Potentials (ODPs) express the stratospheric ozone loss due to emission of a unit mass of a given compound, divided by the ozone loss due to emission of the same mass of a reference compound.

The latest estimates of the ODP of chlorotetrafluoroethane, carried out using 3 different atmospheric models, give results in the range 0.016-0.034 relative to a reference value of 1.0 for CFC-11. The "semi-empirical" ODP was found to be 0.022 (WMO, 1991). The latter value was adopted as the best estimate of the ODP for regulatory purposes in the 1992 Montreal Protocol revision.

The ODP value may vary upwards or downwards in the future, as the models are refined and new kinetic data become available. For instance, use of a recently revised value of the rate constant for the reaction of OH-radicals with CCl_3CH_3 (the reference reaction for atmospheric lifetime calculations), should lead to a downward revision of the ODP of chlorotetrafluoroethane by about 15%.

4.5 GLOBAL WARMING POTENTIAL

Global Warming Potentials (GWPs) express the radiative forcing (increase in earthward infra-red radiation flux) due to emission of a unit mass of a given compound, divided by the radiative forcing due to emission of the same mass of a reference compound.

Based on the lifetimes quoted above, the halocarbon Global Warming Potential (HGWP) of chlorotetrafluoroethane is 0.10 (AER, 1992) relative to a reference value of 1.0 for

¹ Lifetime (LT) is the time necessary for 63% degradation; it is equal to the "half-life" divided by $\ln 2$ (≈ 0.69)

trichlorofluoromethane (CFC-11). This assessment assumes a pulse emission and an infinite integration time horizon (ITH), which is mathematically equivalent to a steady-state calculation.

GWPs may also be expressed relative to CO₂ as the reference substance, and assessed over a finite ITH. For chlorotetrafluoroethane, the corresponding values are 1,500; 440 and 150 (relative to a reference value of 1.0 for CO₂ at each ITH), for ITHs of 20, 100, and 500 y respectively (WMO, 1991; IPCC, 1992).

4.6 TROPOSPHERIC OZONE FORMATION

Chlorotetrafluoroethane is too unreactive in the atmosphere to make any significant contribution to local urban tropospheric ozone formation, and the related "photochemical smog", near the emission sources (WMO, 1989).


4.7 DEGRADATION MECHANISM AND PRODUCTS

(see Figure 1 for reaction scheme)

Support for the basic tropospheric degradation mechanism for chlorotetrafluoroethane proposed in WMO (1989) has been provided by recent laboratory studies (see, for example: WMO, 1991; Edney and Driscoll, 1992; Tuazon and Atkinson, 1993).

Breakdown of chlorotetrafluoroethane in the troposphere will be initiated by OH-radicals and will proceed via various free-radical intermediates to give HCl and CF₃COF. The latter are expected to be removed from the atmosphere within a few days to a few months, by uptake into clouds, rain and the oceans, CF₃COF then being rapidly hydrolysed to trifluoroacetic acid (TFA) and hydrofluoric acid (HF) (AFEAS, 1992; STEP/AFEAS, 1993; Franklin, 1993).

Although a peroxyxynitrate (CF₃CClFO₂NO₂) and a hydroperoxide (CF₃CClFO₂H) may be formed during the degradation of chlorotetrafluoroethane, they are not thought to play a significant role in the atmospheric chemistry of chlorotetrafluoroethane and are probably rather short-lived intermediates.

Figure 1 Tropospheric Degradation Mechanism for HCFC-124

4.8 CONTRIBUTION OF DEGRADATION PRODUCTS TO ENVIRONMENTAL CHLORIDE, FLUORIDE AND TRIFLUOROACETATE AND TO THE ACIDITY OF RAINWATER

Assuming an atmospheric release rate of 25 kt chlorotetrafluoroethane/y (likely upper limit), complete conversion of the latter into HCl, HF, and CF_3COOH (1 mol of each per mol chlorotetrafluoroethane) and uniform scavenging of the acids produced into the global average rainfall of 5×10^{11} kt/y, it follows that the levels of chloride, fluoride, and acidity produced are low compared with those arising from existing sources for the following reasons:

- Cl^- production from chlorotetrafluoroethane would be 7 kt/y, i.e. insignificant compared with the natural atmospheric chloride flux of roughly 1×10^7 kt/y, mainly arising from sea-salt aerosols (WMO, 1989);

- F⁻ production would be 4 kt/y, i.e. very small compared with the estimated atmospheric fluoride flux of 1,000-8,000 kt/y (WMO, 1989);
- the contribution of chlorotetrafluoroethane to the fluoride concentration in rainwater would be 7 ppt (w/w); this should be compared with typical fluoride concentrations in "background" rainwater of around 10 ppb (w/w), i.e. 1,500 times greater, and with levels of about 1 ppm (w/w), used for the fluoridation of drinking water, i.e. 150,000 times greater (WMO, 1989);
- the contribution of chlorotetrafluoroethane to trifluoroacetate in rainwater would be about 40 ppt (w/w);
- the hydrochloric, hydrofluoric, and trifluoroacetic acids formed from chlorotetrafluoroethane and scavenged in rainwater would represent an acidity of a little over 5×10^8 mol H⁺/y, i.e. 20,000 times less than the acidity arising from natural and anthropogenic emissions of SO₂ and NO_x (UKRGAR, 1990). Thus the contribution of chlorotetrafluoroethane to acid rain would be negligible.

4.9 BIODEGRADATION

In a closed-bottle assay for ready biodegradability, chlorotetrafluoroethane in aqueous solution appeared to be stable (1-2% loss) over a 4 wk period (Tobeta, 1992).

SECTION 5. ENVIRONMENTAL LEVELS AND HUMAN EXPOSURE

No observations of chlorotetrafluoroethane in the atmosphere or other environmental compartments have yet been reported.

SECTION 6. EFFECTS ON ORGANISMS IN THE ENVIRONMENT

No data are available on the effects of chlorotetrafluoroethane on environmental organisms.

SECTION 7. KINETICS AND METABOLISM

7.1 ANIMAL STUDIES

Limited metabolic data have been published on chlorotetrafluoroethane. Olson *et al* (1991) exposed rats to 558,000 mg/m³ (10,000 ppm) chlorotetrafluoroethane for two hours and found both inorganic fluoride and trifluoroacetic acid in their urine, both being products of an oxidative metabolism. Increases in urinary fluoride levels in rats exposed by subchronic inhalation to chlorotetrafluoroethane had also been reported earlier by Brewer (1977a) and Malley (1991a). Additional evidence for an oxidative metabolism was provided by Harris *et al* (1992) who exposed rats for 6 hours to 558,000 mg/m³ (10,000 ppm) chlorotetrafluoroethane and found trifluoroacetylated proteins in liver and trifluoroacetic acid in urine, similar to earlier results reported by Harris *et al* (1991). The oxidative metabolism was approximately 5 fold less for chlorotetrafluoroethane compared to halothane (1-chloro-1-bromo-2,2,2-trifluoroethane) and dichlorofluoroethane. Work is currently underway (Anders, 1993a,b) to investigate the metabolism of inhaled chlorotetrafluoroethane in rats, mice, and hamsters.

7.2 HUMAN STUDIES

No data were found for humans relative to the absorption, distribution, metabolic transformation or elimination of chlorotetrafluoroethane.

SECTION 8. EFFECTS ON EXPERIMENTAL ANIMALS AND *IN VITRO* TEST SYSTEMS

8.1 SINGLE EXPOSURE

In recent studies by Kelly (1990) groups of 6 male rats were exposed, nose-only, for single 4-h periods to atmospheres containing chlorotetrafluoroethane vapour at concentrations of 267,840, 902,800, 1,283,400, or 1,674,000 mg/m³ (48,000, 162,000, 230,000 or 300,000 ppm). Aside from a slight weight loss one day post-exposure, rats exposed at the lowest exposure level showed no clinical signs of toxicity during or after exposure. At concentrations of 902,800 and 1,283,400 mg/m³ (162,000 and 230,000 ppm), no mortality occurred but rats did show a temporary weight loss, decreased acoustic-startle responses, prostration, lethargy and incoordination; these anaesthetic effects disappeared shortly after exposure. At 1,674,000 mg/m³ (300,000 ppm), the same clinical signs occurred, but 6 of 6 rats died during the 4-h exposure. Under the conditions of this study, the 4-h LC₅₀ of chlorotetrafluoroethane in rats was estimated to be between 1,283,000 and 1,674,000 mg/m³ (230,000 and 300,000 ppm). Wada (1977) had previously reported a 10-min approximate lethal concentration of 2,460,000 mg/m³ (440,000 ppm) for mice; exposure at 2,230,000 mg/m³ (400,000 ppm) for 10 min produced no mortality. Anaesthesia was observed at 837,000 mg/m³ (150,000 ppm) and no effects were seen at 558,000 mg/m³ (100,000 ppm).

In an earlier study by Coate (1976), a single 6-h exposure to chlorotetrafluoroethane at a concentration of 2,010,000 mg/m³ (360,000 ppm) was not lethal to rats but rapidly depressed motor activity; this was followed by anaesthesia after one hour. Rats exposed to 558,000 mg/m³ (100,000 ppm) were active for 30 min but then inactive until termination of exposure. Rats exposed to 2,010,000 mg/m³ (360,000 ppm) gained less weight than either the controls or the lower exposure group between days 1 and 8.

When dogs were exposed to chlorotetrafluoroethane for 10 min at exposure concentrations ranging from 2,232,000 to 3,906,000 mg/m³ (400,000 to 700,000 ppm), light surgical anaesthesia was rapidly induced. Complete recovery occurred within 7-10 minutes post-exposure (Van Poznak and Artusio, 1960).

8.2 REPEATED EXPOSURE

Trochimowicz *et al* (1977) had reported no adverse effects in 10 rats exposed by inhalation to chlorotetrafluoroethane for 6 h/d, 5 d/wk for 2 wk at a concentration of 558,000 mg/m³

(100,000 ppm). A four-week inhalation study was conducted by Malley (1990) with five groups of 10 male and 10 CD®BR female rats. The exposure regimen was 6 h/d, 5 d/wk for 4 wk at concentrations of 0, 2,790, 11,160, 55,800 or 279,000 mg/m³ (0, 500, 2,000, 10,000 or 50,000 ppm) chlorotetrafluoroethane. Compound related effects on body weight, food consumption, mortality, clinical laboratory parameters, organ weights, and tissue morphology did not occur at any exposure concentration. During the exposure period, rats exposed to 279,000 mg/m³ (50,000 ppm) appeared to be lethargic and uncoordinated. However, at the end of the daily exposure period, when the animals were removed from the chambers, no evidence of lethargy or incoordination was observed. The no-observable-effect level (NOEL) was considered to be 55,800 mg/m³ (10,000 ppm).

In a 90-day inhalation toxicity study by Malley (1991a), 20 CD®BR rats/sex/level were exposed to chlorotetrafluoroethane for 6 h/d, 5 d/wk at concentrations of 0, 27,900, 83,700, or 279,000 mg/m³ (0, 500, 15,000 or 50,000 ppm). Ten rats/sex/level were designated for a neurotoxicity evaluation using a Functional Observational Battery (FOB) and for an additional one-month recovery period. There were no compound-related effects on body weight, food consumption, mortality, clinical signs, haematology, organ weights, and tissue morphology at any exposure concentration. During exposure, rats exposed to 279,000 mg/m³ (50,000 ppm) were less responsive to stimuli compared to control rats as demonstrated by a decreased reaction to a knock on the exposure chamber wall. In addition, on the day following the last exposure, male rats exposed to 83,700 or 279,000 mg/m³ (15,000 or 50,000 ppm) chlorotetrafluoroethane had decreased arousal times (in 4 of 10 and 6 of 10 rats, respectively). Relative to blood chemistry measurements, at the 45-day clinical evaluation interval, male rats exposed to 83,700 and 279,000 mg/m³ (15,000 or 50,000 ppm) had lower serum triglyceride concentrations than controls and female rats at 279,000 mg/m³ (50,000 ppm) showed increased alkaline phosphatase activity. At all exposure levels, fluoride concentration in blood and urine was elevated at several sampling intervals during exposure and also at one month post-exposure. Male rats exposed at the two highest exposure levels also showed a mild diuresis, probably a result of increased osmotic activity from the excreted fluoride ion rather than a direct effect of the compound. The no-observable-effect level (NOEL) for this 90-day study was considered to be 27,900 mg/m³ (5,000 ppm) for male rats and 83,700 mg/m³ (15,000 ppm) for female rats.

Malley (1991b) also conducted a 90-day inhalation toxicity study in 20 CD®-1(ICR)BR mice/sex/group exposed for 6 h/d, 5 d/wk at the same concentration levels of chlorotetrafluoroethane as used in the rat study (see above). Responses in mice were similar to the responses described for rats. There were no compound-related effects on mortality, food consumption, clinical signs, haematology, organ weights and tissue morphology at any exposure concentration for either male or female mice. In male mice killed at 90-days, body weights were 6

to 7% lower than controls (not dose related) and hepatic β -oxidation activity was approximately two-fold higher at all exposure levels, while serum triglyceride concentrations were decreased at the two highest exposure levels. During exposure, both male and female mice exposed at 279,000 mg/m³ (50,000 ppm) chlorotetrafluoroethane were less responsive to stimuli (knocking on chamber window) compared to controls. All the effects seen in female or in male mice during exposure disappeared after 30 days post-exposure. Although all these effects in male mice were slight and of negligible toxicological significance, a NOEL was not achieved for male mice. For female mice, the NOEL was 83,700 mg/m³ (15,000 ppm).

One 90-day inhalation study in rats using a sample of unknown purity had been conducted by Brewer (1977a) at exposure levels up to 27,900 mg/m³ (5,000 ppm) chlorotetrafluoroethane and was reported earlier by ECETOC (1990). However, it will not be discussed here since the recent studies in rodents (Malley, 1991a,b) at similar and higher exposure levels, and conducted for even longer durations, have not confirmed the findings reported by Brewer (1977a).

The minimal, reversible effects seen in the preceding studies are not considered to be adverse. Therefore the subchronic no-observable-adverse-effect level (NOAEL) of chlorotetrafluoroethane for rats or mice is considered to be 279,000 mg/m³ (50,000 ppm).

8.3 LONG-TERM EXPOSURE

No data available.

A lifetime inhalation toxicity/carcinogenicity bioassay in rats under the sponsorship of the Program for Alternative Fluorocarbon Toxicity Testing (PAFT) is currently in progress. Approximately 85 rats/sex/group are being exposed to chlorotetrafluoroethane for 6 h/d, 5 d/wk for 104 wk at concentrations of 0, 27,900, 83,700 or 279,000 mg/m³ (0, 5,000, 15,000 or 50,000 ppm). The exposure phase of this study began in October 1992 and no results are available at the present time.

8.4 SKIN AND EYE IRRITATION/ALLERGIC SENSITIZATION

As chlorotetrafluoroethane is a gas at ambient temperature, studies to assess skin and eye irritation as well as skin sensitization have not been carried out. No evidence of skin or mucosal irritation was seen in rats exposed by inhalation on an acute or a repeated basis (see section 8.2).

8.5 SPECIAL STUDIES: CARDIOVASCULAR AND RESPIRATORY EFFECTS

In an experimental screening study (Mullin, 1976) in dogs, inhalation exposure to high concentrations of chlorotetrafluoroethane for 5 minutes, followed by an intravenous epinephrine challenge ($\sim 8 \mu\text{g/kg}$) induced cardiac sensitization at concentrations of $146,000 \text{ mg/m}^3$ ($26,000 \text{ ppm}$) and above. The NOEL cardiac sensitization in this study was $55,800 \text{ mg/m}^3$ ($10,000 \text{ ppm}$).

In repeated exposure studies (see Section 8.2) by the inhalation route, chlorotetrafluoroethane concentrations as high as $279,000$ to $558,000 \text{ mg/m}^3$ ($50,000$ to $100,000 \text{ ppm}$) produced no clinical or histopathological evidence of respiratory irritation in mice or rats respectively.

Older studies to evaluate anaesthetic potency of chlorotetrafluoroethane have been reported earlier (ECETOC, 1990).

8.6 DEVELOPMENTAL TOXICITY/REPRODUCTIVE PERFORMANCE

In a developmental toxicity study by Alvarez (1990), 24 pregnant CD[®]BR rats/group were exposed by whole-body inhalation to chlorotetrafluoroethane for 6 h/d on days 6 through 15 of gestation at concentrations of 0, 27,900, 83,700 or $279,000 \text{ mg/m}^3$ (0, 5,000, 15,000 or $50,000 \text{ ppm}$). No evidence of embryotoxicity, foetotoxicity, or teratogenicity was seen at any exposure level. Maternal toxicity was demonstrated at $279,000 \text{ mg/m}^3$ ($50,000 \text{ ppm}$) by a decreased rate of weight gain and food consumption during the first 4 days of exposure and by a reduced response to auditory stimuli during exposure but not shortly thereafter. Under these study conditions, the NOAEL for chlorotetrafluoroethane was $83,700 \text{ mg/m}^3$ ($15,000 \text{ ppm}$) for the dam and $279,000 \text{ mg/m}^3$ ($50,000 \text{ ppm}$) for the conceptus.

Using the same target exposure levels as the preceding study, Schroeder (1991) exposed 20 pregnant New-Zealand white rabbits/group by whole-body inhalation to chlorotetrafluoroethane vapour for 6 h/d on days 6 through 18 of gestation. There was no evidence of embryotoxicity, foetotoxicity, or teratogenicity at any exposure level. Similar to the preceding results in rats, slight maternal effects were seen at $279,000 \text{ mg/m}^3$ ($50,000 \text{ ppm}$) as evidenced by decreased food consumption and decreased in-chamber activity. The no-observable-effect level (NOEL) for maternal effects was reported as $83,700 \text{ mg/m}^3$ ($15,000 \text{ ppm}$) while the NOEL for developmental toxicity was considered to be $279,000 \text{ mg/m}^3$ ($50,000 \text{ ppm}$).

In a limited developmental toxicity study using a sample of unknown purity, Brewer (1977b) exposed 20 pregnant Albino rats to $27,900 \text{ mg/m}^3$ ($50,000 \text{ ppm}$) chlorotetrafluoroethane for 6 h/d on days 6

through 15 of gestation. No evidence of maternal toxicity, foetotoxicity, embryotoxicity, or teratogenicity was seen at this single exposure level. This study was reported in more detail earlier (ECETOC, 1990).

No data were available to evaluate the reproductive performance of male and female animals exposed to chlorotetrafluoroethane. However, in rats and mice exposed for 90 days by inhalation to chlorotetrafluoroethane concentrations as high as 279,000 mg/m³ (50,000 ppm) (Malley, 1991a,b), no histopathological effects were seen in the reproductive organs/tract of male or female animals.

8.7 MUTAGENICITY

The *in vitro* mutagenicity potential of chlorotetrafluoroethane has been evaluated in several bacteria and cells. In a series of plate and suspension assays using *Salmonella typhimurium* (Brusick, 1976; Barsky, 1976; Longstaff *et al*, 1984; Reynolds, 1990; May, 1991), strains TA1535, TA1537, TA1538, TA98 and TA100 were exposed to various concentrations of chlorotetrafluoroethane in the presence and absence of an S-9 metabolic activation system. None of these studies showed any evidence of mutagenicity potential. Chlorotetrafluoroethane also showed no mutagenicity potential in other *in vitro* assay with *Escherichia coli* (May, 1991) and *Saccharomyces cerevisiae* (Brusick, 1976). Furthermore chlorotetrafluoroethane showed no evidence of clastogenic activity in cultured Chinese Hamster Ovary (CHO-K1) cells (Edwards, 1991) or in human lymphocytes with or without S-9 metabolic activation (Bentley, 1990).

In one *in vivo* mutagenicity study (Rickard, 1990), chlorotetrafluoroethane was tested for its ability to induce micronuclei in bone marrow polychromatic erythrocytes of male and female mice. The animals were exposed, head-only, to either 0 or 552,400 mg/m³ (0 or 99,000 ppm) vapour in air for 6 h/day on two consecutive days. Bone marrow smears were prepared 24 and 48 h after exposure. Chlorotetrafluoroethane did not induce micronuclei and was considered negative for *in vivo* mutagenicity potential.

In conclusion, chlorotetrafluoroethane is not mutagenic *in vitro* and *in vivo*.

8.8 CARCINOGENICITY

No data to evaluate the carcinogenicity potential of chlorotetrafluoroethane were available. However, an inhalation carcinogenicity bioassay in rats by the inhalation route is currently underway (Section 8.3).

SECTION 9. EFFECTS ON HUMANS

There are no reported adverse health effects which can be ascribed to chlorotetrafluoroethane.

The American Industrial Hygiene Association's Workplace Environmental Exposure Level (WEEL) Committee assigned chlorotetrafluoroethane an occupational exposure limit (8-hour time-weighted average) of 1,000 ppm (5,580 mg/m³) (AIHA, 1992).

BIBLIOGRAPHY

- AER, 1992. Atmospheric and Environmental Research, Cambridge, Massachusetts, USA. Private communication to AFEAS, 23 January 1992.
- AFEAS, 1992. Proceedings of the AFEAS Workshop: Atmospheric Wet and Dry Deposition of Carbonyl and Haloacetyl Halides. Brussels, 22 September 1992.
- Alvarez L, 1990. Teratogenicity Study of HCFC-124 in Rats (HLR 482-90). Conducted by Haskell Laboratory for Toxicology and Industrial Medicine, E.I. duPont de Nemours & Company, Newark, DE 19711, USA.
- AIHA, 1992. Workplace Environmental Exposure Level (WEEL), Guide for 2-Chloro-1,1,1,2-Tetrafluoroethane. American Industrial Hygiene Association, 345 White Pond Drive, P.O. Box 8390, Akron, Ohio 44320, USA.
- Anders MW, 1993a. Metabolism and Toxicity of Hydrochlorofluoro-carbons: Current Knowledge and Needs for the Future. *Environ. Hlth. Perspec.* 96:185-191.
- Anders MW, 1993b. Metabolism Studies in Rats, Mice, and Hamsters. Work Underway at the University of Rochester, NY, USA, under Sponsorship of the Program for Alternative Fluorocarbon Toxicity Testing (PAFT).
- Barsky FC, 1976. *In Vitro* Microbial Mutagenicity Studies of Ethane, 1-Chloro-1,1,1,2-Tetrafluoro (HLR 349-76). Conducted by Haskell Laboratory for Toxicology and Industrial Medicine, E.I. duPont de Nemours & Company, Newark, DE 19711, USA.
- Bentley KS, 1990. *In Vitro* Evaluation of HCFC-124 for Chromosome Aberrations in Human Lymphocytes (HLR 367-90). Conducted by Haskell Laboratory for Toxicology and Industrial Medicine, E.I. duPont de Nemours & Company, Newark, DE 19711, USA.
- Brewer WE, 1977a. Ninety-Day Subacute Inhalation Toxicity Study with Genetron 124 in Albino Rats (IBT No. 8562-09345). Conducted by Industrial Bio-Test Laboratories, Inc for Allied Chemical Corporation, Morristown, NJ, USA.
- Brewer WE, 1977b. Teratogenic Study via Inhalation with Genetron 124 in Albino Rats (IBT No. 8562-09345). Conducted by Industrial Bio-Test Laboratories, Inc. for Allied Chemical Corporation, Morristown, NJ, USA.
- Brusick DJ, 1976. Mutagenicity Evaluation of Genetron 124 (LBI No. 2547). Conducted by Litton Bionetics, Inc., for Allied Chemical Corporation, Morristown, NJ, USA.
- Coate WB, 1976. LC50 of G124 in Rats (HLA No. 165-163). Conducted by Hazleton Laboratories American, Inc. for Allied Chemical Corporation, Morristown, NJ, USA.
- ECETOC, 1990. 1-chloro-1,2,2,2-tetrafluoroethane Joint Assessment of Commodity Chemicals No 12. European Center for Ecotoxicology and Toxicology of Chemicals, Brussels.
- Edney EO and Driscoll DJ, 1992. Chlorine Initiated Photooxidation Studies of Hydrochlorofluorocarbons (HCFCs) and Hydrofluorocarbons (HFCs): Results for HCFC-22 (CHClF_2); HFC-41 (CH_3F); HCFC-124 ($\text{CClF}_2\text{CHF}_3$); HFC-125 (CF_3CHF_2); HFC-134a ($\text{CF}_3\text{CH}_2\text{F}$); HCFC-142b (CClF_2CH_3); and HFC-152a (CHF_2CH_3). *Int. J. Chem. Kinet.* 24, 1067-1081.
- Edwards CN, 1991. *In Vitro* Assessment of the Clastogenic Activity of HCFC-124 in Cultured Chinese Hamster Ovary CHO-K1 Cells. (LSR No. 91/PAR002/0885a), conducted by Life Science Research (LSR) Limited for the Program for Alternative Fluorocarbon Toxicity Testing (PAFT).
- Franklin J, 1993. The Atmospheric Degradation and Impact of 1,1,1,2-Tetrafluoroethane (Hydrofluorocarbon 134a). *Chemosphere*, 27(8), 1565-1601.
- Harris JW, Pohl LR, Martin JL, and Anders MW, 1991. Tissue Acylation by the Chlorofluorocarbon Substitute 2,2-Dichloro-1,1,1-Trifluoroethane. *Proc. Natl. Acad. Sci. USA* 88: 1407-1410.
- Harris JW, Jones JP, Martin JL, LaRosa AC, Olson MJ, Pohl LR, and Anders MW, 1992. Pentahaloethane-Based Chlorofluorocarbon Substitutes and Halothane: Correlation of *In Vivo* Hepatic Protein Trifluoroacetylation and Urinary Trifluoroacetic Acid Excretion with Calculated Enthalpies of Activation. *Chem. Res. Toxicol.* 5(5):720-725.
- IPCC, 1992. Intergovernmental Panel on Climate Change. Climate Change 1992. The Supplementary Report to the IPCC Scientific Assessment. Houghton, J.T., Callander, B.A. and Varney, S.K., Eds. Cambridge University Press.
- Kelly DP, 1990. Four-Hour Inhalation Approximate Lethal Concentration (ALC) of HCFC-124 in Rats (HLR 71-90). Conducted by Haskell Laboratory for Toxicology and Industrial Medicine, E.I. duPont de Nemours & Company, Newark, DE 19711, USA.
- Longstaff E, Robinson M, Bradbrook C, Styles JA and Purchase IFH, 1984. Genotoxicity and Carcinogenicity of Fluorocarbons: Assessment by Short-Term *In Vitro* Tests and Chronic Exposure in Rats. *Toxicol. Appl. Pharmacol.* 72:15-31.
- Malley LA, 1990. Subchronic Inhalation Toxicity: Four-Week Study with HCFC-124 in Rats (HLR 257-90). Conducted by Haskell Laboratory for Toxicology and Industrial Medicine, E.I. duPont de Nemours & Company, Newark, DE 19711, USA.
- Malley LA, 1991a. Subchronic Inhalation Toxicity: 90-Day Study with HCFC-124 in Rats (HLR 79-91). Conducted by Haskell Laboratory for Toxicology and Industrial Medicine, E.I. duPont de Nemours & Company, Newark, DE 19711, USA.

- Malley LA, 1991b. Subchronic Inhalation Toxicity: 90-Day Study with HCFC-124 in Mice (HLR 695-91). Conducted by Haskell Laboratory for Toxicology and Industrial Medicine, E.I. duPont de Nemours & Company, Newark, DE 19711, USA.
- May K, 1991. HCFC-124 in Gaseous Phase: Assessment of Mutagenic Potential in Amino-Acid Auxotrophs of *Salmonella typhimurium* and *Escherichia Coli* (the Ames Test) (LSR No. 91/PAR001/0842). Conducted by Life Science Research Limited (UK) for the Program for Alternative Fluorocarbon Toxicity Testing (PAFT).
- Mullin LS, 1976. Cardiac Sensitization Study in Beagle Dogs on 1-Chloro-1,2,2,2-Tetrafluoroethane (HLR220-76). Conducted by Haskell Laboratory for Toxicology and Industrial Medicine, E.I. duPont de Nemours & Company, Newark, DE 19711, USA.
- Olson MJ, Johnson JT, O'Gara JF, and Surbrook SE Jr, 1991. Metabolism *In Vivo* and *In Vitro* of the Refrigerant Substitute 1,1,1,2-Tetrafluoro-2-Chloroethane. Drug Metab. Dispos. 19:1004-1011.
- Reynolds VL, 1990. Mutagenicity Testing of HCFC-124 in the *Salmonella Typhimurium* Plate Incorporation Assay (HLR 191-90). Conducted by Haskell Laboratory for Toxicology and Industrial Medicine, E.I. duPont de Nemours & Company, Newark, DE 19711, USA.
- Rickard LB, 1990. Mouse Bone Marrow Micronucleus Assay of HCFC-124 (HLR 52-90). Conducted by Haskell Laboratory for Toxicology and Industrial Medicine, E.I. duPont de Nemours & Company, Newark, DE 19711, USA.
- Schroeder RE, 1991. An Inhalation Developmental Toxicity Study in Rabbits with HCFC-124, Project No. 89-3514). Conducted by Bio/dynamics, Inc., East Millstone, NJ 08875-2360, USA, for the Program for Alternative Fluorocarbon Toxicity Testing (PAFT).
- STEP/AFEAS, 1993. Proceedings of the STEP-HALOCSIDE/AFEAS Workshop: Kinetics and Mechanism for the Reactions of Halogenated Organic Compounds in the Troposphere. Dublin, 23-25 March 1993.
- Tobeta Y, 1992. Test on biodegradability of HCFC 124 by microorganisms (closed bottle method). Final Report. Kurume Research Laboratories, Chemical Biotesting Center, Chemicals Inspection and Testing Institute. Japan.
- Trochimowicz HJ, Moore BL and Chiu T, 1977. Subacute Inhalation Toxicity Studies on Eight Fluorocarbons. Toxicol. Appl. Pharmacol. 41: 198-199.
- Tuazon EC and Atkinson R, 1993. Tropospheric Transformation Products of a Series of Hydrofluorocarbons and Hydrochlorofluorocarbons. J. Atmos. Chem., 17 (2):179-199.
- UKRGAR, 1990. Acid Deposition in the United Kingdom 1986-1988. Third Report of the United Kingdom Review Group on Acid Rain, September 1990.
- Van Poznak A and Artusio JF Jr, 1960. Anesthetic Properties of a Series of Fluorinated Compounds. 1. Fluorinated Hydrocarbons. Toxicol. Appl. Pharmacol. 2:363-373.
- Wada H, 1977. Anesthetic Activity and Fatality (F-123, 123a, 124 and 11). Attachment to Correspondence to M.B. Berenbaum, Allied Chemical Corporation, Morristown, NJ, USA, from H. Wada, Daikin Kogyo Co., Ltd., May 16, 1977.
- WMO, 1989. World Meteorological Association. Global Ozone Research and Monitoring Project, Report No. 20. Scientific Assessment of Stratospheric Ozone: 1989. Volume II, Appendix: AFEAS Report.
- WMO, 1991. World Meteorological Association. Global Ozone Research and Monitoring Project, Report No. 25. Scientific Assessment of Ozone Depletion.

MEMBERS OF THE TASK FORCE

D. Mayer	HOECHST D - Frankfurt am Main
M. Collins	ICI CHEMICALS AND POLYMERS GB - Cheshire
C. De Rooij	SOLVAY B - Brussels
J. Franklin	SOLVAY B - Brussels
S. Magda	SOLVAY PHARMA DEUTSCHLAND D - Hannover
G. Malinverno	AUSIMONT/MONTECATINI GROUP I - Milan
R. J. Millischer	ELF ATOCHEM F - Paris
G. M. Rusch	ALLIED SIGNAL Inc. USA - Morristown, NJ
H. J. Trochimowicz	E. I. DU PONT DE NEMOURS USA - Newark, DE
W. Haebler	ECETOC B - Brussels

MEMBERS OF THE SCIENTIFIC COMMITTEE (Peer Review Committee)

W. F. Tordoir (Chairman), Head of Occupational Health and Toxicology Division	SHELL NL - Den Haag
H. Verschuuren, (Vice-Chairman) Head of Toxicology Department	DOW EUROPE CH - Horgen
O. C. Bøckman, Scientific Advisor	NORSK HYDRO N - Porsgrunn
N. G. Carmichael, Toxicology Director Worldwide	RHÔNE-POULENC F - Lyon
H. De Henau, European Technical Centre Professional and Regulatory Services	PROCTER AND GAMBLE B - Brussels
A. De Morsier, Head, Ecotoxicology	CIBA-GEIGY CH - Basel
P. A. Gilbert, Head, Environment Division	UNILEVER UK - Port Sunlight
I. J. Graham-Bryce, Head of Environmental Affairs	SHELL NL - Den Haag
B. Hildebrand, Director, Experimental Toxicology	BASF D - Ludwischafen
J. R. Jackson, Director Medicine and Health Science	MONSANTO EUROPE B - Brussels
K. Künstler, Head Biological Research	HENKEL D - Düsseldorf
H. Lagast, chief Medical Officer	SOLVAY B - Brussels
E. Löser, Head of Institute of Industrial Toxicology	BAYER D - Wuppertal
R. Millischer ¹ , Chief Toxicologist	ELF ATOCHEM F - Paris
I. F. H. Purchase ¹ , Director, Central Toxicology Laboratory	ZENECA UK - Macclesfield

¹ Stewards responsible for primary peer review

TECHNICAL REPORTS

No. Title

- No. 1 Assessment of Data on the Effects of Formaldehyde on Humans. May 81
- No. 2 The Mutagenic and Carcinogenic Potential of Formaldehyde. May 81
- No. 3 Assessment of Test Methods for Photodegradation of Chemicals in the Environment. Aug 81
- No. 4 The Toxicology of Ethylene Glycol Monoalkyl Ethers and its Relevance to Man. Jul 82
- No. 5 Toxicity of Ethylene Oxide and its Relevance to Man. Sep 82
- No. 6 Formaldehyde Toxicology: an Up-Dating of the ECETOC Technical reports 1 and 2. Sep 82
- No. 7 Experimental Assessment of the Phototransformation of Chemicals in the Atmosphere. Sep 82
- No. 8 Biodegradation Testing: An Assessment of the Present Status. Nov 83
- No. 9 Assessment of Reverse-Phase Chromatographic Methods for Determining Partition Coefficients. Dec 83
- No. 10 Considerations Regarding the Extrapolation of Biological Data in Deriving Occupational Exposure Limits. Feb 84
- No. 11 Ethylene Oxide Toxicology and its Relevance to Man : An Up-Dating of ECETOC Technical Report n°5. Mar 84
- No. 12 The Phototransformation of Chemicals in Water : Results of a Ring-Test. Jun 84
- No. 13 The EEC 6th Amendment : A Guide to Risk Evaluation for Effects on the Environment. Mar 84
- No. 14 The EEC 6th Amendment : A Guide to Risk Evaluation for Effects on Human Health. Mar 94
- No. 15 The Use of Physical-Chemical Properties in the 6th Amendment and their Required Precision, Accuracy and Limiting Values. Jun 84
- No. 16 A review of Recent Literature on the Toxicology of Benzene. Dec 84
- No. 17 The Toxicology of Glycol Ethers and its Relevance to Man : An Up-Dating of ECETOC Technical Report n°4. Apr 85
- No. 18 Harmonisation of Ready Biodegradability Tests. Apr 85
- No. 19 An Assessment of Occurrence and Effects of Dialkyl-o-Phthalates in the Environment. May 85
- No. 20 Biodegradation Tests for Poorly-Soluble Compounds. Feb 86
- No. 21 Guide to the Classification of Carcinogens, Mutagens and Teratogens Under the 6th Amendment. Feb 86
- No. 22 Classification of Dangerous Substances and Pesticides in the EEC Directives. A Proposed Revision of Criteria for Inhalational Toxicity. 87
- No. 23 Evaluation of the Toxicity of Substances to be Assessed for Biodegradability. Nov 86
- No. 24 The EEC 6th Amendment : Prolonged Fish Toxicity Tests. Oct 86
- No. 25 Evaluation of Fish Tainting. Jan 87
- No. 26 The Assessment of Carcinogenic Hazard for Human Beings Exposed to Methylene Chloride. Jan 87
- No. 27 Nitrate and Drinking Water. Jan 88
- No. 28 Evaluation of Anaerobic Biodegradation. Jun 88
- No. 29 Concentrations of Industrial Organic Chemicals Measured in the Environment: The Influence of Physico- Chemical Properties, Tonnage and Use Pattern. Jun 88
- No. 30(5) Existing Chemicals : Literature Reviews and Evaluations. 94
- No. 31 The Mutagenicity and Carcinogenicity of Vinyl Chloride : A Historical Review and Assessment. Jul 88
- No. 32 Methylene Chloride (Dichloromethane) : Human Risk Assessment Using Experimental Animal Data. May 88
- No. 33 Nickel and Nickel Compounds : Review of Toxicology and Epidemiology with Special Reference to Carcinogenesis. Feb 89
- No. 34 Methylene Chloride (Dichloromethane) : An Overview of Experimental Work Investigating Species, Differences in Carcinogenicity and their Relevance to Man. Mar 89
- No. 35 Fate, Behaviour and Toxicity of Organic Chemicals Associated with Sediments. Jan 90
- No. 36 Biomonitoring of Industrial Effluents. Apr 90
- No. 37 Tetrachloroethylene : Assessment of Human Carcinogenic Hazard. May 90
- No. 38 A Guide to the Classification of Preparations Containing Carcinogens, Mutagens and Teratogens. Jul 90
- No. 39 Hazard Assessment of Floating Chemicals After an Accidental Spill at Sea. Jul 90
- No. 40 Hazard Assessment of Chemical Contaminants in Soil. Aug 90
- No. 41 Human Exposure to N-Nitrosamines, Their Effects and a Risk Assessment for n-Nitrosodiethanolamine in Personal Care Products. Aug 90
- No. 42 Critical Evaluation of Methods for the Determination of N-Nitrosamines in Personal Care and Household Products. Feb 91
- No. 43 Emergency Exposure Indices for Industrial Chemicals. Mar 91
- No. 44 Biodegradation Kinetics. Sep 91
- No. 45 Nickel, Cobalt and Chromium in Consumer Products: Allergic Contact Dermatitis. Mar 92
- No. 46 EC 7th Amendment: Role of Mammalian Toxicokinetic and Metabolic Studies in the Toxicological Assessment of Industrial Chemicals. May 92
- No. 47 EC 7th Amendment: 'Toxic to Reproduction' - Guidance on Classification. Aug 92
- No. 48 Eye Irritation: Reference Chemicals Data Bank. Aug 92
- No. 49 Exposure of Man to Dioxins: A Perspective on Industrial Waste Incineration. Sep 92
- No. 50 Estimating the Environmental Concentrations of Chemicals Using Fate and Exposure Models. Nov 92
- No. 51 Environmental Hazard Assessment of Substances. Jan 93
- No. 52 Styrene Toxicology Investigations on the Potential for Carcinogenicity. Nov 92
- No. 53 DHTDMAC: Aquatic and Terrestrial Hazard Assessment. CAS No. 61789-80-8. Feb 93
- No. 54 Assessment of the Biodegradation of Chemicals in the Marine Environment. Aug 93
- No. 55 Pulmonary Toxicity of Polyalkylene Glycols. (in preparation)
- No. 56 Aquatic Toxicity Data Evaluation. Dec 93
- No. 57 Polypropylene Production and Colorectal Cancer. Feb 94
- No. 58 Assessment of Non-Occupational Exposure to Chemicals. May 94
- No. 59 Testing For Worker Protection. May 94
- No. 60 Trichloroethylene: Assessment of Human Carcinogenic Hazard. May 94

Responsible Editor: D. A. Stringer, ECETOC
Av. E. Van Nieuwenhuyse, 4 (Bte 6)
B - 1160 Brussels, Belgium
D-1994-3001-106